



This patient group direction (PGD) must only be used by registered Paramedics or Nurse Practitioners who have been named and authorised by their organisation to practice under it. The most recent and up to date final signed version of the PGD should be used.

Patient group direction

for the administration of

Tranexamic Acid

by registered Paramedics / Advanced Nurse Practitioners

in Re-Pat Ltd

Version number:1.0

Change history


Version number	Change details	Date
1.1	Alteration to job titles on PGD development and authorisation page.	6/8/25

Document details

Reference number	2301776
Valid from	1/1/25
Renew date	31/12/2027
Expiry date	30/12/2027

PGD development

People responsible for PGD development

Name	Job title and organisation	Signature	Date
Richard Drakeford	Director		5/12/24
Dr James Cronin	Consultant Intensivist		
Neil Hayward	Pharmaceutical Consultant		
Jordan Cronshaw	Specialist Paramedic		

PGD authorisation

People responsible for PGD authorisation

Name	Job title and organisation	Signature	Date
Dr James Cronin	Consultant Intensivist		
Neil Hayward	Pharmaceutical Consultant		
Jordan Cronshaw	Specialist Paramedic		

Training and competency of registered Paramedics and Advanced Nurse Practitioners

Requirements of working under the PGD

Qualifications and professional registration	Current registration with the HCPC as a Paramedic. Current registration with the NMC as a Advanced Nurse Practitioner While working for Re-Pat Ltd Has successfully completed an approved university module in applied pharmacology, including individual applied pharmacology or other approved university modules that contain applied pharmacology as part of their content.
---	--

Initial training	Paramedics/Advanced Nurse Practitioners (ANP) must have undertaken appropriate training and successfully completed the competencies to undertake clinical assessment of patient leading to diagnosis of the conditions listed. Paramedics/ANP must be competent to recognise and treat unintended but expected side effects including loss of airway reflexes, respiratory depression and anaphylaxis.
Competency assessment	Paramedics / ANP should self-declare that they are competent to use this PGD, assuring themselves that they have the necessary clinical skills and knowledge for treatment of the conditions included and use of the drugs involved. Paramedics / ANP should also understand the legislation surrounding use of PGDs and their responsibilities as a PGD user.
Ongoing training and competency	The clinician must meet the requirements of the prevailing level of education required for PGD use at this level of practice. attending regular medicines management update training. All ongoing regular training requirements (e.g. statutory and mandatory training) as required by Re-Pat Ltd for this role must be completed. The clinician is responsible for keeping themselves aware of any changes to the recommendations for the medicine listed. It is the responsibility of the individual to keep up to date with continued professional development and to work within the limitations of their own individual scope of practice.

Clinical condition

Condition-specific information

Clinical condition or situation to which this PGD applies	Patients with signs of actual or suspected severe haemorrhage resulting from: • Major traumatic injuries, where significant internal or external haemorrhage (other than gastrointestinal haemorrhage) is known or suspected.
--	--

Inclusion criteria	Traumatic Haemorrhage Treatment of known or suspected traumatic internal or external haemorrhage (other than gastrointestinal haemorrhage), where bleeding started less than 3 hours ago. • In trauma: the clinician should assess the extent of traumatic haemorrhage using a combination of patient physiology, anatomical injury pattern, mechanism of injury and the patient's response to initial resuscitation. The above would include women who have recently given birth but have suffered subsequent trauma. • Children aged under 12 years old who have known or suspected severe haemorrhage from major traumatic injuries - if in doubt, these cases can be discussed with the duty senior on call clinician who can be contacted via the Critical Care Desk (CCD) in EOC. OR Post-Partum Haemorrhage (PPH) • PPH (bleeding from the genital tract >500ml) which usually occurs within 4 hours (but up to 24 hours) after delivery. This can be associated with haemodynamic instability. Tranexamic acid should be given after the administration of a uterotonic drug (misoprostol / syntometrine) except in individuals for whom uterotonic drugs are contraindicated (rare). • Woman with a post-partum haemorrhage when uterine trauma (rupture) is suspected. Bleeding may be intra-abdominal. Or Head Injuries • Tranexamic acid should be administered to any patient aged 18 years and over, who have a known or suspected head injury with a GCS of 12 or less, where the injury has occurred within the last 3 hours
----------------------------------	---

Exclusion criteria	Any haemorrhage condition not listed in the inclusion criteria Known allergy to tranexamic acid or any of the excipients (water for injection). Bleeding started more than 3 hours ago. Known or suspected gastrointestinal haemorrhage. Obvious resolution of haemorrhage. Post-partum haemorrhage before the administration of a uterotonic drug unless uterotonic drug is contraindicated or if trauma is the suspected cause. Critical interventions required (must only be given after critical interventions have been performed: i.e. airway managed; control or splinting of major haemorrhage etc. May be administered en route to hospital). Antepartum haemorrhage. Ensure exclusion is recorded in patient records.
---------------------------	--


Cautions (including any relevant action to be taken)	Where a caution is present the practitioner should be aware of the possible effects of administration but should continue to administer where the benefit outweighs risk. Patients with a history of convulsions. High dose regimes have been associated with convulsions; however, in the low dose regime recommended here, the benefit from giving tranexamic acid for severe haemorrhage outweighs the risk of convulsions. An increase in seizure rate may be due to the antagonistic effect of tranexamic acid on GABA receptors. Treat seizures as per JRCALC guidance. Patients with a history of acute venous or arterial thrombosis. In the low dose regime recommended here, the benefit from giving tranexamic acid for severe haemorrhage outweighs the risk of thrombotic events. This information should be passed to the receiving hospital. In the low dose regime recommended here, the benefit from giving tranexamic acid for severe haemorrhage outweighs the risk of thrombotic events. This information should be passed to the receiving hospital. In the low dose regime recommended here, the benefit from giving tranexamic acid for severe haemorrhage outweighs the risk of accumulation. This information should be passed to the receiving hospital. Patients receiving oral contraceptives may have an increased risk of thrombosis. Patients with severe renal impairment. There is a risk of accumulation of tranexamic acid. In Patients with massive haematuria. Avoid if risk of ureteric obstruction. Rapid injection may cause hypotension and loss of consciousness. Do not administer through the same line as blood products or penicillin antibiotics (including co-amoxiclav). Breast Feeding and Pregnancy: Tranexamic acid has been subject to some controlled clinical studies in pregnancy. Tranexamic acid does cross the placenta It's unknown what effect on the baby is. There is no evidence of teratogenicity in animal studies, however, only use if potential benefit outweighs the risk. Tranexamic acid's safe use during lactation has not been established. There is a small amount present in breast milk, although any antifibrinolytic effect in the infant is unlikely. The benefits of administration must outweigh any risk.
---	---

Arrangements for referral for medical advice	Patients who receive tranexamic acid should be fully monitored (ECG, NIBP and SpO2) and conveyed to the nearest appropriate hospital or handed over to an enhanced care team who are able to manage a patient who has had tranexamic acid administered. The receiving clinical team must be verbally informed, and the patient record should clearly show: • At what time the patient had tranexamic acid administered. • How much tranexamic acid the patient has had administered. • Whether the patient had capacity to provide informed consent.
Action to be taken if patient excluded	Ensure exclusion is recorded in patient records.
Action to be taken if patient declines treatment	Ensure the patient understands the information and rationale for the proposed administration and is therefore able to make an informed choice. It may be possible in the context of severe haemorrhage, that the patient will not be able to make an informed choice. Therefore, the clinician should act in the best interests of the patient at all times.

Details of the medicine

Medicines information

Name, form and strength of medicine	Tranexamic acid 500mg/5ml solution for injection ampoules
Legal category	POM
Indicate any off-label use (if relevant)	Tranexamic Acid's use in severe haemorrhage following trauma is currently off label. However, its use is supported by national and internationally recognised guidelines (4, 5), a large scale randomised controlled trial (6) the military (7) and other well established prehospital critical care systems. Its use in paediatric trauma is supported by national consensus statement (8) and recent papers (12). Tranexamic acid's use in post-partum haemorrhage is evidenced by a recent randomised, double-blind, placebo-controlled trial (9) and national obstetric guidance (10).

Route/method of administration	Intravenous injection (IV) Intraosseous injection (IO) *Intramuscular injection (IM) The IV or IO dose should be administered slowly over 10 minutes. *Where IV access is not achievable promptly, and the IO route is not appropriate, the IM route can be considered into a large muscle as below. Where two or more injections need to be administered, they should be given at separate sites, preferably in a different limb, they should be administered at least 2.5cm apart. IM Injection Sites and Appropriate Volumes <table border="1" data-bbox="472 488 1197 725"> <thead> <tr> <th>Site</th><th>Maximum volume in Adult Patients</th></tr> </thead> <tbody> <tr> <td>Deltoid</td><td>0.5 - 2 ml</td></tr> <tr> <td>Ventrogluteal</td><td>2.5 - 5 ml</td></tr> <tr> <td>Rectus Femoris</td><td>5ml</td></tr> <tr> <td>Vastus Lateralis</td><td>5ml</td></tr> </tbody> </table>  Injection Sites IM Needle Length <i>Vastus Lateralis and Rectus Femoris IM</i> The length of needle required is dependent upon the injection site chosen by the clinician. For intramuscular injections in infants, children and adults a 25mm (1 inch) long needle should be used. In larger adults or obese patients a 38mm (1.5 inch) long needle may be required. Intraosseous injection of tranexamic acid has been reported in a recent paper from the British Defence Medical Services with both military and civilian casualties receiving IO administration of tranexamic acid in 82 patients with no associated complications (11). Intravenous administration is preferable but if there is only IO access treatment should proceed without delay.	Site	Maximum volume in Adult Patients	Deltoid	0.5 - 2 ml	Ventrogluteal	2.5 - 5 ml	Rectus Femoris	5ml	Vastus Lateralis	5ml
Site	Maximum volume in Adult Patients										
Deltoid	0.5 - 2 ml										
Ventrogluteal	2.5 - 5 ml										
Rectus Femoris	5ml										
Vastus Lateralis	5ml										

Dose and frequency	Traumatic Haemorrhage Single dose administration only. In adults and children aged 12 years and over: • 1g (10ml) administered over 10 minutes when administering IV/IO. In children under 12 years of age: estimate the patient's weight (refer to JRCALC 'age page'): • 15mg/kg (maximum dose 1g (10ml)) administered over 10 minutes when administering IV/IO. These cases can be discussed, if advice is needed, with the duty senior on call clinician who can be contacted via the Critical Care Desk (CCD) in EOC. <i>Note:</i> The dose shown in this table is rounded to the nearest 50mg for ease of administration <table><tr><td>3yrs</td><td>14kg</td><td>200mg</td><td>2mls</td><td>100mg/ml</td></tr><tr><td>4yrs</td><td>16kg</td><td>250mg</td><td>2.5mls</td><td>100mg/ml</td></tr><tr><td>5yrs</td><td>19kg</td><td>300mg</td><td>3mls</td><td>100mg/ml</td></tr><tr><td>6yrs</td><td>21kg</td><td>300mg</td><td>3mls</td><td>100mg/ml</td></tr><tr><td>7yrs</td><td>23kg</td><td>350mg</td><td>3.5mls</td><td>100mg/ml</td></tr><tr><td>8yrs</td><td>26kg</td><td>400mg</td><td>4mls</td><td>100mg/ml</td></tr><tr><td>9yrs</td><td>29kg</td><td>450mg</td><td>4.5mls</td><td>100mg/ml</td></tr><tr><td>10yrs</td><td>32kg</td><td>500mg</td><td>5mls</td><td>100mg/ml</td></tr><tr><td>11yrs</td><td>35kg</td><td>500mg</td><td>5mls</td><td>100mg/ml</td></tr></table> Head Injury (adults aged 18 years and over only) Single dose administration only. • 1g (10ml) administered over 10 minutes when administering IV/IO. Post-partum Haemorrhage (in adults and children 12 years and over) • 1g (10ml) administered slowly when administering IV/IO. A second dose can be administered under this PGD for PPH if bleeding continues after 30 minutes of the first dose being administered. Any further doses are not permissible under this PGD within 24 hours.	3yrs	14kg	200mg	2mls	100mg/ml	4yrs	16kg	250mg	2.5mls	100mg/ml	5yrs	19kg	300mg	3mls	100mg/ml	6yrs	21kg	300mg	3mls	100mg/ml	7yrs	23kg	350mg	3.5mls	100mg/ml	8yrs	26kg	400mg	4mls	100mg/ml	9yrs	29kg	450mg	4.5mls	100mg/ml	10yrs	32kg	500mg	5mls	100mg/ml	11yrs	35kg	500mg	5mls	100mg/ml
3yrs	14kg	200mg	2mls	100mg/ml																																										
4yrs	16kg	250mg	2.5mls	100mg/ml																																										
5yrs	19kg	300mg	3mls	100mg/ml																																										
6yrs	21kg	300mg	3mls	100mg/ml																																										
7yrs	23kg	350mg	3.5mls	100mg/ml																																										
8yrs	26kg	400mg	4mls	100mg/ml																																										
9yrs	29kg	450mg	4.5mls	100mg/ml																																										
10yrs	32kg	500mg	5mls	100mg/ml																																										
11yrs	35kg	500mg	5mls	100mg/ml																																										

Administration notes	Visually inspect solution to ensure solution is clear and free from: Particulate matter prior to administration. Doses UNDER 500mg IV / IO / IM administration: Draw up the contents of ONE 500mg in 5ml ampoules into a labelled 5ml syringe (100mg/1ml). Doses OVER 500mg IV / IO administration: Draw up the contents of TWO 500mg in 5ml ampoules into a labelled 10ml syringe (100mg/1ml). IM administration: Draw up the contents of TWO 500mg in 5ml ampoule into TWO labelled 5ml syringes (100mg/1ml). TWO 5ml syringes are required to administer the total 1g dose across separate injection sites. Tranexamic acid is for administration by intravenous or intraosseous injection. Where IV access is not achievable promptly, and the IO route is not appropriate, the IM route can be considered into a large muscle. The IV/IO dose should be administered over 10 minutes. This can be given as 10 aliquots administered 1 minute apart where practicable. After administration flush with 0.9% Sodium Chloride. Do not administer through same line as blood products or penicillin antibiotics. All patients must be fully monitored including SpO2, ECG and BP. Rapid injection may cause hypotension and loss of consciousness. Resuscitation equipment should always be available
-----------------------------	---

Maximum or minimum treatment period	Traumatic Haemorrhage and Head Injuries Single dose permitted under this PGD. Post-partum haemorrhage A second dose can be administered under this PGD if bleeding continues after 30 minutes of the first dose being administered. Any further doses are not permissible under this PGD within 24 hours.
Adverse effects	Common side effects: • Nausea Vomiting Diarrhoea Uncommon: Dermatitis allergic Unknown frequency: Hypersensitivity reactions including anaphylaxis Convulsions - particularly in case of misuse Malaise Hypotension, with or without loss of consciousness (generally following a too fast intravenous injection, exceptionally after oral administration) Arterial or venous thrombosis at any sites Report adverse effects at: https://yellowcard.mhra.gov.uk/ Hypersensitivity reactions including anaphylaxis Convulsions - particularly in case of misuse Visual disturbances including impaired colour vision

Records to be kept	Record that valid, informed consent was given by the patient. In cases where the patient lacks mental capacity, a record of how mental capacity was assessed and how the administration of this medicine was in the best interest of the patient. Patient's name, address, date of birth Contact details of GP (if registered) Diagnosis or working diagnosis Patient's estimated weight when under 12 Dose given and route given by Batch number and expiry date of drugs given Time of administration Advice given to patient Signature and name of staff who administered medication Details of any adverse reactions and action taken
---------------------------	---

Patient information

Information for patients

Written information to be given to patient or carer	The patient record must show: • Which medications have been administered. • The strength of medications administered. • The dose of the medications administered. • The batch number of medications administered. • What time medications were administered. • The effect (intended or unintended) of medications administered The patient record must demonstrate that informed patient consent has been gained. If the patient lacks mental capacity a record of how the administration of the medicine is in the best interest of the patient. In cases where the patient has mental capacity: • Consent had been gained to administer medication. • The patient must be informed why treatment is required and what the intended effects of the medication are • Potential side-effects of the medication must be explained
--	--

Follow-up advice to be given to patient or carer

Patients who have received tranexamic acid should be conveyed to the nearest appropriate hospital or handed over to an enhanced care team who are able to manage a patient who has had tranexamic acid administered; OR in the cases of patients where life extinct is recognised patients may be left in the community in the normal way.

The receiving clinical team must be verbally informed, and the patient record should clearly show:

- Which medications have been administered.
- How much tranexamic acid the patient has had administered.
- Whether the patient had capacity to provide informed consent.

Appendices

Appendix A: key references

1. Tranexamic acid 100 mg/ml solution for injection - Summary of Product Characteristics (SmPC) - (emc) (medicines.org.uk)
Accessed online July 2023
2. British National Formulary by NICE Tranexamic Acid.
Accessed online March 2021

<https://bnf.nice.org.uk/drug/tranexamic-acid.html>

3. British National Formulary for Children by NICE Tranexamic Acid. Accessed online March 2021
<https://bnfc.nice.org.uk/drug/tranexamic-acid.html>
4. National Institute for Health and Care Excellence (2016) Major trauma: assessment and initial management, NICE guideline [NG39]
5. Spahn et al. The European guideline on management of major bleeding and coagulopathy following trauma. (2019) Critical care, 23(98)
6. Crash-2 Collaborators. (2011). The importance of preliminary treatment with tranexamic acid in bleeding trauma patients: an exploratory analysis of the CRASH-2 randomised controlled trial. The Lancet, 377(9771), 1096-1101.
7. Morrison, J. J., Dubose, J. J., Rasmussen, T. E., & Midwinter, M. J. (2012). Military application of tranexamic acid in trauma

emergency resuscitation (MATTERs) study. Archives of surgery, 147(2), 113-119.

8. Royal College of Paediatrics and Child Health (RCPCH). Evidence Statement Major trauma and the use of tranexamic acid in children November 2012

https://www.stemlynsblog.org/wp-content/uploads/2015/03/CEM6706-121112_TXA-evidence-statement_final11.pdf

9. Collaborators, W. T. (2017). Effect of early tranexamic acid administration on mortality, hysterectomy, and other morbidities in women with post-partum haemorrhage (WOMAN): an international, randomised, double-blind, placebo- controlled trial. The Lancet, 389(10084), 2105-2116.

10. Royal College of Obstetricians and Gynaecologists. Postpartum haemorrhage, prevention and management (Green Top Guideline No. 52) (2016)

11. Lewis, P., & Wright, C. (2014). Saving the critically injured trauma patient: a retrospective analysis of 1000 uses of intraosseous access. Emerg Med J, emermed- 2015.

12. Beno, S., Ackery, A. D., Callum, J., & Rizoli, S. (2014). Tranexamic acid in paediatric trauma: why not? Critical Care, 18(4), 313.

13. Brown, S. N., Kumar, D., Millins, M., & Mark, J. (Eds.). (2016). UK

ambulance services Clinical practice guidelines JRCALC 2016.

Bridgwater: Class Professional

14.Patient Group Directions. Medicines Practice Guide NICE August 2013. Updated March 2017

15.Updated WHO Recommendation on Tranexamic Acid for the Treatment of Postpartum Haemorrhage (2017). Accessed July 2023

16.Crash 3 Trial Collaborators (2019). Effects of tranexamic acid on death, disability, vascular occlusive events and other morbidities in patients with acute traumatic brain injury (CRASH-3): a randomised, placebo-controlled trial. Lancet; 394: 1713-1723

17.National PGD Tranexamic-acid-National-PGD-template-V2.1-FINAL-July-2023.doc (live.com) Last updated July 2023

Appendix B: Paramedics / Advanced Nurse Practitioners agreement to practise

I have read and understood the patient group direction and agree to administer this medicine only in accordance with this PGD.

Details of participating individual

Name	Signature	Senior authorising representative	Date