

# SAVE BIG ON ELANCO PRODUCTS

**INTERCEPTOR<sup>PLUS</sup>**  
(milbemycin oxime/praziquantel) OR **Interceptor**  
(milbemycin oxime)

**\$6** | **\$15**

OFF 6 DOSES | OFF 12 DOSES

OFFER CODE: INT2025

**Credelio**  
(lotilaner)

**\$15** | **\$35**

OFF 6 DOSES | OFF 12 DOSES

OFFER CODE: CRD2025

**Galliprant**  
(grapiprant tablets)

**\$20** | **\$60**

OFF 30-COUNT | OFF 90-COUNT

OFFER CODE: GPT2025

**Zenrelia**<sup>TM</sup>  
(ilunocitinib tablets)

**\$9** | **\$20**

OFF 30-COUNT | OFF 90-COUNT

OFFER CODE: ZEN2025

**Credelio Quattro**  
(lotilaner, moxidectin, praziquantel, and pyrantel chewable tablets)

**\$30** | **\$65**

OFF 6 DOSES | OFF 12 DOSES

OFFER CODE: CQ2025

**Credelio CAT**  
(lotilaner)

**\$10**

OFF 6 DOSES

OFFER CODE: CAT2025

**Seresto**

**\$6**

OFF 1 COLLAR

OFFER CODE: SER2025

**INTERCEPTOR<sup>PLUS</sup>**  
(milbemycin oxime/praziquantel) OR **Interceptor**  
(milbemycin oxime) OR **Credelio**  
(lotilaner)

**\$25** | **\$75**

OFF 6/6 DOSES | OFF 12/12 DOSES

OFFER CODE: INCR25

**Trifexis**  
(spinosad + milbemycin oxime)

**\$20**

OFF 6 DOSES

OFFER CODE: TFX2025

**Canine Parvovirus Monoclonal Antibody**  
USDA Conditionally Approved

**\$200**

OFF 1 TREATMENT

OFFER CODE: PARVO2025



For rebate details, scan the QR code or visit  
[ElancoRebates.com](https://ElancoRebates.com)

## INDICATIONS

Zenrelia is a prescription medication used to control itching and inflammation associated with skin allergies for dogs over 12 months of age.

## IMPORTANT SAFETY INFORMATION

See package insert including the Boxed Warning. For full prescribing information speak with

**WARNING: VACCINE-INDUCED DISEASE AND INADEQUATE IMMUNE RESPONSE TO VACCINES.** Based on results of the vaccine response study, dogs receiving Zenrelia are at risk of fatal vaccine-induced disease from modified live virus vaccines and inadequate immune response to any vaccine. Discontinue Zenrelia for at least 28 days to 3 months prior to vaccination and withhold Zenrelia for at least 28 days after vaccination. Dogs should be up to date on vaccinations prior to starting Zenrelia. Do not use in dogs less than 12 months old or dogs with a serious infection. Dogs should be monitored for the development of infections because Zenrelia may increase the chances of developing an infection. Neoplastic conditions (benign and malignant) were observed during clinical studies. The most common side effects were vomiting, diarrhea and tiredness. Zenrelia has not been tested in dogs used for breeding, pregnant, or lactating dogs and has not been evaluated in combination with glucocorticoids, cyclosporine, or other immune suppressive drugs.

## ASK US HOW YOU CAN SAVE ON THESE PRODUCTS!