

Dosing for Infants and Children

From Your Healthcare Professional



If possible, use weight to dose; otherwise use age.

mL = milliliter
tsp = teaspoon

DOSE: Every 4 hours as needed. DO NOT GIVE MORE THAN 5 DOSES IN 24 HOURS.



Infants' TYLENOL® Oral Suspension

Active Ingredient:
Acetaminophen 160 mg
(in each 5 mL)

Available in:



Use only as directed.



Children's TYLENOL® Oral Suspension

Active Ingredient:
Acetaminophen 160 mg
(in each 5 mL or 1 tsp)

Available in:



Use only as directed.

WEIGHT	AGE			
6-11 lbs	0-3 mos	1.25 mL		—
12-17 lbs	4-11 mos	2.5 mL		—
18-23 lbs	12-23 mos	3.75 mL		—
24-35 lbs	2-3 yrs	5 mL		5 mL (1 tsp)
36-47 lbs	4-5 yrs	—		7.5 mL (1½ tsp)
48-59 lbs	6-8 yrs	—		10 mL (2 tsp)
60-71 lbs	9-10 yrs	—		12.5 mL (2½ tsp)
72-95 lbs	11 yrs	—		15 mL (3 tsp)

IMPORTANT INSTRUCTIONS for Proper Use

Today's Date: _____

This dosing recommendation from your doctor will expire in 14 DAYS.

- Read and follow the label on all TYLENOL® products.
- Take every 4 hours as needed. Do NOT exceed more than 5 doses in 24 hours.
- Do NOT use with any other product containing acetaminophen.
- Use only the dosing device that comes with a specific product.
- **All Infants' TYLENOL® and Children's TYLENOL® Oral Suspension products have the same acetaminophen concentration (160 mg/5 mL).**