

August 1 - October 31, 2026

Save now on select Elanco solutions.

Elanco™

Category	Qualifying Products	Offer Details
Cydectin®	Cydectin® (moxidectin) Pour-On 1L, 2.5L, 5L, and 10L	Receive a \$7/liter rebate, no minimum liter purchase and flat rebate value assigned per SKU.
	Cydectin® (moxidectin) Oral Drench 1L and 4L	
	Cydectin® (moxidectin) Injectable	Receive \$25/unit (2 unit minimum)
Insecticides	Clean-Up™ II Pour-On Insecticide with IGR	Earn a \$25 rebate on Elanco insecticides for every \$500 spent on qualifying products.
	CyLence® Pour-On Insecticide	
	CyLence Ultra® Insecticide Cattle Ear Tag	
	Permethrin® CDS Pour-On Insecticide	
	Agita® 10 WG	
Ear Tags	Corathon®	2% rebate: 240-4,999 tags 4% rebate: 5000 and up tags
	CyLence Ultra® Cattle Ear Tag	
	Patriot™ Insecticide Cattle Ear Tag	
Injectables	Baytril® 100 (enrofloxacin)	Baytril® 250ml: \$30/bottle Baytril® 500ml: \$60/bottle
	Pradalex™ (pradofloxacin injection)	Pradalex™ 100ml: \$10/bottle Pradalex™ 250ml: \$25/bottle
	Micotil® 300 (tilmicosin injection) Prior to using this drug, it is important to read the entire package insert including the boxed warning.	Micotil®: \$65/bottle
	Increxxa™ (tulathromycin injection)	Increxxa™ 100ml: \$30/bottle Increxxa™ 250ml: \$75/bottle Increxxa™ 500ml: \$140/bottle
Vaccines	Titanium® T3, T5 and T5L5	10 dose: \$3/bottle rebate 50 dose: \$5/bottle rebate
	NUPLURA® PH and NUPLURA® PH+5	50 dose: \$15/bottle rebate
	Scour Bos®	10 dose: \$3/bottle rebate 50 dose: \$15/bottle rebate
	Vira Shield® 6 + VL5 Somnus, 6 + Somnus, and 6 + VL5 HB	10 dose: \$3/bottle rebate 50 dose: \$20/bottle rebate
	ReproSTAR® L5 HB	
	Vib Shield® Plus L5	



Claim your rebate!
ElancoCattleRebates.com

Minimum \$50
rebate required
across program

BAYTRIL 100 INDICATIONS

Single-Dose Therapy: Baytril 100 is indicated for the treatment of bovine respiratory disease (BRD) associated with *Mannheimia haemolytica*, *Pasteurella multocida*, *Histophilus somni* and *Mycoplasma bovis* in beef and non-lactating dairy cattle; and for the control of BRD in beef and non-lactating dairy cattle at high risk of developing BRD associated with *M. haemolytica*, *P. multocida*, *H. somni* and *M. bovis*.

Multiple-Day Therapy: Baytril 100 is indicated for the treatment of BRD associated with *M. haemolytica*, *P. multocida* and *H. somni* in beef and non-lactating dairy cattle.

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian and prohibits the extra-label use of this drug in food-producing animals.

Not for human use. Keep out of reach of children. Avoid contact with eyes and skin. Individuals with a history of hypersensitivity to quinolones should avoid this product. Not for use in female dairy cattle 20 months of age or older, including dry dairy cows, and veal calves; a withdrawal period has not been established for this product in pre-ruminating calves. Quinolone-class drugs should be used with caution in animals with known or suspected Central Nervous System (CNS) disorders. Baytril 100 has a pre-slaughter withdrawal time of 28 days for cattle. See [package insert](#) for additional safety information.

PRADALEX INDICATIONS

Pradalex is indicated for the treatment of bovine respiratory disease (BRD) associated with *Mannheimia haemolytica*, *Pasteurella multocida*, *Histophilus somni* and *Mycoplasma bovis* in cattle intended for slaughter (beef calves 2 months of age and older, growing beef steers, growing beef heifers, and beef bulls intended for slaughter), and in cattle intended for breeding less than 1 year of age (replacement beef and dairy heifers less than 1 year of age and beef and dairy bulls less than 1 year of age). Not for use in cattle intended for breeding 1 year of age and older (replacement beef and dairy heifers 1 year of age and older, beef and dairy bulls 1 year of age and older, and beef and dairy cows), beef calves less than 2 months of age, dairy calves, and veal calves.

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian and prohibits the extra-label use of this drug in food-producing animals. To ensure responsible antimicrobial drug use, use of pradofloxacin should be limited to treatment of bovine respiratory disease (BRD) in cattle only after consideration of other non-fluoroquinolone therapeutic options.

Not for use in humans. Keep out of reach of children. Avoid contact with eyes and skin. Individuals with a history of hypersensitivity to quinolones should avoid this product. Not for use in animals intended for breeding greater than 1 year of age. Not for use in beef calves less than 2 months of age, dairy calves and veal calves; a withdrawal period has not been established for this product in pre-ruminating calves. Quinolones should be used with caution in animals with known or suspected central nervous system (CNS) disorders. Mild to moderate inflammatory changes of the injection site may be seen in cattle treated with Pradalex. Pradalex has a pre-slaughter withdrawal time of 4 days for cattle. See [package insert](#) for additional safety information.

MICOTIL 300 INDICATIONS

Micotil is indicated for the treatment of bovine respiratory disease (BRD) associated with *Mannheimia haemolytica*, *Pasteurella multocida* and *Histophilus somni*. It is indicated for the control of respiratory disease in cattle at high risk of developing BRD associated with *M. haemolytica*.

IMPORTANT SAFETY INFORMATION FOR MICOTIL

Before using this product, it is important to read the entire product insert, including the boxed human warning. For full prescribing information call 1 888 545 5973 or visit www.elancolabels.com/us/micotil-300.

HUMAN WARNING: Not for human use. Injection of this drug in humans has been associated with fatalities. Keep out of reach of children. Administer only with a tube-fed safety syringe. Do not use in automatically powered syringes, single-use syringes, or other delivery devices. Exercise extreme caution to avoid accidental self-injection. In case of human injection, consult a physician immediately and apply ice or cold pack to injection site while avoiding direct contact with the skin. Emergency medical telephone numbers are 1-800-722-0987 or 1-800-428-4441. Avoid contact with skin, eyes, or mucous membranes.

Micotil is available only by prescription. Micotil must be used with the quick-fit connector and tube-fed safety syringe included to avoid accidental self-injection. Contact Elanco or your distributor for this equipment if needed. For subcutaneous injection in cattle and sheep only; do not administer intravenously. Injection of this antibiotic has been shown to be fatal in swine and non-human primates and may be fatal in horses, goats, and other species. Not for use in use in lactating dairy cattle 20 months of age or older. Use in lactating dairy cattle or sheep may cause milk residues. Injection site swelling and inflammation, lameness, collapse, anaphylaxis/anaphylactoid reactions, decreased food and water consumption, and death have been reported in cattle. Micotil has a pre-slaughter withdrawal time of 42 days. See [package insert](#) for full prescribing information.

INCREXXA INDICATIONS

Beef and Non-Lactating Dairy Cattle: Increxxa is indicated for the treatment of bovine respiratory disease (BRD) and control of respiratory disease in cattle at high risk of developing BRD associated with *Mannheimia haemolytica*, *Pasteurella multocida*, *Histophilus somni* and *Mycoplasma bovis*; for the treatment of infectious bovine keratoconjunctivitis (IBK) associated with *Moraxella bovis*; and for the treatment of bovine foot rot (interdigital necrobacillosis) associated with *Fusobacterium necrophorum* and *Porphyromonas levis*.

Suckling Calves, Dairy Calves and Veal Calves: Increxxa is indicated for the treatment of BRD associated with *M. haemolytica*, *P. multocida*, *H. somni*, and *M. bovis*.

Important Safety Information (ISI)

Increxxa is available only by prescription. Not for human use. Keep out of reach of children. Do not use in animals previously found to be hypersensitive to the drug. Not for use in chickens or turkeys. Do not use in female dairy cattle 20 months of age or older, including dry dairy cows. Injection site reactions and anaphylaxis/anaphylactoid reactions have been reported in cattle. Increxxa has a pre-slaughter withdrawal time of 18 days for cattle. See [package insert](#) for full prescribing information.

Keep Cydectin® out of reach of children.

The label contains complete use information, including cautions and warnings. Always read, understand, and follow the label directions.

Baytril is sold by Elanco or its affiliates and are not Bayer products. The Baytril trademark is owned by Bayer and used under license.

Agita, Clean-Up, Corathon, Cydectin, CyLence, CyLence Ultra, Increxxa, Micotil, NUPLURA, Patriot, Permethrin, Pradalex, ReproSTAR, Scour Bos, Titanium, Vib Shield, Vira Shield, Elanco and the diagonal bar logo are trademarks of Elanco or its affiliates. ©2026 Elanco or its affiliates. PM-US-26-1247

Program Terms & Conditions:

Ways to claim your rebate:

1. Online Claims: Scan the QR code to be directed to the rebate account set up form to create a rebate account and submit your rebate claims, or visit ElancoCattleRebates.com
2. Mail-in Claims: Complete a mail-in claim form and submit with proof of purchase (invoice) via mail.
3. Supplier Reported Claims (EDI): Qualifying purchases billed through an EDI-reporting distributor will be aggregated at the end of the program. EDI-reporting distributor must report EDI sales to Elanco. Qualifying customers will automatically receive a rebate check in the mail.

Program Terms:

- Beef and dairy producers only are eligible for this program. Veterinarians, resellers, and swine producers are ineligible. Producers enrolled in any special Elanco pricing programs or purchases below Elanco's minimum advertised price are not eligible for this rebate offer.
- Purchases are cumulative August 1 - October 31, 2026. Eligible purchases must be invoiced by November 30, 2026.
- \$50 minimum rebate check.
- EDI Purchases over \$500 will be paid automatically if the check is a minimum of \$50.
- Any EDI purchases under \$500 during the program period will not be automatically paid and require submission of an online or mail in claim to receive a rebate if the \$50 minimum is hit.
- Allow at minimum 6 weeks for processing following program conclusion.
- Eligible products for the offer include: Agita®, Baytril® 100, Clean-Up II™ Pour-On Insecticide, Cydectin® Oral Drench, Cydectin® Injectable, Cydectin® Pour-On, CyLence® Pour-On Insecticide, CyLence Ultra® Insecticide, CyLence Ultra® Insecticide Ear Tag, Corathon®, CyLence Ultra®, Increxxa™, Micotil®, NUPLURA® PH, NUPLURA® PH+5, Patriot™, Permethrin® CDS Pour-On Insecticide, Pradalex™, ReproSTAR® L5 HB, Scour Bos®, Titanium®, Vib Shield®, Vira Shield® 6+ Somnus, Vira Shield® 6 + VL5 HB, Vira Shield® 6 + VL5 Somnus. Offers valid while supplies last.
- Products that are on backorder and qualify for the rebate offer must ship during the program term to receive the rebate. Products that are on backorder and do not ship during the program term will not qualify for any rebate in this program. All product returns will be applied toward the subsequent program payout period.
- Producers who submit coupons via mail or online will not be eligible for EDI reported claims.
- Limit 10 mail or online claim submissions per household. Multiple proofs of purchase can be used per submission. One rebate will be calculated per submission. Qualifying rebates will be calculated for all invoices included in a submission and will not be calculated across multiple submissions.
- Mail and online rebate claim form and invoice(s) must be postmarked or submitted by November 30, 2026.
- Elanco reserves the right to vary the terms and conditions of this program or to cancel this program at any time upon notice through the website Elanco.com.

For more information regarding this rebate program and eligibility, contact your Elanco Animal Health representative or email us at ElancoConsumerRebates@apfco.com

Elanco

TM