Guide to the Science of Climate Change in the 21st Century

Chapter 12 Climate Change – Natural Forces

> David H. Manz PhD., P. Eng., AOE, FCAE 2020



Eco-print, "Inspiration", Nora Manz 2020

Table of Contents

Chapter 12	Climate Change – Natural Forces
	12.1 Introduction
	12.2 Tectonic activity
	12.3 Solar radiation, sun spots and cosmic radiation
	12.4 Milankovitch Cycles – changes in Earth's orbit and
	orientation to the sun
	12.5 Volcanic activity
	12.6 Global dimming
	12.7 Information support

List of Figures

Figure 12.1 Earth climate system.

Figure 12.2 A segment of Earth from its center, the core, to its outer surface consisting of the lithosphere and the crust.

Figure 12.3 Note the Earth's crust consists of the ocean crust and the much thicker continental crust.

Figure 12.4 Plates as they exist today.

Figure 12.5 Earth supercontinent, Pangaea, 240 million years or more.

Figure 12.6 Continental drift.

Figure 12.6 Thrust and reverse fault movement resulting in mountain formation.

Figure 12.7 Subduction.

Figure 12.8 Movement of magma within the Earth's crust.

Figure 12.9 Ring of fire. Volcanic activity on the Pacific Rim.

Figure 12.10 Solar cycle variations – daily and annual.

Figure 12.11 Sunspots on surface of the sun.

Figure 12.12 Difference between a solar maximum and a solar minimum.

Figure 12.13 Variation of Earth's orbit around the Sun – eccentricity.

Figure 12.14 Variation of the tilt of the Earth's axis of rotation – axial tilt or obliquity.

Figure 12.15 Variation of Earth's rotational axis – precession.

Figure 12.16 Rotation of Earth's orbit around the sun – apsidal precession.

Figure 12.17 Back and forth tilt of the plane of Earth's orbit around the Sun relative to a reference plane – orbital oscillation.

Figure 12.18 Berger astronomical model of orbital variability, past, present and future.

Figure 12.19 Volcanic eruption releasing ash and a variety of aerosols and gases.

Figure 12.20 Mt. Pinatubo and global circulation of ash.

Figure 12.21 Volcanic eruptions since 1870.

Figure 12.22 Sun-blocking aerosols around the world since the 1991 eruption of Mount Pinatubo according to satellite estimates. NASA

Figure 12.23 Contrails 2001.

Figure 12.24 Forest fires – eastern China.

Figure 12.25 Forest fires 2020 – western United States, NASA.

Figure 12.26 Pollution - Golden Gate Bridge – San Francisco.

Figure 12.27 Dust storm sweeping across the Middle East – NASA.

12.0 Climate Change – Natural Forces

12.1 Introduction

Earth's climate is constantly being *forced* to change by a variety of natural occurring phenomena that directly or indirectly affect the energy budget.

Initial response may result in amplification of the original forcing phenomena, an effect known as *positive feedback*. The resulting impact on climate change may be much more significant than that which would have resulted were the original forcing acting without positive feedback.

Natural occurring phenomena affecting Earth's climate include:

- Tectonic processes resulting in movement of continents.
- Increase or decrease in intensity of solar radiation from the Sun.
- Changes in Earth's orbit and orientation to the Sun.
- Volcanic activity.

These interact with Earth's climate system as shown in Figure 12.1.



Figure 12.1 Earth climate system.

12.2 Tectonic activity

Tectonic processes that influence climate system variability include plate motion, changes in continental geography, mountain building and erosion, the production and subduction of seafloor crust, and related changes in bio-geo-chemical cycles, particularly the carbon cycle.

Figure 12.2 shows a segment of Earth from its center, the core, to its outer surface, the lithosphere and crust. The lighter crust floats on the lithosphere. As shown in Figure 12.3 there are two types of crust, the ocean crust and the continental crust which is much thicker.



Figure 12.2 A segment of Earth from its center, the core, to its outer surface consisting of the lithosphere and the crust.



Figure 12.3 Note the Earth's crust consists of the ocean crust and the much thicker continental crust.

The Earth's crust and lithosphere are divided into portions known as plates that move more-orless independently of each other. The crust and lithosphere below it move together. Figure 12.4 illustrates the plates as they exist today. Note that a plate might include both ocean and continental types of crust.

Approximately 240 million years BP the plates were orientated to form a single continental mass or continent known as Pangaea as shown in Figure 12.5 and the top illustration in Figure 12.6. As a result of plate movement, (known as tectonic drift), the continental masses separated to ultimately form present day continents. The plates continue to move. The Himalayas continue to be forced upwards – Mount Everest gains four mm per year.



Figure 12.4 Plates as they exist today.



Figure 12.5 Earth supercontinent, Pangaea, 240 million years or more.

HOW DO WE KNOW THE CONTINENTS MOVED?

In the early 20th century, German researcher Alfred Wegener noted that fossil remains of the same extinct animal or plant can be found across several continents that are not now adjacent to one another. This suggests that in the distant past, the configuration of continents was different than it is today, a theory Wegener called "continental drift."



Figure 12.6 Continental drift.

In the process of moving the various plates collide with each other. One of the plates may be thrust vertically to form mountains as shown in Figure 12.6. The subducting process may also result in volcanic activity as illustrated in Figure 12.7. As the subducting ocean crust and associated lithosphere sink the result may be local uplift of the resisting plate and/ or the lighter subducting crust may melt. The mixed lighter molten rock, magma, moves upward, breaks through the overlying lithosphere and crust to spill out over the surface to form a volcano. The occurrence of this process around the Pacific Ocean is illustrated in Figure 12.8 and is known as the 'Ring of Fire' named after the volcanoes that are formed as a consequence plate collision.

Volcanos may form from plumes of magma flowing directly from the mantle. This is shown in Figure 12.7 and 12.8. Volcanoes resulting from plate movement around the Pacific are shown in Figure 12.9 where many of largest eruptions in the past 10,000 years have occurred.



Figure 12.6 Thrust and reverse fault movement resulting in mountain formation.



Figure 12.7 Subduction.



Figure 12.8 Movement of magma within the Earth's crust.



Figure 12.9 Ring of fire. Volcanic activity on the Pacific Rim.

Another phenomenon is related to the melting of the Greenland and Antarctic ice sheets. As they formed the weight of the ice caused the underlying land mass to sink into the crust. As they melt and the water drains into the surrounding ocean the underlying land masses rise in a process known as isostatic rebound (also called continental rebound, post-glacial rebound or isostatic adjustment). Land masses once covered by kilometers thick layers of ice during the last ice age are still rebounding even though the ice essentially disappeared twelve thousand years ago.

Tectonic processes impact ocean currents, atmospheric circulation, and volcanic activity. Ocean circulation will be affected by the spatial relationships of the continents to each other but the changes can be expected to be very gradual occurring over many millions of years similar to mountain building processes that will affect atmospheric circulation. By comparison volcanic activity occurs continuously and can have immediate effect and for this reason is treated separately.

12.3 Solar radiation, sunspots and cosmic radiation

Solar radiation reaching the Earth has been discussed quite extensively in Chapter 4. Solar irradiance is the power per unit area, Watts per square metre (Wm^{-2}) of solar radiant energy reaching the Earth as observed at the top of the atmosphere or earth's surface. The measure of solar irradiance received over a period of time of energy per unit area, Joules per square metre (Jm^{-2}) , is known as solar radiant energy per square metre, solar irradiation, solar insolation or insolation.

Solar irradiance varies over a ten to twelve-year period. Each period is called a solar cycle. The variations in strength of solar irradiance over each cycle are in the order of 0.05% to 0.1%. These small variations in solar irradiance, on a daily and annual basis, are shown in Figure 12.9. In the absence of anthropogenic forcing these variations have been shown to be significant.

The solar cycle is observed by variations in the Sun's magnetic activity and on the surface of the Sun by the presence of sunspots and solar flares. Sunspots are colder areas of the Sun's surface and are observed as shown in Figure 12.10 and are of particular interest because of possible association with forcing of climate change. Solar flares are a flash of increased brightness on the Sun that may result in the ejection of coronal mass that interact with Earth's magnetic field – often observed as Northern Lights but have no apparent affect on climate. The Sun's magnetic activity protects Earth from cosmic radiation originating from outside the solar system. Fluctuations in the Sun's magnetic field inversely affect the amount of cosmic radiation reaching the Earth – the greater the activity and stronger the magnetic field, the less cosmic radiation reaches the Earth.



Figure 12.10 Solar cycle variations – daily and annual.

At the beginning, midway and end of a solar cycle there are the least number of sunspots, known as a solar minimum (least amount of solar activity) and one- quarter and three-quarter way into the cycle there are the most sunspots, known as the solar maximums (most amount of solar activity). Figure 12.11 illustrated sun spots on the surface of the Sun compared to the size of the Earth. Figure 12.12 illustrates the difference between a solar maximum and a solar minimum.

Because the occurrence and number of sunspots are directly correlated to the solar cycle and the Sun's magnetic field their presence can be correlated to amount of cosmic radiation reaching the Earth – in particular its atmosphere. It had been hypothesized that because cosmic radiation can induce aerosol formation that this must lead to increased cloud formation and global cooling. The sequence of steps were the production of aerosols, growth of aerosols, and the production of water particles that result in cloud formation and reflection of solar radiation. The European Organization for Nuclear Research, CERN, has demonstrated that cosmic radiation can induce aerosol formation. However, several investigations have disproved the hypothesis that aerosol production would lead to significant increases in cloud cover that would result in global cooling.



Figure 12.11 Sunspots on surface of the sun.





12.4 Milankovitch Cycles – changes in Earth's orbit and orientation to the Sun.

Perhaps the most important of the natural forces affecting Earth's climate are the Milankovitch Cycles named after the Serbian geophysicist and astronomer Milutin Milankovitch. He hypothesized that variations in Earth's orbit known as eccentricity, the axial tilt of the Earth known as obliquity, and rotation of the axial tilt known as precession resulted in cyclical variation in how solar radiation was received by the Earth and that these variations could be used to explain the periodic formation and disappearance of the ice ages. Figures 12.13, 12.14 and 12.15 illustrate eccentricity, axial tilt and precession. Note that were a few other investigators who had made similar hypothesis but were unable to demonstrate the validity of their ideas because of the lack evidence of the occurrence and timing of ice age phenomena. This information only became available toward the end of the nineteenth century.



Figure 12.13 Variation of Earth's orbit around the Sun – eccentricity.

Figure 12.14 Variation of the tilt of the Earth's axis of rotation – axial tilt or obliquity.

Figure 12.15 Variation of Earth's rotational axis – precession.

There are two other cycles, rotation of Earth's orbit around the Sun in the same plane known as apsidal precession and the movement of the inclination of Earth's orbit around the sun, up and down, known as orbital inclination. These are shown in Figures 12.16 and 12.17.

Figure 12.18 shows variations in Earth's orbit and the resulting changes in solar energy flux at high latitude.



Figure 12.16 Rotation of Earth's orbit around the sun – apsidal precession.



Figure 12.17 Back and forth tilt of the plane of Earth's orbit around the Sun relative to a reference plane – orbital oscillation.



Figure 12.18 Berger astronomical model of orbital variability, past, present and future.

12.5 Volcanic activity

Volcanic activity occurs as a result of tectonic processes as discussed in 12.1. It frequently results in the ejection of molten rock, ash and gas from below the Earth's crust as illustrated in Figure 12.19. The impact may be local or global as did the eruption of Mt. Pinatubo in the Philippines shown in Figure 12.20. NASA estimates that the maximum effect of the five largest volcanoes over the past century is less than 0.2 degrees cooling globally and 0.4 degrees regionally. Volcanoes produce approximately 1% of the carbon dioxide produced by human activity. Aerosols may persist up to two years or longer in the northern hemisphere which can result in greater areas of snow and ice cover and regional cooling. The stratosphere may warm several degrees which in turn may strengthen the polar vortex in the northern hemisphere and the polar jet stream resulting in warmer winters. The ozone layer may also be depleted because of the presence of chlorine and other CFCs similar to those produced by human activity.



Figure 12.19 Volcanic eruption releasing ash and a variety of aerosols and gases.



Figure 12.20 Mt. Pinatubo and global circulation of ash.

Figure 12.21 illustrates the significant volcanic activity since 1870. The greater the optical depth the more aerosols ejected into the atmosphere. Figure 12.22 shows the most recent volcanic eruptions and how their effects have dissipated with time.



Figure 12.21 Volcanic eruptions since 1870.



Figure 12.22 Sun-blocking aerosols around the world since the 1991 eruption of Mount Pinatubo according to satellite estimates. NASA

12.6 Global dimming

Global dimming is caused by forest fire, pollution, contrails and dust storms. The effect is the same as volcanic activity as described 12.4. Global dimming caused by contrails as shown in Figure 12.23 be more significant than the GHG emissions from the jet plane engines. Global dimming due to forest fires is shown in Figures 12.24 and 12.25. Global dimming due to pollution is shown in Figure 12.26. The effect of dust storms is shown in Figure 12.27.



Figure 12.23 Contrails 2001.



Figure 12.24 Forest fires – eastern China.



NASA's Aqua satellite captured this true-color image of the United States on Sep. 15, 2020, showing the fires in the West, the smoke from those fires drifting over the country, several hurricanes converging from different angles, and Hurricane Sally making landfall. Credit: NASA

Figure12.25 Forest fires 2020 – western United States, NASA.



Figure 12.26 Pollution - Golden Gate Bridge – San Francisco.



Figure 12.27 Dust storm sweeping across the Middle East – NASA.

12.7 Information support

Key web sites:

- 1. Solar variability and the total solar irradiance. https://archive.ipcc.ch/publications_and_data/ar4/wg1/en/ch1s1-4-3.html
- 2. Sea level change. https://www.ipcc.ch/site/assets/uploads/2018/02/WG1AR5_Chapter13_FINAL.pdf
- 3. Anthropogenic and natural radiative forcing. https://www.ipcc.ch/site/assets/uploads/2018/02/WG1AR5_Chapter08_FINAL.pdf
- 4. Solar irradiance. https://en.wikipedia.org/wiki/Solar_irradiance
- 5. Insolation. <u>https://www.sciencedirect.com/topics/earth-and-planetary-sciences/insolation</u>
- 6. Solar cycle. <u>https://en.wikipedia.org/wiki/Solar_cycle</u>
- 7. Solar cycle. <u>https://spaceplace.nasa.gov/solar-cycles/en/</u>
- Sunspot cycle. <u>https://scied.ucar.edu/sunspot-</u> cycle#:~:text=The%2011%2Dyear%20sunspot%20cycle,northern%20and%20south ern%20hemispheres%20switch.
- 9. Cosmic radiation and solar activity from ice cores and tree rings. https://www.pnas.org/content/109/16/5967
- 10. Cosmic rays and global warming. <u>https://skepticalscience.com/cosmic-rays-and-global-warming-advanced.htm</u>
- 11. Milankovitch cycles. https://en.wikipedia.org/wiki/Milankovitch cycles
- 12. Milankovitch cycles. https://www.skepticalscience.com/Milankovitch.html
- 13. Astronomical theory of climate change. <u>https://www.researchgate.net/publication/238956033</u> Astronomical theory of cl <u>imate_change</u>
- 14. Volcanoes, explained. <u>https://www.nationalgeographic.com/environment/natural-disasters/volcanoes/#close</u>

- 15. Ring of fire. https://en.wikipedia.org/wiki/Ring of Fire
- 16. Ring of fire. https://www.britannica.com/place/Ring-of-Fire
- 17. Tectonic shift. https://oceanservice.noaa.gov/facts/tectonics.html
- 18. Plate tectonics. <u>https://volcanoes.usgs.gov/vsc/file_mngr/file-139/This_Dynamic_Planet-Teaching_Companion_Packet.pdf</u>
- 19. Earth's crust.

https://www.nationalgeographic.org/encyclopedia/crust/#:~:text=1%2F12-,%E2%80%9CCrust%E2%80%9D%20describes%20the%20outermost%20shell%20of %20a%20terrestrial%20planet.,of%20solid%20rocks%20and%20minerals.

- 20. Eruption of Mount Pinatubo, Philippines. <u>https://pubs.usgs.gov/fs/1997/fs113-97/</u>
- 21. Global effects of Mount Pinatubo. https://earthobservatory.nasa.gov/images/1510/global-effects-of-mount-pinatubo
- 22. How volcanoes influence climate. <u>https://scied.ucar.edu/shortcontent/how-volcanoes-influence-climate</u>
- 23. Volcano hazards. <u>https://www.usgs.gov/natural-hazards/volcano-hazards/volcanic-gases</u>
- 24. Anthropogenic and natural radiative forcing. <u>http://www.climatechange2013.org/images/report/WG1AR5_Chapter08_FINAL.p</u> <u>df</u>
- 25. Volcanoes and climate change. <u>https://earthdata.nasa.gov/learn/sensing-our-planet/volcanoes-and-climate-change</u>
- 26. Volcanic eruptions and climate. http://climate.envsci.rutgers.edu/pdf/ROG2000.pdf
- 27. Volcanoes. <u>https://archive.ipcc.ch/publications_and_data/ar4/wg1/en/ch8s8-7-2-3.html</u>
- 28. Changes in atmospheric constituents and in radiative forcing. https://www.ipcc.ch/site/assets/uploads/2018/02/ar4-wg1-chapter2-1.pdf
- 29. Explosive volcanic activity. https://archive.ipcc.ch/publications_and_data/ar4/wg1/en/ch2s2-7-2.html

- 30. Role of volcanism and solar irradiance. https://archive.ipcc.ch/publications_and_data/ar4/wg1/en/ch9s9-3-3-2.html
- 31. How much do volcanoes influence the climate? <u>https://www.carbonbrief.org/in-brief-how-much-do-volcanoes-influence-the-climate</u>
- 32. Global dimming. https://en.wikipedia.org/wiki/Global dimming
- 33. Contrails. <u>https://www.sciencemag.org/news/2019/06/aviation-s-dirty-secret-airplane-contrails-are-surprisingly-potent-cause-global-warming</u>
- 34. Dust storm sweeps across Middle East. <u>https://earthobservatory.nasa.gov/images/86571/dust-storm-sweeps-across-</u> <u>middle-east</u>
- 35. Paleoclimate. <u>https://www.ipcc.ch/site/assets/uploads/2018/02/ar4-wg1-chapter6-1.pdf</u>
- 36. Oxygen balance. https://earthobservatory.nasa.gov/features/Paleoclimatology_OxygenBalance_
- 37. Proxy climate. https://en.wikipedia.org/wiki/Proxy (climate)#Boreholes