



PREVENTS ADULT FLEA EMERGENCE FOR UP TO 7 MONTHS

ACTIVE INGREDIENT:

(S)-Methoprene (CAS #65733-16-6)	1.2%
OTHER INGREDIENTS*	98.8%
TOTAL:	100.0%

*Contains petroleum distillates

EPA Reg. No. 2724-352 EPA Est. No. 2724-TX-1

CAUTION

INTRODUCTION

PRECOR® IGR CONCENTRATE (PRECOR®) acts on the immature life stages of the flea, preventing the adult flea from developing. Flea eggs deposited on, and larvae crawling onto treated areas will not develop into adult fleas. Existing adult fleas and flea pupae are not affected. If a high population of adult fleas (the biting stage) is present, tank-mix with an adulticide of choice that is EPA registered and properly labeled for flea control. To prevent infestations before they start, apply PRECOR® prior to flea season.

When applied to surfaces where fleas are found, such as carpets, upholstered furniture, and pet sleeping areas, **PRECOR®** prevents adult flea emergence for up to 7 months. Controlling fleas on pets is important to prevent introduction of adult fleas indoors. If fleas are a problem on pets, treat the pets with an effective EPA registered flea control product.

SPECIMEN LABEL

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if absorbed through the skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling.

CIDOT AID

FIKST AID	
Call a poison control center or doctor immediately for	
treatment advice.	
 Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. 	
 Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. 	

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-248-7763 for emergency medical treatment information.

NOTE TO PHYSICIAN: This product contains petroleum distillates. If swallowed: Immediately call a poison control center or doctor. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person.