

Dr. Jean Dodds Vaccination Protocol

9 – 10 weeks of age

Distemper + Parvovirus, MLV

e.g. Merck Nobivac (Intervet Progard) Puppy DPV

14 – 15 weeks of age

Distemper + Parvovirus, MLV

18 weeks of age

Parvovirus only, MLV

Note: New research states that last puppy parvovirus vaccine should be at 18 weeks old.

20 weeks or older, if allowable by law

Rabies – give 3-4 weeks apart from other vaccines

Mercury-free (thimerosal-free, TF)

1 year old

Distemper + Parvovirus, MLV

This is an optional booster or titer. If the client intends *not* to booster after this optional booster or intends *to* retest titers in another three years, this optional booster at puberty is wise.

1 year old

Rabies – give 3-4 weeks apart from other vaccines

3-year product if allowable by law; mercury-free (TF)

Perform vaccine antibody titers for distemper and parvovirus every three years thereafter, or more often, if desired.

Vaccinate for rabies virus according to the law, except where circumstances indicate that a written waiver needs to be obtained from the primary care veterinarian. In that case, a rabies antibody titer can also be performed to accompany the waiver request. Visit [The Rabies Challenge Fund](#) for more information.

W. Jean Dodds, DVM

[Hemopet](#) / [NutriScan](#)

11561 Salinaz Avenue

Garden Grove, CA 92843

Followed & Recommended by:

