Kellen Jex, MD Board Certified in Gastroenterology/ Internal Medicine



**Miss-Lou GI** 136 Jeff Davis Blvd. Suite B Phone: 601-492-2224 Fax: 601-492-2231

Kristin Foster, NP-C

Colonoscopy Procedure Information

Patient Name:

DOB

## DAY BEFORE PROCEDURE:

Do not eat solid foods. Only consume clear liquids throughout the day. Start prep at 5:00 pm. Do not eat or drink after midnight EXCEPT FOR the second dose of your bowel prep (see instructions).

## DAY OF PROCEDURE:

You will take the second dose of your bowel prep starting FIVE hours before your scheduled procedure time (see instructions).

\*Take your morning medications, if any, with a sip of water.

\*If you use an inhaler, take a treatment the morning of your procedure.

\*If you are a diabetic, please call your primary care provider for dosage changes at least by the day before the procedure.

\* If you take BLOOD THINNERS, it is VERY IMPORTANT to inform Dr. Jex so a plan can be made to either stop or continue these medications leading up to your procedure. BLOOD THINNERS include but are not limited to:

- Coumadin (Warfarin)
- Eliquis (Apixaban)
- Xarelto (Rivaroxaban)
- Plavix (Clopidogrel)
- Pletal (Cilostazol)

If you take any of the other following medications, hold these injections for two weeks prior to your procedure:

- Ozempic
- Semaglutide
- Wegovy
- Mounjaro
- Liraglutide
- Tirzepatide

Other Important Considerations:

- If you take INSULIN, you need to discuss this with your primary care provider before the day of your procedure.
- HOLD NSAIDS for SEVEN days prior to your scheduled procedure. YOU MAY NEED TO HOLD ASPIRIN; HOWEVER, please let our staff know if you have had a HEART ATTACK or STROKE. YOU MAY NEED TO CONTINUE YOUR ASPIRIN FOR YOUR PROCEDURE.
- DO NOT TAKE IRON PILLS FOR ONE WEEK LEADING UP TO YOUR PROCEDURE.