Kellen Jex, MD

Board Certified in Gastroenterology/ Internal Medicine



Miss-Lou Gl

136 Jeff Davis Blvd. Suite B

Phone: 601-492-2224 Fax: 601-492-2231

## Kristin Foster, NP-C

# **EGD + Colonoscopy Procedure Information**

Patient Name:			
DOB			

### **DAY BEFORE PROCEDURE:**

Do not eat solid foods. Only consume clear liquids throughout the day. Start prep at 5:00 pm. Do not eat or drink after midnight EXCEPT FOR the second dose of your bowel prep (see instructions).

#### **DAY OF PROCEDURE:**

You will take the second dose of your bowel prep starting FIVE hours before your scheduled procedure time (see instructions).

- \*Take your morning medications, if any, with a sip of water.
- \*If you use an inhaler, take a treatment the morning of your procedure.
- \*If you are a diabetic, please call your primary care provider for dosage changes at least by the day before the procedure.

\*If you take BLOOD THINNERS, it is VERY IMPORTANT to inform Dr. Jex so a plan can be made to either stop or continue these medications leading up to your procedure. BLOOD THINNERS include but are not limited to:

Coumadin (Warfarin), Eliquis (Alpixaban), Xarelto (Rivaroxaban), Plavix (Clopidogrel), Pletal (Cilostazol).

If you take any of the following medications, hold these injections for two weeks prior to your procedure:

- Ozempic
- Semaglutide
- Wegovy
- Mounjaro
- Liraglutide
- Tirzepatide

## **Other Important Considerations:**

- If you take INSULIN, you need to discuss this with your primary care provider before the day of your procedure.
- HOLD NSAIDS for SEVEN days prior to your scheduled procedure. YOU MAY NEED TO HOLD
  ASPIRIN; HOWEVER, please let our staff know if you have had a HEART ATTACK or STROKE. YOU
  MAY NEED TO CONTINUE YOUR ASPIRIN FOR YOUR PROCEDURE.
- DO NOT TAKE IRON PILLS FOR ONE WEEK LEADING UP TO YOUR PROCEDURE.