ESTHETIC BOUTIQUE MED SPA

Post treatment instructions for Neuromodulators.

After treatment possible side effects include: mild headache, bruising at injection site, a sensation of "numbness" or tingling at the injection site, feeling of pressure or fullness at injection site.

- 1. Remain in an upright position for 4 hours.
- 2. Do not lie flat or rub/scrub the area for four (4) hours after injection.
- 3. No massaging of the treated area for 1-3 hours. This can cause the toxin to migrate to a different area.
- 4. Emote the treated area (wrinkle the forehead, smile, frown, etc.) every so often for 1-3 hours after treatment. This will help with the uptake of the Botox.
- 5. Do not go to sauna/steam room or exercise (running, aerobics, weight lifting, etc...) for 24 hours.
- 6. Do not take aspirin or ibuprofen for 24 hours. This includes Advil. Motrin, Aleve, Naproxen, Bayer Baby Aspirin.
- 7. If you can take generic Tylenol, you may take Tylenol for any headaches.
- 8. Mild bruising can be seen after the procedure.
- 9. Arnica Gel is a homeopathic treatment that has great results on post treatment bruising and discomfort. It can be purchased at any Whole Food grocery store.
- 10. It will take 10-14 days for the full effect to be achieved.

Possible side effects include: mild headache, bruising at injection site, a sensation of "numbness" or tingling at the injection site, feeling of pressure or fullness at injection site

Call the Office for any of the following or any concerns:

- If you have a severe headache, fever, weakness, or other significant change in your health, seek emergency care.
- Notify us at 602-332-1660 to speak with us if any drooping of the eyelids (ptosis) or other unexpected side effects of treatment occur.