

EXPANDED PROGRAMME ON IMMUNISATION – EPI ISA! REVISED

Age of Child	Vaccines Needed	How and where it is given
At Birth	BCG Bacilles Calmette Guerin	Right Arm
	OPV (0) Oral Polio Vaccine	Drops by mouth
6 Weeks	OPV (1) Oral Polio Vaccine	Drops by mouth
	RV (1) Rotavirus Vaccine	Liquid by mouth
	DTap-IPV-Hib-HBV (1) Diphtheria, Tetanus, Acellular Pertussis, Inactivated Polio Vaccine, <i>Haemophilus influenzae</i> type b and Hepatitis B Combined	Intramuscular / Left thigh
	PCV (1) Pneumococcal Conjugated Vaccine	Intramuscular / Right thigh
10 Weeks	DTap-IPV-Hib-HBV (2) Diphtheria, Tetanus, Acellular Pertussis, Inactivated Polio Vaccine, <i>Haemophilus influenzae</i> type b and Hepatitis B Combined	Intramuscular / Left thigh
14 Weeks	RV (2) Rotavirus Vaccine*	Liquid by mouth
	DTap-IPV-Hib-HBV (3) Diphtheria, Tetanus, Acellular Pertussis, Inactivated Polio Vaccine, <i>Haemophilus influenzae</i> type b and Hepatitis B Combined	Intramuscular / Left thigh
	PCV (2) Pneumococcal Conjugated Vaccine	Intramuscular / Right thigh
6 Months	Measles Vaccine (1)**	Subcutaneous / Left thigh
9 Months	PCV (3) Pneumococcal Conjugated Vaccine	Intramuscular / Right thigh
12 Months	Measles Vaccine (2)**	Subcutaneous / Right arm
18 Months	DTap-IPV-Hib-HBV (4) Diphtheria, Tetanus, Acellular Pertussis, Inactivated Polio Vaccine, <i>Haemophilus influenzae</i> type b and Hepatitis B Combined	Intramuscular / Left arm
6 Years (both boys and girls)	Td Vaccine Tetanus and reduced strength of Diphtheria Vaccine	Intramuscular / Left arm
12 Years (both boys and girls)	Td Vaccine Tetanus and reduced strength of Diphtheria Vaccine	Intramuscular / Left arm

* Rotavirus Vaccine should NOT be administered after 24 weeks

** Do not administer with any other vaccine