

# Colonoscopy

**This leaflet explains more about your colonoscopy and what you can expect when you come to hospital. It also details the instructions on how to prepare for this test, including when to stop eating, how to cleanse your bowel and what tablets you may need to stop.**

**It is very important that no food or milk is consumed in the 24-hour period prior your examination. Please read the fasting and laxative instructions overleaf.**

## What is a colonoscopy?

Colonoscopy is a camera examination of the bowel. It helps to find the cause of your abdominal symptoms and to diagnose and sometimes treat diseases of the bowel. The procedure takes 30-45 minutes but please allow two to three hours in the hospital for the whole process to be completed.

## What are the risks?

This test is very safe, but very rarely there can be a problem, for example:

- The sedative can affect your breathing making it slow and shallow.
- There is a small risk of missing polyps (small growths) or other abnormalities. It is very rare to miss bowel cancer. Successful laxative preparation decreases this risk.
- When a biopsy is taken or a polyp is removed, on rare occasions there may be bleeding.
- Perforation, which is a tear in the wall of the colon is very rare (< 1:2,000 risk). If this happens it would require a short stay in hospital including treatment with antibiotics and possibly surgical repair.

## Are there any alternatives?

For some conditions it may be possible to perform a CT scan. The disadvantage of this procedure is that a biopsy cannot be taken, or a polyp removed. The radiation from the scan may make this unsuitable for some patients.

## If my symptoms have stopped before the colonoscopy should I still come for the test?

**Yes.** It is important that you still come for the test. Your doctor has organised this test to ensure you have no problems in your large bowel. Although the symptoms may have gone, it remains important to examine the bowel.

## How can I prepare for my colonoscopy?

- A clear view of the colon is required for this examination to be successful and so your bowel must be as empty as possible. You will have been prescribed either Citrafleet or Picolax or Moviprep which are laxatives designed to clear your bowel of stool. Please read the instructions below carefully.
- If you are diabetic or taking Warfarin, or blood thinning medications such as Apixaban, Rivaroxaban, Dabigatran, Clopidogrel, Ticagrelor or Prasugrel please contact us on the phone numbers on Page 5 of this leaflet under **Contact us**.
- Take other medications as normal, except iron tablets or stool bulking laxatives such as Fybogel or Movicol which should be stopped one week before your examination.
- If you are taking the contraceptive pill, please take additional precautions for one week following the bowel preparation.
- Please do not take oral medications one hour before or one hour after taking the bowel preparation.
- Please bring a list of your regular medications and bring any inhalers or sprays with you.

## Colonoscopy dietary and laxative preparation

Please follow the instructions below. **Failure to comply with the laxative preparation may lead to cancellation of your procedure for safety reasons.**

### Three days before the examination

Eat a light diet avoiding cereals, wholemeal bread, salads, fruits or any food containing fibre, nuts or seeds. Please remove the skins from all vegetables and tomatoes. Instead please eat low fibre food such as rice, pasta, fish, chicken, peeled potatoes, dairy products, white bread.

### The day before the examination

**DO NOT EAT FOOD FOR 24 HOURS BEFORE THE TESTS.** Instead take only *clear fluid* e.g., water, herbal tea, black tea or coffee (can have sugar but no milk), lemonade, sports drinks, orange squash. You may have clear soup (broth) or yellow / green jelly. You can continue clear fluids up until two hours before the examination.

**PLEASE DRINK 2-3 LITRES OF CLEAR FLUID DURING THE FASTING PERIOD.**

### If taking Citrafleet/Picolax sachets:

#### At 4pm on the day before the examination

Drink one sachet (mix with a glass of water). The drink may fizz. Drink a full glass of water after drinking the Citrafleet or Picolax. It can start to work within as little as 30 minutes but can take up to five hours to work. Please drink plenty of *clear fluids* throughout the afternoon and evening as above.

#### At 6am on the day of the examination

Drink the second sachet of Citrafleet or Picolax, even if you think the first one has already worked.

Please drink a further 500mls to a litre of *clear fluid* up to two hours before the colonoscopy.

For afternoon appointments you can take the second sachet at 9am.

**You must stop drinking two hours prior to the procedure.**

## **If taking Moviprep sachets:**

### **At 4pm on the day before the examination**

Make up your first jug of Moviprep by mixing both sachets A and B in a litre of water.

Stir the solution until clear. Drink a glassful every 10-15 minutes until it is finished.

It is also important to drink an additional 500ml of water or clear fluid during the evening.

### **At 6am on the day of the examination (or earlier if required)**

Make up your second litre of Moviprep repeating the steps above and drink it over 1-2 hours, drinking a glassful every 10-15 minutes until it is finished. Following this please drink an extra 500mls of clear fluid. For afternoon appointments you can start the second litre at 9am.

**You must stop drinking two hours prior to the procedure.**

## **What if my bowel preparation hasn't worked for hours after taking the laxative?**

**Please be patient.** We anticipate that the laxative will work within a few hours as outlined in the manufacturer's guidelines, but this can sometimes take a little longer depending on your age, diet, if you have diabetes and whether you suffer from constipation. If by the morning of your appointment you have had no result, please contact us for advice. Once it does start working please stay close to a toilet as sometimes no warning may be given.

## **What If the laxative makes me feel sick or nauseated?**

Occasionally, nausea maybe experienced. If this happens stop drinking the preparation for 15 minutes and then start again, slowly at first. If you experience nausea or vomiting, go to your local pharmacy and ask for anti-sickness tablets. Contact us if your symptoms persist.

## **Advice for diabetic patients**

The bowel preparation can upset your diabetes temporarily. For further advice about your diabetes, please contact your GP or your diabetic nurse.

## **Tips on bowel preparation**

Apply barrier cream (eg Sudocrem) to your anal area, as repeated bowel actions can make you sore. If you feel uncomfortable, stop temporarily, keep moving about and you should feel better. No alcohol should be drunk the day before or on the day of your examination.

## On arrival at the endoscopy unit

- A nurse will check your details, including medications and allergies
- The test will be explained again and you will be asked to sign a consent form.
- You will be asked to change into a gown.

## Asking for your consent

It is important that you feel involved in decisions about your care. You will be asked to sign a consent form to say that you agree to have the treatment and understand what it involves. You can withdraw your consent at any time, even if you have said 'yes' previously. If you would like more details about our consent process, please ask for a copy of our policy.

## What happens during colonoscopy?

- A sedative injection is offered. It makes you relaxed but usually not asleep. It can in some patients affect memory of the procedure. Some patients choose not to have the sedation as the test is easily tolerated in many patients. Entonox (gas and air) is available as an alternative.
- The colonoscope is passed into the bowel and air is introduced. You may feel discomfort as if you want to go to the toilet and short-lasting cramps can occur.
- A biopsy may be taken, or a polyp may be removed during the examination. You cannot usually feel this, but you can often visualise what is happening.

## Will I feel any pain?

You may feel some discomfort from the air that is pumped into the large bowel so that the endoscopist can view the lining adequately. You may also feel intermittent mild cramping when the colonoscope moves around the bowel, this is not usually painful. A mild sedative and painkiller are usually given for the procedure, but some people prefer not to have any. Entonox is also available: This is a "gas and air" mixture that can be used during the colonoscopy. It gives a short-lasting pain relief and patients can go home unaccompanied 30 minutes later. It is not suitable for patients with certain lung conditions or who have had certain eye/ear procedures.

## What happens after colonoscopy?

Following the tests, you will be taken to the recovery area where you will be monitored for 45 minutes to 1 hour. After this you will be able to get dressed and have some refreshments.

If you have had sedation you will not be allowed to drive and must arrange for someone to accompany you home. The medication given during the test will prohibit you from driving for 24 hours after the examination. Please do not plan to take public transport home unless accompanied. If you are unable to arrange transportation we can arrange a taxi to take you

home however you are responsible for the fare. You will need a responsible adult at home for at least 12 hours. Once your escort has arrived to accompany you home a nurse will give you a copy of the endoscopy report as well as some important discharge advice. If you have had no sedation or choose to have Entonox only you can leave unaccompanied. A normal diet can be resumed once the procedures are complete.

## **Results / Will I have a follow-up appointment?**

Upon completion of the colonoscopy the findings will be discussed with you. We will be able to tell you of any visual findings, however any samples will need to be sent to the laboratory for testing. This can take up to three weeks.

## **Useful sources of information**

<https://www.nhs.uk/video/Pages/Colonoscopy.aspx>

includes further explanation and videos about colonoscopy.

## **Contact us**

If you have any questions or concerns about your examination, please contact us:

[info@ramwellsurgicalservicesltd.co.uk](mailto:info@ramwellsurgicalservicesltd.co.uk)

02079521468 (Practice Manager)

