(neuroHPI2)

Numbness: O none	O right arm O right hand	O left arm
O left hand O	right leg O right foot O left leg	O left foot

Muscle spasms: O none O neck O mid back O lumbar O right arm O left arm O right leg O left leg

## **Duration of Symptoms:**

O 1	O 2	O 3	O 4	0	5	Ο	6	0	7	0	8	0	9	0	10	0	11	0	12
O sev	veral	O da	y(s)	0	we	eek(	s)	0	mo	onth	n(s)	0	yea	ar(s	5)	0	ch	ron	ic

Aggrevating factors: O nothing in particular O bending O lifting O twisting O activity O standing O walking O extending head

- O twisting head O reaching O leaning forward
- O getting up from a seated position O straightening
- O working at job O driving O laying down

## Alleviating factors: O nothing in particular O sitting

- O laying flat O rest O moving around
- O leaning on a grocery cart O laying on side
- O laying flat with legs flexed and elevated O sleeping
- O raising arm over head

(neuroHPI3)

**Treatment**: O nothing to date O physical therapy

- O epidural steroid injections O narcotic pain meds
  - O pain management O muscle relaxants
  - O nonsteroidal antiinflammatory meds O rest O heat
  - O cold/ice O trigger point injections O chiropractor
  - O oral steroids O accupuncture O massage O TENS unit
- O rhizotomies O back brace O neck brace O botox
- O traction

Length of conservative treatment: O nothing to date O 1 week

O 1 month O at least 6 months O several days O several weeks O several months O several years

Bowel incontinence: O Yes O No

## Bladder incontinence: O Yes O No

O Chronic stress incontinence