## Case 2A:

Hank is a 54-year-old male with opioid use disorder, in sustained remission. He was medically exposed to opioid pain medication after he was in a car accident in his 30s, and he began misusing them via insufflation. He was abstinent for 12 months after attending an inpatient treatment program 10 years ago, but he returned to use for 2 weeks when his mom died. He has remained in recovery since that time, initially being prescribed medication for opioid use disorder (mOUD) which he tapered to discontinuation after 1 year. His blood pressure is controlled on lisinopril, and his lipids are controlled on rosuvastatin. Hank has been an IT Specialist at the same company for the last 5 years.

Rate the level of risk:

1 2 3 4 5 6 7 8 9 10 Least Most

## Case 2B:

Brenda is a 38-year-old hairdresser who is an alcoholic who has been clean for 5 years. She began drinking in her 20s which progressively escalated over time to the point that she was occasionally missing work and began feeling depressed. She decided to go to rehab 5 years ago and was prescribed naltrexone at that time. Her discharge diagnoses were alcohol abuse and alcohol-induced depressive disorder; however, her mood was noted to improve during the treatment course. After discharge, she began attending Alcoholics Anonymous and continues to attend regularly. She discontinued naltrexone 4 years ago due to an elevation in her liver enzymes after which her LFTs returned to normal; however, she managed to stay clean and sober since rehab was completed.

Rate the level of risk:

1 2 3 4 5 6 7 8 9 10
Least Most