

TECHNOLOGICAL ORGANIZATION IN THE LATE HOLOCENE: RESULTS FROM THE CLEARVIEW SITE (XMH-1303)

Briana N. Doering

Department of Anthropology, University of Wyoming, 1000 E. University Ave., Dept. 3431, Laramie, WY 82071;
bdoerin2@uwyo.edu

Julie A. Esdale

Center for the Environmental Management of Military Lands, Colorado State University, Directorate of Public Works,
ATTN: IMFW-PWE (Esdale), 1046 Marks Street, #6000, Ft. Wainwright, AK 99703-6000; julie.esdale@colostate.edu

Senna D. Catenacci

Museum of Anthropological Archaeology, University of Michigan, 3010 School of Education, 610 E. University Ave.,
Ann Arbor, MI 48109; sennac@umich.edu

ABSTRACT

Roughly 1500 years ago, a group of hunter-gatherers left behind numerous tools and thousands of pieces of lithic production debris at the Clearview site, situated on a rise overlooking the middle Tanana Valley and the Alaska Range in central Alaska. This paper considers the spatial relationships between tool type, debitage production, and material to demonstrate the technological organization at this late Holocene occupation. The diverse assemblage considered here comprises several tools (projectile points, bifacial knives, end and side scrapers, expedient flake tools, burins, and microblades) and raw materials, including obsidian. A comprehensive analysis of this assemblage reveals that the site's occupants undertook a complex lithic reduction sequence focused on intermediate stages of bifacial reduction and late-stage microblade production. Generally, this assemblage appears to represent a residential camp of mobile foragers similar to those associated with the Northern Archaic tradition. The results presented here suggest that significant late Holocene behavioral shifts (e.g., increased subsistence specialization, decreased mobility, increased use of metal and bone tools) occurred after Clearview was occupied.

INTRODUCTION

The middle Tanana Valley has provided archaeologists with a wealth of material data related to subsistence, technology, and mobility spanning the last 14,000 years (Goebel and Potter 2016; Holmes 2008). Archaeologists in this region have developed a comprehensive understanding of human responses to climatic and ecological conditions for the first several thousand years of human history in eastern Beringia (Goebel et al. 2011; Graf and Bigelow 2011; Potter 2011). However, single-occupation sites from later time periods, particularly the mid- to late Holocene,

are rarely subjected to comprehensive testing and synthetic analysis that could broaden anthropological understandings of human behavior in the late Holocene and inform our understanding of life in the middle Tanana region.

The lithic assemblage from the Clearview site (XMH-1303) builds on the wealth of data archaeologists have collected through nearly a century of research. This single-component site has an associated lithic assemblage with diagnostic information that can be related to technological organization, raw material use, and settlement

patterning. Moreover, radiocarbon dates from this site indicate that it was occupied between two ashfall events that may be linked to the Athabaskan transition (Mullen 2012; Workman 1979). Here, we present the results of a comprehensive analysis of this site's lithic assemblage in order to provide additional comparative information for other late Holocene occupations and site use across the region.

BACKGROUND

SETTING AND PREVIOUS RESEARCH

The Clearview site is in the middle Tanana Valley just north of the Alaska Range. The site is in an intermediate ecological zone at the transition from lowlands to uplands in the foothills of the Granite Range, which extends to the southeast (Fig. 1; Gallant et al. 1995). Clearview is situated on a small rise at 460 m above sea level and is so

named because of the site's exceptional 360-degree view of the Tanana Valley and Donnelly Dome to the south and west, the Granite Mountains to the southeast, and the Yukon-Tanana Uplands to the northeast. Banjo Lake and a small unnamed lake lie to the north and south of the site, respectively. The site is on the perimeter of productive upland and lowland areas that offer a diverse array of resources in a setting that provides a unique vantage point of the surrounding area.

Archaeologists from Colorado State University's Center for Environmental Management of Military Lands (CEMML) recovered the first artifacts from the Clearview site in 2006 (Robertson et al. 2013). The Clearview site was located through shovel testing during a survey for the construction of the Battle Area Complex at Fort Wainwright, Alaska, and was found eligible for the National Register of Historic Places (Carlson et al. 2017; Robertson et al. 2007).




Figure 1. The Clearview Site (XMH-1303) in relation to other late Holocene sites in the middle Tanana River Valley.

CEMML archaeologists returned to Clearview in 2009 for additional site testing, which resulted in 40 systematically placed 1 x 1 m units that defined the site's area and tentatively identified the central activity area of the site (Robertson et al. 2013). Following these investigations, this site became a part of the Jarvis Creek Archaeological District (Carlson et al. 2017). Archaeologists have identified dozens of sites in the Jarvis Creek Archaeological District, but few have been comprehensively tested, and many lack the stratigraphic context or materials that are appropriate for radiocarbon dating (Robertson et al. 2009, 2013). Further, very few of the sites that archaeologists have tested and radiocarbon dated have late Holocene components. Therefore, Clearview has the potential to further our knowledge of behavior during this time as one of the few sites within the Jarvis Creek Archaeological District that has been comprehensively tested and dated to the late Holocene.

THE LATE HOLOCENE IN CENTRAL ALASKA

Anthropologists have provided detailed documentation of central Alaska life for almost a century, beginning with Rainey (1940) and Osgood (1937) in the 1930s. Archaeological and ethnographic research conducted over the last century suggests that central Alaskans pursued a semisedentary, logistical subsistence strategy based on accumulating seasonally abundant resources such as salmon and caribou (Potter 2016). Central Alaskans moved seasonally between large residential encampments, beginning with salmon camps in the summer, transitioning to fall hunting camps, and later moving on to winter trapping camps. Archaeological evidence from late Holocene sites in central Alaska such as Dixthada (TNX-004) and Nenana River Gorge (HEA-062) shows evidence for large seasonal aggregations with a range of material culture, including copper tools and pottery (Plaskett 1977; Shinkwin 1979), which resemble ethnographically documented summer fish camps (Andrews 1975; Osgood 1937). This extensive research provides a general account of life during the late Holocene that could be improved by additional archaeological investigations at sites that represent different roles within the seasonal round.

Archaeological research focused on the earlier periods of Alaska history has provided material evidence from sites across central Alaska, particularly in the middle Tanana Valley (Goebel and Potter 2016). Archaeologists have doc-

umented dozens of radiocarbon-dated occupations from the late Pleistocene through the late Holocene with assemblages that are indicative of small, short-term hunting camps with little evidence for repeated occupations (Potter 2008, 2016). However, evidence from the latest pre-European sites and ethnographic records (cited above) suggests that central Alaskans occupied large seasonal camps repeatedly. This discrepancy has led archaeologists to argue that either: (1) taphonomic processes have obscured these larger seasonal camps or village sites (Yesner 1996), or (2) central Alaskans pursued a highly mobile subsistence strategy until 1500 to 1000 years ago, when groups in the region began to return to camps seasonally (Potter 2008). Additional archaeological evidence from the late Holocene could address this discrepancy in occupation strategies spanning the Holocene.

Researchers have interpreted the long cultural record of central Alaska in terms of Binford's (1980) model of residential and logistical mobility, based in part on observations he made of Alaska subsistence hunters in the Brooks Range. Residential mobility is associated with a series of medium-sized generalized base or residential camps, where a wide variety of subsistence behavior is carried out and consumers move to resources. In contrast, logistically mobile groups occupy several smaller, task-oriented sites that move resources to a larger central residence. Despite the potential archaeological ambiguity between these systems of residential mobility, archaeologists have drawn heavily on this model to contextualize the significant changes documented in the central Alaska material record and have suggested that central Alaskans shifted from a system of high residential mobility to a logistically mobile, semisedentary system during the late Holocene.

Anthropologists have argued that increasing sedentism and logistical mobility coincided with changes in subsistence and technology in central Alaska. Archaeologists have recovered very little material evidence for extensive fishing in central Alaska prior to the late Holocene (Potter 2016), when consistent salmon exploitation began. Additionally, archaeologists have argued that stone tool technology was replaced by bone, antler, and copper tools during the late Holocene, based on several assemblages from central Alaska and Yukon (Holmes 2008; Potter 2008). Finally, evidence from ice patches in Yukon suggests that atlatl and dart technology was replaced by the bow and arrow at this time (Hare et al. 2004). These behavioral changes in subsistence and technology are comparable to other North

American hunter-gatherer transitions to semisedentism that took place throughout the Holocene (Chatters and Prentiss 2005; DeBoer 1988).

The late Holocene behavioral transition, inferred from the shifts in the archaeological record outlined above, is also known as the “Athabascan transition” (Helmer et al. 1977). Explanations for these changes have centered around volcanic eruptions emanating from Mount Churchill in the Wrangell–St. Elias mountain ranges in southeast Alaska (Workman 1979). Two late Holocene volcanic events, ca. 1700 cal BP and 1100 cal BP, produced thick layers of ash with possible ecological effects in eastern or central Alaska, and are known as the North and East Lobes of the White River Ash (WRA), respectively (Jensen et al. 2014; Lynch et al. 2018; Mullen 2012). Geologists have described the later East Lobe WRA event in particular as one of the most significant in the region’s history (Péwé 1975), and biologists have linked this event and subsequent decimation of lichen populations to a documented genetic bottleneck in the region’s caribou herds (Kuhn et al. 2010). Anthropologists have suggested that Athabascan oral histories that reference a massive volcanic eruption tie this behavioral change to the ashfall and provide further evidence for the event’s dramatic impact on central Alaska life and subsistence (Moodie et al. 1992; Mullen 2012; Workman 1979). Moreover, paleoenvironmental reconstructions suggest that the impacts of the WRA events may have stood out in an otherwise stable climatic regime (Anderson et al. 2003; Kaufman et al. 2004). The Clearview assemblage offers archaeological context for the impact of the North and East Lobe WRA events, which occurred before and after the site’s occupation, respectively.

Additional archaeological evidence from sites that date to this period can provide anthropologists with an improved chronological and theoretical understanding of the late Holocene behavioral transition and potential impacts of the White River Ash events. Clearview was occupied ca. 1500 cal BP, on the verge of the Athabascan transition. The North Lobe WRA event occurred ca. 200 years prior to Clearview’s occupation, and the East Lobe WRA event occurred ca. 350 years later. The North Lobe WRA event likely would have had a more direct effect on central Alaskans due to the dispersal of the ash northward rather than eastward. Thus, the Clearview assemblage should reflect any long-term behavioral shifts that took place following the North Lobe WRA event ca. 1700 cal BP (Lynch et al. 2018; Mullen 2012). Evidence from

late Holocene hunting camps can also serve as a bridge between the expansive archaeological data sets from earlier periods of Alaska history and the wealth of knowledge found in ethnographic accounts. Therefore, the Clearview site, a comprehensively excavated and securely dated late Holocene hunting camp, provides an important link that can improve our understanding of Alaska’s past.

TECHNOLOGICAL ORGANIZATION

Large lithic assemblages present several proxies for behavior that could be used to determine whether the North Lobe WRA event led to increased sedentism and a shift to logistical mobility among central Alaskans. Archaeologists in Alaska and elsewhere have successfully applied artifact class richness, local and exotic material use, and lithic reduction strategy to reconstruct occupation length, mobility patterns, and subsistence strategies. Further, these behavioral correlates can be extrapolated to regional site patterning from different periods of central Alaska history to better evaluate and explain the timing of the Athabascan transition.

Artifact class richness represents the number of tool types present in an assemblage and serves as a proxy for site use (Bamforth 1986; Cowan 1999; Shott 1986). High artifact class richness indicates longer occupations and/or generalized site use associated with residential bases (Habu 2002). In contrast, low artifact class richness indicates specialization and/or shorter occupations, associated with logistical camps. Therefore, artifact class richness conveys broader behavioral patterns related to the functional and organizational role of sites that can be used to compare across assemblages. Additionally, a typological debitage analysis permits a synthesis of the reduction sequence within an assemblage and can build upon information offered by formal tools to show how an assemblage fits in to broader behavioral and settlement patterns (Andrefsky 2005:201–210; Ostahowski and Kelly 2014). At Clearview, numerous and diverse artifacts and debitage would indicate a longer, residential occupation, whereas a logistical hunting camp would exhibit late-stage debitage related to sharpening and retouch (Surovell 2009).

Raw material variability represents broader patterns of mobility and exchange, particularly when exotic materials are present (Andrefsky 1994). Archaeologists have suggested that a lower ratio of nonlocal or exotic materials to local materials correlates with longer occupation lengths (Surovell 2009:79). Within Alaska assemblages,

the presence and source of obsidian also demonstrates broader trends in mobility and trade that the North Lobe WRA event may have disrupted due to its proximity to Wiki Peak, an important Alaska obsidian source (Reuther et al. 2011). Indeed, previous raw material sourcing research on pyrometamorphic rocks, or clinkers, traded in the Northwest Territories, Canada, has indicated that the East Lobe WRA disrupted trade in the western subarctic (Kristensen et al. 2019). Decreased use of Wiki Peak obsidian could represent larger disruptions in mobility or trade related to the preceding North Lobe WRA event.

The association between raw material use and tool type can further indicate patterns in skill acquisition and division of labor (Amick 1999) and serve as a proxy for environmental variability, predictability, and risk mitigation. A reliance on complex, patterned reduction strategies can result in more efficient use of raw materials (Elston and Brantingham 2002) but necessitates a great deal of skill and can be risky in resource-poor environments (Bamforth and Finlay 2008). If the North Lobe WRA dramatically increased ecological unpredictability and risk, we would expect to see reliance on generalized and easily maintained tools, such as bifaces, and absence of complex, patterned reduction or specialization at Clearview. In sum, if the volcanic activity associated with the North Lobe WRA event resulted in a rapid shift in central Alaskan behavior, Clearview should demonstrate reduced trade in exotics, debitage consistent with a generalized toolkit, and a low artifact class richness associated with a shift to logistical mobility.

METHODS

EXCAVATION METHODS

Between 2016 and 2018, CEMML archaeologists conducted additional testing as part of mitigation related to

the construction of a fire break around the range's target complex. These excavations expanded the previously excavated 1 x 1 m test units into a 6 x 5 m block to identify specific lithic production areas within the area of highest artifact concentration, as well as three 2 x 1 m blocks that investigated other areas with high potential to yield additional artifacts. All excavated artifacts will be permanently housed at the University of Alaska Museum of the North (UA2016-136, UA2017-92, UA2018-71). Excavation methods remained consistent during the four seasons of excavation completed between 2009 and 2018. Each 1 x 1 m unit was excavated in arbitrary 5 cm levels by 50 x 50 cm quadrants. Through this excavation strategy, approximately 10% of the total site area was excavated. Diagnostic materials were three-point provenienced using a Sokkia Set 6 Total Station, and all material was screened through 1/8-inch hardware cloth.

During excavation, charcoal samples were collected for chronological control, and strata were sampled and recorded for geoarchaeological analysis. Four samples of culturally associated wood charcoal from two locations were submitted for AMS radiocarbon dating at the National Ocean Sciences Accelerator Mass Spectrometry facility at the Woods Hole Oceanographic Institution (Table 1). The first location, N 501 E 97, was targeted for sampling because of the association between a visible charcoal lens and several diagnostic artifacts at the base of Stratum III. Three radiocarbon dates were submitted from this context. The second, N 497 E 97, also represented a cluster of charcoal associated with a dense cluster of lithic debitage that was used to corroborate results from the first location. Soil samples were collected at 5 cm intervals to understand the impact of soil acidity on preservation at Clearview. Soil pH was assessed by Matt Ferderbar at the Cold Regions Research and Engineering Laboratory, Fort Wainwright.

Table 1. Radiocarbon results and calibrated dates from Clearview.

NOSAMS lab no.	Field sample no.	Northing	Easting	RCYBP (1σ)	Cal BP (2σ)
OS-130783	478	497.828	97.29	1250 ± 40	1168–1278
OS-130784	467	501.467	97.015	1720 ± 40	1545–1715
OS-130785	468	501.323	97.323	1540 ± 30	1365–1524
OS-130786	474	501.37	97.268	1550 ± 30	1377–1527

Radiocarbon dates calibrated with Calib 7.1 using the IntCal13 curve (Stuiver et al. 2019)

Lithic Analysis

All excavated materials were analyzed at the University of Michigan Museum of Anthropological Archaeology following widely practiced identification methods (Andrefsky 2005; Esdale 2009). Tools and debitage were analyzed separately. Tool and tool fragments were weighed, and material type was assessed through comparison to toolstone types found in local drainages (e.g., Jarvis Creek) and neighboring sites (e.g., Banjo Lake). Finally, tools and tool fragments were analyzed in comparison to known tool types from central Alaska and sorted into six broad technological categories: uniface, biface, burin, blade, microblade, and retouched flake.

A typological analysis of lithic debitage took place in three general phases. First, artifacts were counted, weighed, and cleaned, and the raw material of each piece was identified through a visual analysis. Jeff Rasic, National Park Service, analyzed the obsidian debitage separately with X-ray fluorescence with a portable Bruker Tracer III-V. Second, lithic pieces with an intact bulb of percussion, platform, and terminating edge were counted and separated for additional analysis. These pieces of debitage were individually weighed and assigned a size class on a base two scale, beginning at 1 cm². Next, these pieces were assessed individually for presence of cortex, heat treatment, and use-wear.

Finally, each piece of debitage was assigned one of 13 production-phase categories, following Esdale (2009). General production phase categories distinguished between early reduction, bifacial reduction, unifacial reduction, and microblade reduction. Early reduction flakes were further separated into primary decortication (> 50% cortex), secondary decortication (10%–50% cortex), and interior flakes (0%–10% cortex). Debitage related to bifacial reduction was separated into early thinning, late thinning, edge preparation, alternate, and bifacial pressure flakes. Microblade reduction debitage was sorted into core face rejuvenation flakes, platform rejuvenation flakes, and linear flakes.

Following this visual analysis of lithic materials, the results were compared using statistical methods to assess the variation between material type, tool type, and phase of production. Statistical comparisons between material and tool types were made using a Fisher's exact test. Fisher's exact tests evaluate categorical data in a 2x2 contingency table and are similar to chi-square tests but better accom-

modate comparisons between small (e.g., $n < 10$) and large samples. Therefore, this test is well suited to Clearview's lithic assemblage. In this study, results with $p < 0.05$ are considered significant.

Spatial Analysis

As a single-component site, spatial data associated with the recovered artifacts from the Clearview site were considered in the horizontal plane. This analysis revealed the boundaries of artifact clusters, specific areas of activity across the site, and spatial relationships between raw materials and tool types. Two-dimensional spatial data was input into ArcGIS Desktop 10.6 as raster and point data. Raster data comprised lithic debitage, and point data comprised individual diagnostic tools and tool fragments. Artifact distribution was assessed based on raw material type, debitage category, and presence of cortex.

To identify other underlying patterns in tool manufacture and use, debitage recovered in the central block of Clearview was considered in two phases following methods standard in geospatial statistical analysis and artifact patterning (Wandsnider 1996). First, two distinct activity areas, or centers of debitage production, were spatially designated through a k -means cluster analysis, and debitage results were either attributed to one of these two spatial clusters or excluded from the analysis if they were outside these central activity areas. Second, Fisher's exact tests were applied to assess differences in lithic production based on debitage type, phase, and raw material between these two activity areas. In sum, we evaluate technology production at Clearview based on spatial and artifact attributes.

RESULTS**EXCAVATION RESULTS**

During four seasons of fieldwork at Clearview, CEMML archaeologists excavated 76 m², resulting in the recovery of 5138 lithic artifacts and an average density of 67 artifacts per m². The artifact density at the Clearview site is consistent with a single-component site. Archaeological materials were primarily recovered in Levels 2 and 3, approximately 10–20 cm below surface (Fig. 2). Flake frequencies by unit produced unimodal distributions across the site, indicating that the materials recovered from the Clearview site were deposited synchronically. No faunal materials were recovered during these excavations.




Figure 2. Stratigraphy at Clearview. Drawn in the field by Katherine Mulliken.

Stratigraphy

Macromorphological indicators suggest that Pleistocene glacial processes and Holocene aeolian activity likely shaped the parent material at the Clearview site. Further, its stratigraphic context is very similar to the neighboring Banjo Lake site (Esdale et al. 2015). The sediments at Clearview can be organized into three primary stratigraphic units: glacial outwash, silts with evidence for several episodes of soil formation, and humic mat. The deepest stratigraphic unit comprised poorly sorted glacial outwash, likely derived from subglacial eskers and kames (Reger et al. 2008). In 2009, excavations recovered no archaeological materials in this stratum, and all subsequent excavations were terminated at contact with this stratigraphic unit, approximately 30–40 cm below surface.

Above the glacial deposit, a thick layer of silt is further divided into four horizons based on color with evidence of soil development, likely related to the succession of several coniferous boreal forests throughout the Holocene (Ping et al. 2008). Cultural materials primarily appeared in a weak B horizon (Bw) 10–20 cm below surface in the upper silt. Finally, Stratum I represents the humic mat of the organic horizon. Although some vertical mixing of artifacts may have occurred, disturbance to the stratigraphic

integrity of the site appears minimal due to the concentrations of lithic material, which will be discussed in greater detail in the following section.

Site Chronology and Soil Chemistry

All four charcoal samples collected from Clearview resulted in radiocarbon dates that are consistent with a late Holocene occupation of the site, ca. 1500 cal BP (Table 1). The results of a pH analysis on soils collected from different stratigraphic units at Clearview show that Stratum III, which contains the majority of the cultural material at Clearview, is also the most acidic unit. With a pH of 4.39, this stratum falls within the pH range least conducive to the preservation of faunal material (Table 2).

ANALYTICAL RESULTS

The Clearview assemblage contains 55 tools, tool fragments, and cores. Both expedient and formal tools and tool fragments are present in the assemblage (Figs. 3–5). Over half of these are complete tools (56.4%), including expedient flake tools, burins, blades, and bifacial points and knives. The other items present in the assemblage are bifacial fragments (e.g., projectile point bases and tips) or

Table 2. Results of soil pH analysis on each stratigraphic unit.

Field sample no.	Stratum	pH 1:5 DI H ₂ O	pH 1:5 0.1M CaCl ₂
149	I	4.8	4.77
150	II	4.51	3.84
151	III	4.39	3.38
152	IV	5.28	5.06
148	V	5.83	5.11
153	VI	5.46	4.41

microblade cores. The assemblage also contains 82 microblades and microblade fragments, or linear flakes with evidence for use-wear. The range of technology present at Clearview signifies the broad range of activities that may have taken place during the occupation of the site.

More than 4400 pieces of debitage were recovered during excavations at the Clearview site. Debitage, including complete flakes and shatter, represents 86% of the total artifact assemblage. Complete flakes (debitage with intact platforms and identifiable bulbs of percussion) represent 34% of the total flake assemblage (Table 3). The average weight of complete flakes was 0.71 g,




Figure 3. Bifaces (a–c) and biface fragments (d–f) recovered from Clearview (University of Alaska Museum of the North accession numbers UA2011-401, UA2016-136, UA2017-92, UA2018-71). Photographed by Whitney McLaren.




Figure 4. Microblade core fragments (a–b), blade (c), and microblades and microblade fragments (d–j) recovered from Clearview (University of Alaska Museum of the North accession numbers UA2011-401, UA2016-136, UA2017-92, UA2018-71). Photographed by Whitney McLaren.




Figure 5. End scrapers recovered from Clearview (a–c), including one composite end scraper and shaft straightener (c) (University of Alaska Museum of the North accession numbers UA2011-401, UA2016-136, UA2017-92, UA2018-71). Photographed by Whitney McLaren.

Table 3. Results of debitage analysis by production phase and material type.

Raw material	Initial core reduction					Microblade Reduction				
	Primary decortication	Secondary decortication	Interior flake	<i>n</i>	%	Core face rejuvenation	Platform rejuvenation	Linear flake	<i>n</i>	%
Andesite	4	9	8	21	45.7			1	1	2.2
Banded gray chert	1	1	3	5	11.4			9	9	20.5
Black chert	4	13	34	51	8.3	1	3	80	84	13.6
Brown chert								1	1	6.7
Chalcedony		1	4	5	5.1	1	1	2	4	4.0
Gray chert	9	7	25	41	7.9	1	2	29	32	6.2
Obsidian								16	16	69.6
Red chert			1	1	4.0					
Rhyolite	1	3	14	18	2.4	2	1	17	20	2.6
Quartz		1	1	2	28.6					
Quartzite										
White chert										
Total	19	35	90	144	6.6	5	7	155	167	7.7
%	13.1	24.3	62.5			3.0	4.1	92.8		

Raw material	Bifacial reduction					Unifacial reduction				All
	Early thinning	Late thinning	Edge preparation	Alternate	Bifacial pressure	<i>n</i>	%	Unifacial pressure	%	<i>n</i>
Andesite	10	7	7			24	52.2			46
Banded gray chert	15	7	1	1	5	29	65.9	1	2.3	44
Black chert	218	129	78	20	31	476	77.0	7	1.1	618
Brown chert	8	3	3			14	93.3			15
Chalcedony	27	31	17	5	9	89	89.9	1	1.0	99
Gray chert	123	160	69	34	57	443	85.5	2	0.4	518
Obsidian	3	2			2	7	30.4			23
Red chert	6	9	2	2	5	24	96.0			25
Rhyolite	103	308	113	83	112	719	95.0			757
Quartz	1	2	2			5	71.4			7
Quartzite	2	1	1			4	100.0			4
White chert	4	3	2		1	10	100.0			10
Total	520	662	295	145	222	1844	85.1	11	0.5	2166
%	28.2	35.9	16.0	7.9	12.0					

and 78% of these flakes were 1 cm² or smaller. Finally, only 54 pieces of debitage exhibited any evidence of cortex. These data suggest that the debitage deposited at the Clearview site was primarily related to intermediate tool production and maintenance.

Raw Materials at Clearview

A visual analysis of color, grain size, and luster revealed that at least 13 individual cobbles or raw materials were brought to this site. Of these, there were seven subcategories of semisedimentary chert or chalcedony, three subcategories of volcanic material, and two subcategories of metamorphic rock (Table 3). Within these subcategories, any additional variations in color and texture were determined to be too minimal to warrant additional objective subdivision. Additionally, only three artifacts in the assemblage demonstrated possible evidence of heat treatment, including coloration, heat fracture, or pot-lidding. This, combined with the absence of hearths at the site, indicates that these artifacts may have been heat treated off-site. The raw materials present within Clearview's assemblage show that a wide variety of local and nonlocal stone was brought to the site, and heat treatment was uncommon.

Local materials dominated the lithic assemblage from Clearview. Black chert, gray chert, and rhyolite comprised 85% of the assemblage. A short survey of the seasonal Jarvis Creek, 2 km west of the site, resulted in the recovery of large cobbles of each of these materials. Previous toolstone surveys suggest that these materials are also easily found in eroding glacial kames throughout the area (Esdale et al. 2015). These sources are within 20 km of the site, or a day's walk, and meet Surovell's (2009:78) definition of local toolstone. Cortex present on primary reduction debitage appears to be cobble cortex, further indicating that these materials were collected from riverbeds rather than mined from geological sources. Based on the results of a comparative visual analysis, the overwhelming majority of raw materials used at Clearview are locally abundant.

The assemblage contains at least one type of obsidian derived from a nonlocal source. Alaska archaeologists have generated comprehensive geochemical profiles of Alaska obsidian using portable X-ray fluorescence and have identified four sources of this raw material across the region (Reuther et al. 2011). One obsidian microblade was conclusively sourced to Wiki Peak, located more than 300 km to the southeast in the Wrangell–St. Elias Mountain range

(Doering et al. 2019). Seven additional microblade fragments were tentatively sourced to Batza Tena, though these fragments were too small for a confident quantitative sourcing assessment. Nevertheless, it is clear from these data that occupants of Clearview used obsidian from at least one distant source. Aside from obsidian, exotic or nonlocal materials within the assemblage are more difficult to assess with certainty. Other potential nonlocal materials include fine-grained red and white chert, chalcedony, and jasper (Esdale et al. 2015).

Early Stage Core Reduction

Early reduction debitage, identified by the presence of cortex on individual pieces, flake scars, and overall size, represented only 9.3% of the total assemblage. Only cobble cortex was identified in the assemblage, indicating that no materials were quarried from bedrock outcrops. None of these pieces were produced on exotic raw materials, and 80.7% were produced on rhyolite, black chert, or gray chert. Further, no cobble cores or tested cobbles were recovered during excavations at Clearview. Overall, early reduction debitage comprises a small part of the overall assemblage and is made from the dominant local raw materials.

Bifacial Technology

Bifacial projectile points and projectile point fragments represent approximately one-third of the tools within the Clearview assemblage (32.7%). More than half of the bifacial technology in the assemblage is fragmentary, with only five complete bifaces. Nevertheless, fragmentary and complete bifacial technology in the assemblage indicates that at least three styles of bifacial technology were used at the site: bifacial knives, lanceolate projectile points, and straight-based projectile points. Only two of the 18 bifaces or biface fragments were made on a potentially nonlocal type of chert.

In comparison, nearly two-thirds of intact debitage (74.8%) is related to the reduction of large flakes or blanks into bifaces, reflecting the importance of intermediate bifacial reduction at the site. However, bifacial pressure flakes, typically removed with soft percussion to sharpen or resharpen the edge of a biface, were not common and represent only 12.0% of bifacial debitage. Additionally, the assemblage contains only one biface blank and no bifacial cores associated with early bifacial production. The lack of blanks and bifacial pressure flakes is surprising

given the quantity of bifaces and biface fragments at the site and extensive evidence for intermediate bifacial reduction in the assemblage. Finally, very few pieces of bifacial debitage were produced on nonlocal materials (6.0%) in contrast to microblades and related debitage (see below), reflecting general trends for raw material use observed in bifacial tool fragments recovered from the site.

Blade Technology

Three microblade cores or core fragments, three blade fragments, and 155 microblades are present in the Clearview assemblage. However, no microblade core tablets or blade cores were recovered. The three microblade cores recovered at Clearview are all consistent with a wedge-shaped style that is common in Alaska assemblages from the mid- to late Holocene (Coutouly 2012). Two cores were made on biface fragments and one was made on a large flake, suggesting that the use life of raw materials at Clearview was extended by converting spent bifaces into microblade cores. Further, several crested blades in the assemblage provide additional evidence that microblades were commonly made on expended bifaces or biface fragments. This style is common in the small number of Alaska assemblages containing microblades that have been dated to the late Holocene (Holmes 2008).

Two of the three microblade cores are made from potentially nonlocal or rare red chert and agate, and the third is made of rhyolite, which suggests that microblade production served to conserve rare toolstone. Three blade fragments were recovered from Clearview. Two of these blade fragments refit and show signs of retouching. These blades are nearly 2 cm wide and significantly larger than the microblades in the assemblage, and no cores or core fragments were recovered, suggesting that these finished blades were brought to the site. All three fragments are made from chert, though the two refitting and retouched fragments are the only examples of dark red fine-grained chert in the assemblage, suggesting that this material may be locally rare or exotic.

Microblades and debitage related to microblade production were present but not abundant in the overall assemblage (11.0%). This indicates that bifacial production was significantly more important than microblade production. Further, debitage related to microblade production contains many exotic pieces (12.1%), including 15 pieces of obsidian, one of which was confidently sourced to Wiki Peak. The results of a Fisher's exact test that compared the

use of exotic and local materials in the production of bifacial and microblade technology indicated a statistically significant ($p = 0.01$) difference between the raw materials selected for biface and microblade reduction.

Other Lithic Technologies

Unifacial scrapers and fragments represent 20% of the overall tool assemblage. Of these, one is a side scraper while the vast majority appear to have been used as end scrapers. Additionally, one unifacial tool appears to be a composite end scraper and shaft straightener. Unifacial technology at Clearview was overwhelmingly produced on locally available raw materials such as gray chert and black chert, though there are two complete chalcedony end scrapers present in the assemblage. This indicates that bifacial technology and unifacial technology raw material use strategies were approximately equivalent at Clearview.

Complete pieces of debitage linked to unifacial production represented a very small part of the overall debitage assemblage (0.7%), in contrast to the number of unifacial tools and tool fragments recovered ($n = 11$) during excavations at Clearview. The low number of intact debitage related to unifacial production may relate to the difficulty of distinguishing between bifacial and unifacial debitage, particularly pressure flakes, and the short reduction sequence of unifacial tool technology (Esdale 2009). However, this may also indicate that unifacial tools were used at Clearview but not produced on the same scale as bifacial or microblade technology.

The assemblage contains three transverse burins and five diagnostic burin spalls that were generated through burination, or edge modification of a flake for use as a graver or scraper (Barton et al. 1996). While the three burins were made from locally abundant raw materials, the burin spalls comprised red and white cherts that may be less abundant in the region. With such a small sample, it is not possible to determine whether the difference in raw material selection for specific technologies is statistically significant. However, burin production certainly employed a variety of raw materials.

Finally, utilized and retouched flakes were likely used as expedient tools and represent 23.6% of the tools at Clearview. These may have been used as flake knives during the occupation of the site and were made on locally available raw materials, including rhyolite, black chert, and gray chert.




Figure 6. Lithic technology recovered from the excavated area of the site. Illustrated by Briana Doering.




Figure 7. Debitage clusters by functional type; one square = 1 m².

Spatial Analysis

To preliminarily identify spatial patterns in artifact use and manufacture at Clearview, the distribution of diagnostic tools and tool fragments was compiled in ArcGIS Desktop 10.6 (Fig. 6). A series of *k*-means cluster analyses of tools, raw materials, and complete debitage conducted in RStudio showed two likely activity areas within the central excavation block. The composition of the two activity areas was compared to determine whether any significant differences in raw material and tool types existed, which could relate to differences in tool production activities. The difference in number of debitage pieces related to microblade and bifacial production in the two activity areas is not statistically significant ($p = 0.49$), indicating that both tool types were produced in similar numbers in both areas (Fig. 7). Early and intermediate bifacial reduction also occurred in both loci with no significant differences in distribution ($p = 0.55$). However, late bifacial reduction occurred at a slightly higher rate than early reduction in the smaller southern cluster ($p = 0.044$).

In contrast, raw material use varied significantly between the two areas (Fig. 8). Specifically, the difference in the amount of black chert and rhyolite tools and debitage is statistically significant ($p < 0.001$) between the two clusters, as was the amount of black chert and gray chert tools and debitage ($p < 0.001$), and gray chert and rhyolite ($p < 0.001$). These data suggest that black chert and rhyolite were primarily used in the northern activity area, and gray chert was primarily used in the southern activity area (Fig. 8). However, local and nonlocal material distributions do not differ significantly ($p = 0.09$) between the two clusters, indicating that the production of tools on nonlocal materials took place in both activity areas.

TECHNOLOGICAL ORGANIZATION AT CLEARVIEW

The results of the comprehensive lithic analysis presented above provide several correlates of human behavior that suggest the North Lobe WRA event did not generate a sudden wide-scale shift in mobility, diet, and organization among central Alaskans. First, the high artifact class richness and diversity of raw materials at Clearview indicate that the site was used for a relatively long period of time and implies that the site likely served as a residential base, not a logistical hunting camp. Second, exotic raw materials recovered from Clearview, including Wiki Peak and likely Batza Tena obsidian, suggest that patterns of long-distance exchange and/or high regional mobility persisted after the North Lobe WRA event. Finally, the complex reduction sequence and diverse toolkit described above does not appear consistent with expectations for a resource-poor environment. Combined, the aspects of the lithic assemblage indicate that the North Lobe WRA event did not produce a sudden, drastic shift in behavior among central Alaskans.

The extensive excavations at Clearview failed to recover any faunal materials that could have shown how subsistence strategies may have shifted during this period. Archaeologists have associated the lack of faunal materials at Alaska archaeological sites with acidic soils, which are common in coniferous boreal forests (Ping et al. 2008; Yesner 2001). Previous research has shown that faunal remains are best preserved in neutral (pH = 7) or slightly alkaline (pH = 7.5–8) soil environments (Nicholson 1996). In contrast, acidic soils with a pH of 3.5–4.5 provide the worst environment for faunal preservation, though low soil




Figure 8. Debitage clusters by material type; one square = 1 m². Illustrated by Briana Doering.

pH can also limit the growth of destructive microbes and does not necessarily connote a poor preservation environment (Manifold 2012). However, low soil pH, along with the complete lack of faunal material at Clearview, suggests that any faunal remains deposited during the site's occupation have subsequently decomposed in these acidic soils. Nevertheless, Clearview offers a wealth of lithic data that can be related to other sites to reveal larger patterns in mobility and land use.

The lithic assemblage from Clearview is comparable to Northern Archaic residential camps in the vicinity, such as Banjo Lake. This neighboring site features a similar and complex reduction sequence, diverse raw materials, obsidian sourced to Batza Tena, and assorted tool types (Esdale et al. 2015). However, the reduction sequence at Banjo Lake is incomplete across raw material types, meaning that the assemblage does not contain artifacts that represent the full sequence of lithic production for any single material type. This suggests that tools were made or maintained on whatever raw materials were available. Additionally, the site features extensive evidence for late-stage bifacial reduction and retouch. In contrast, Clearview's assemblage represents the full sequence of lithic reduction in artifacts of each material type, and few late-stage bifacial reduction and retouch flakes were recovered during extensive excavations. This important contrast in intra-assemblage variability suggests that the residents of Clearview may have been preparing tools for use at a logistical camp rather than refining tools for immediate use (Binford 1979). However, more assemblage and subsistence data from this and other sites from the late Holocene are necessary to identify larger patterns in mobility and landscape use at this transitional period.

CONCLUSION

The broad range of lithic technology and raw materials present at Clearview indicate that a group of residentially mobile hunter-gatherers occupied the site for a relatively long period of time. As a peripheral upland site with extensive evidence for large mammal hunting, represented indirectly by large lanceolate projectile points and hide working tools, Clearview is comparable to a summer or fall camp, according to Potter's (2016:548) model of seasonal subsistence for the Northern Archaic tradition. The focus on intermediate lithic production suggests that Clearview's residents may have been transitioning from

high mobility to semisedentism, but additional excavations and analysis from other central Alaska sites from this period are necessary to conclusively reconstruct shifting late Holocene settlement patterns. The lithic assemblage associated with this single occupation provides critical information relating late Holocene subsistence, technology, and mobility that can be used to clarify broader archaeological understandings of human activity through Alaska's history.

The research presented here provides further evidence that materials excavated as part of larger public works projects have the potential to address some of the most interesting debates in Alaska anthropology. The lithic assemblage recovered from Clearview provides a critical lens on central Alaska life around a time of significant behavioral changes. Modeling patterns of site use and residential mobility using a typological assessment of debitage can further our understanding of hunter-gatherer mobility and subsistence in central Alaska. Additional research on lithic assemblages from the late Holocene can enhance our understanding of life at the cusp of a major behavioral transition.

ACKNOWLEDGEMENTS

This article and the analysis presented here represent the culmination of three years of field and lab work completed in cooperation with the Tanana Chiefs Conference, on whose traditional lands Clearview lies. U.S. Army Garrison Alaska provided access to Clearview (XMH-1303) on Fort Wainwright under ARPA Permit 2016-001. Several archaeologists from the Center for Environmental Management of Military Lands conducted fieldwork at Clearview from 2016 to 2018. The data presented here owes much to the leadership of CEMML crew chiefs Heather Hardy, Whitney McLaren, Joshua Lynch, and Kate Yeske, and the excavation expertise of crew members Pierce Bateman, John Blong, Nicolette Edwards, Michael Grooms, Hilary Hilmer, Julia Kenyon, Brooks Lawler, Kelly Meierotto, Katherine Mulliken, Robert Nethkin, Allie Peltó, and Dougless Skinner. Whitney McLaren also spent her Christmas Eve in the lab photographing the artifacts. Scott Shirar facilitated the loan of previously excavated artifacts from the Museum of the North. At the University of Michigan, Raven Garvey provided space for the analysis of these artifacts, which was further supported by the curatorial efforts of collections manager Lauren

Fuka and administrative assistant Amy Winchester. Raven Garvey, John Speth, and Lauren Pratt aided in the development of the theoretical model considered here. Caitlin Holloway's feedback on the structure and argumentation of this article, as well as Robert Bowman's comments on Clearview's stratigraphy, immensely improved the excavation results presented here. Finally, Elizabeth Noll provided expert and invaluable editorial guidance, and this manuscript benefited from several helpful argumentation and formatting suggestions made by two anonymous reviewers and the editors. Any errors are the sole responsibility of the authors.

REFERENCES

- Amick, Daniel S.
1999 Raw Material Variation in Folsom Stone Tool Assemblages and the Division of Labor in Hunter-Gatherer Societies. In *Folsom Lithic Technology: Explorations in Structure and Variation*, edited by Daniel S. Amick, pp. 169–187. International Monographs in Prehistory: Archaeology Series, no. 12. Ann Arbor.
- Anderson, Patricia M., Mary E. Edwards, and Linda B. Brubaker
2003 Results and Paleoclimate Implications of 35 Years of Paleoecological Research in Alaska. *Developments in Quaternary Sciences* 1:427–440.
- Andrews, Elizabeth F.
1975 Salcha: An Athapaskan Band of the Tanana River and Its Culture. Master's thesis, Department of Anthropology, University of Alaska Fairbanks.
- Andrefsky, William
1994 Raw-Material Availability and the Organization of Technology. *American Antiquity* 59(1):21–34.
2005 *Lithics: Macroscopic Approaches to Analysis*. 2nd ed., *Cambridge Manuals in Archaeology*. Cambridge University Press, Cambridge.
- Bamforth, Douglas B.
1986 Technological Efficiency and Tool Curation. *American Antiquity* 51(1):38–50.
- Bamforth, Douglas B., and Nyree Finlay
2008 Introduction: Archaeological Approaches to Lithic Production Skill and Craft Learning. *Journal of Archaeological Method and Theory* 15(1):1–27.
- Barton, C. Michael, Deborah I. Olszewski, and Nancy R. Coinman
1996 Beyond the Graver: Reconsidering Burin Function. *Journal of Field Archaeology* 23(1):111–125.
- Binford, Lewis R.
1979 Organization and Formation Processes: Looking at Curated Technologies. *Journal of Anthropological Research* 35(3):255–73.
1980 Willow Smoke and Dogs' Tails: Hunter-Gatherer Settlement Systems and Archaeological Site Formation. *American Antiquity* 45(1):4–20.
- Carlson, Eric S., Julie A. Esdale, and Joshua J. Lynch
2017 Archaeological Districts on Fort Wainwright. Center for Environmental Management of Military Lands, Colorado State University, Ft. Collins.
- Chatters, James C., and William C. Prentiss
2005 A Darwinian Macro-Evolutionary Perspective on the Development of Hunter-Gatherer Systems in Northwestern North America. *World Archaeology* 37(1):46–65.
- Coutouly, Yan Axel Gómez
2012 Pressure Microblade Industries in Pleistocene-Holocene Interior Alaska: Current Data and Discussions. In *The Emergence of Pressure Blade Making*, edited by Pierre M. Desroisiers, pp. 347–374. Springer, New York.
- Cowan, Frank L.
1999 Making Sense of Flake Scatters: Lithic Technological Strategies and Mobility. *American Antiquity* 64(4):593–607.
- DeBoer, Warren R.
1988 Subterranean Storage and the Organization of Surplus: The View from Eastern North America. *Southeastern Archaeology* 7(1):1–20.
- Doering, Briana N., Julie A. Esdale, and Senna Catenacci
2019 Clearview (XMH-01303): Results of 2016–2018 Archaeological Excavations. Center for Environmental Management of Military Lands, Colorado State University, Ft. Collins.
- Elston, Robert G., and P. Jeffrey Brantingham
2002 Microlithic Technology in Northern Asia: A Risk-Minimizing Strategy of the Late Paleolithic and Early Holocene. *Archaeological Papers of the American Anthropological Association* 12(1):103–116.

- Esdale, Julie A.
2009 Lithic Production Sequences and Toolkit Variability: Examples from the Middle Holocene, Northwest Alaska. Ph.D. dissertation, Department of Anthropology, Brown University, Providence.
- Esdale, Julie A., Aaron Robertson, and William Johnson
2015 Banjo Lake: A Middle Holocene Site in the Tanana Valley. *Alaska Journal of Anthropology* 13(1):35–56.
- Gallant, Alisa L., Emily F. Binnian, James M. Omernik, and Mark B. Shasby
1995 Ecoregions of Alaska [map]. Scale 1:5,000,000. United States Geological Survey Professional Paper 1567, Washington, D.C. <https://pubs.er.usgs.gov/publication/pp1567>
- Goebel, Ted, and Ben A. Potter
2016 First Traces: Late Pleistocene Human Settlement of the Arctic. In *The Oxford Handbook of the Prehistoric Arctic*, edited by T. Max Friesen and O.K. Mason, pp. 223–253. Oxford University Press, New York.
- Goebel, Ted, Bryan Hockett, Kenneth D. Adams, David Rhode, and Kelly Graf
2011 Climate, Environment, and Humans in North America's Great Basin During the Younger Dryas, 12,900–11,600 Calendar Years Ago. *Quaternary International* 242(2):479–501.
- Graf, Kelly E., and Nancy H. Bigelow
2011 Human Response to Climate During the Younger Dryas Chronozone in Central Alaska. *Quaternary International* 242:434–51.
- Habu, Junko
2002 Jomon Collectors and Foragers: Regional Interactions and Long-Term Changes in Settlement Systems among Prehistoric Hunter-Gatherers in Japan. In *Beyond Foraging and Collecting: Evolutionary Change in Hunter-Gatherer Settlement Systems*, edited by Ben Fitzhugh and Junko Habu, pp. 53–72. Fundamental Issues in Archaeology. Kluwer Academic, Plenum Publishers, New York.
- Hare, P. Gregory, Sheila Greer, Ruth Gotthardt, Richard Farnell, Vandy Bowyer, Charles Schweger, and Diane Strand
2004 Ethnographic and Archaeological Investigations of Alpine Ice Patches in Southwest Yukon. *Arctic* 57(3):260–272.
- Helmer, James W., S. Van Dyke, and F. J. Kense, editors
1977 *Problems in the Prehistory of the North American Subarctic: The Athapaskan Question*. Archaeological Association of the University of Calgary, Alberta.
- Holmes, Charles E.
2008 The Taiga Period: Holocene Archaeology of the Northern Boreal Forest, Alaska. *Alaska Journal of Anthropology* 6(1–2):69–81.
- Jensen, Britta J.L., Sean Pyne-O'Donnell, Gill Plunkett, Duane G. Froese, Paul D.M. Hughes, Michael Sigl, Joseph R. McConnell, Matthew J. Amesbury, Paul G. Blackwell, and Christel van den Bogaard
2014 Transatlantic Distribution of the Alaskan White River Ash. *Geology* 42(10):875–878.
- Kaufman, Darrell S., Thomas A. Ager, N. John Anderson, Patricia M. Anderson, John T. Andrews, Pat J. Bartlein, Linda B. Brubaker, Larry L. Coats, Les C. Cwynar, and Mathieu L. Duvall
2004 Holocene Thermal Maximum in the Western Arctic (0–180 W). *Quaternary Science Reviews* 23(5–6):529–560.
- Kristensen, Todd J., Thomas D. Andrews, Glen MacKay, Ruth Gotthardt, Sean C. Lynch, John M. Duke, Andrew J. Locock, and John W. Ives
2019 Identifying and Sourcing Pyrometamorphic Artifacts: Clinker in Subarctic North America and the Hunter-Gatherer Response to a Late Holocene Volcanic Eruption. *Journal of Archaeological Science: Reports* 23:773–790.
- Kuhn, Tyler S., Keri A. McFarlane, Pamela Groves, Arne Ø. Mooers, and Beth Shapiro
2010 Modern and Ancient DNA Reveal Recent Partial Replacement of Caribou in the Southwest Yukon. *Molecular Ecology* 19(7):1312–1323.
- Lynch, Joshua J., Ted Goebel, Kelly Graf, Jeffrey T. Rasic, and Steven C. Kuehn
2018 Archaeology of the Uppermost Tanana Basin: Results of a Survey of the Nabesna and Chisana Rivers, East-Central Alaska. *Alaska Journal of Anthropology* 16(1):21–43.
- Manifold, Bernadette M.
2012 Intrinsic and Extrinsic Factors Involved in the Preservation of Non-Adult Skeletal Remains in Archaeology and Forensic Science. *Bulletin of the International Association for Paleodontology* 6(2):51–69.

- Moodie, D. Wayne, A.J. W. Catchpole, and Kerry Abel
1992 Northern Athapaskan Oral Traditions and the White River Volcano. *Ethnohistory* 39(2):148–171.
- Mullen, Patrick O.
2012 An Archaeological Test of the Effects of the White River Ash Eruptions. *Arctic Anthropology* 49(1):35–44.
- Nicholson, Rebecca A.
1996 Bone Degradation, Burial Medium and Species Representation: Debunking the Myths, an Experiment-Based Approach. *Journal of Archaeological Science* 23(4):513–533.
- Osgood, Cornelius B.
1937 *The Ethnography of the Tanaina*. Yale University Publications in Anthropology no. 16. Yale University, New Haven.
- Ostahowski, Brian E., and Robert L. Kelly
2014 Alm Rockshelter Lithic Debitage Analysis: Implications for Hunter-Gatherer Mobility Strategies in the Big Horn Mountains, Wyoming. In *Lithics in the West: Using Lithic Analysis to Solve Archaeological Problems in Western North America*, edited by Douglas H. MacDonald, William Andrefsky, and Pei-Lin Yu, pp. 120–141. University of Montana Press, Missoula.
- Péwé, Troy L.
1975 *Quaternary Geology of Alaska*. Geological Survey Professional Paper 835. U.S. Government Printing Office, Washington, D.C.
- Ping, C.L., G.J. Michaelson, J.M. Kimble, V.E. Romanovsky, Y.L. Shur, D.K. Swanson, and D.A. Walker
2008 Cryogenesis and Soil Formation along a Bioclimate Gradient in Arctic North America. *Journal of Geophysical Research: Biogeosciences* 113(G3).
- Plaskett, David C.
1977 The Nenana River Gorge Site: A Late Prehistoric Athapaskan Campsite in Central Alaska. Master's thesis, Department of Anthropology, University of Alaska Fairbanks.
- Potter, Ben A.
2008 Exploratory Models of Intersite Variability in Mid to Late Holocene Central Alaska. *Arctic* 61(4):407–425.
2011 Late Pleistocene and Early Holocene Assemblage Variability in Central Alaska. In *From the Yenisei to the Yukon: Interpreting Lithic Assemblage Variability in Late Pleistocene/Early Holocene Beringia*, edited by F. Ted Goebel and Ian Buvitt, pp. 215–233. Texas A&M University Press, College Station.
- 2016 Holocene Prehistory of the Northwestern Subarctic. In *The Oxford Handbook of the Prehistoric Arctic*, edited by T. Max Friesen and O.K. Mason, pp. 537–562. Oxford University Press, New York.
- Rainey, Froelich
1940 Archaeological Investigation in Central Alaska. *American Antiquity* 5(4):299–308.
- Reger, R., D. Stevens, and D. Solie
2008 Surficial Geology of the Alaska Highway Corridor, Delta Junction to Dot Lake. In *Preliminary Interpretive Report*. Alaska Division of Geological and Geophysical Surveys, Fairbanks.
- Reuther, Joshua D., N.S. Slobodina, J.T. Rasic, John P. Cook, and Robert J. Speakman
2011 Gaining Momentum: Late Pleistocene and Early Holocene Archaeological Obsidian Source Studies in Interior and Northeastern Beringia. In *From the Yenisei to the Yukon: Interpreting Lithic Assemblage Variability in Late Pleistocene/Early Holocene Beringia*, edited by Ted Goebel and Ian Buvitt, pp. 270–286. Texas A&M University Press, College Station.
- Robertson, Aaron C., Julie A. Esdale, James Quinn, Heather D. Hardy, and Vanessa Aziz
2013 Archaeological Data Recovery: Battle Area Complex, Donnelly Training Area, Fort Wainwright, Alaska 2009. Center for Environmental Management of Military Lands, Colorado State University, Ft. Collins.
- Robertson, Aaron C., Julie A. Esdale, Molly Proue, Scott Shirar, C. Kanani Paraso, and Christopher Ciancibelli
2009 Archaeological Survey and Mitigation: Donnelly Training Area, Fort Wainwright, Alaska 2007. Center for Environmental Management of Military Lands, Colorado State University, Ft. Collins.
- Robertson, Aaron C., Molly Proue, Patrick Hall, Scott Shirar, and C. Kanani Paraso
2007 Archaeological Survey, Evaluation, and Mitigation: Donnelly Training Area, Fort Wainwright, Alaska 2006. Center for Environmental Management of Military Lands, Colorado State University, Ft. Collins.
- Shinkwin, Anne D.
1979 Dakah De'nin's Village and the Dixthada Site: A Contribution to Northern Athapaskan Prehis-

- tory. *Musée National de l'Homme. Collection Mercure. Commission Archéologique du Canada. Publications d'Archéologie. Dossier Ottawa* (91):1–197.
- Shott, Michael J.
1986 Technological Organization and Settlement Mobility: An Ethnographic Examination. *Journal of Anthropological Research* 42(1):15–52.
- Stuiver, M., P. J. Reimer, and R. W. Reimer
2019 Calib 7.1. <http://calib.org/>
- Surovell, Todd A.
2009 *Toward a Behavioral Ecology of Lithic Technology: Cases from Paleoindian Archaeology*. University of Arizona Press, Tucson.
- Wandsnider, LuAnn
1996 Describing and Comparing Archaeological Spatial Structures. *Journal of Archaeological Method and Theory* 3(4):319–384.
- Workman, William B.
1979 The Significance of Volcanism in the Prehistory of Subarctic Northwest North America. In *Volcanic Activity and Human Ecology*, edited by Payson Sheets and Donald Grayson, pp. 339–371. Academic Press, New York.
- Yesner, David R.
1996 Human Adaptation at the Pleistocene-Holocene Boundary (circa 13,000–8,000 BP) in Eastern Beringia. In *Humans at the End of the Ice Age: The Archaeology of the Pleistocene-Holocene Transition*, edited by L. G. Straus, Berit V. Eriksen, J. M. Erlandson, and David R. Yesner, pp. 255–276. Plenum Press, New York.
- 2001 Human Dispersal into Interior Alaska: Antecedent Conditions, Mode of Colonization, and Adaptations. *Quaternary Science Reviews* 20(1–3):315–327.