

Informed Consent for SkinVive™ by Juvéderm®

Patient Name (printed)

British Columbia Botox® Clinics, by Dr. Ward

SkinVive™ by Juvéderm® is a hyaluronic acid (HA) injectable treatment that delivers long-lasting, deep hydration for up to 9 months following a single treatment. The HA molecule itself can attract and bind more than 1000 times its weight in water. It may also increase expression of water and glycerol transportation proteins (e.g., Aquaporin-3). Together, this micro-injection treatment noticeably rejuvenates and hydrates skin 1-month after treatment.

The following safety information comes directly from the product monograph:

- Inflammatory reactions (redness, oedema, erythema, etc.) which may be associated with itching and/or pain on pressure and/or paresthesia, occurring after the injection. These reactions may last for a week.
- Haematomas.
- Induration or nodules at the injection site.
- Staining or discolouration of the injection site might be observed, especially when HA dermal filler is injected too superficially and/or in thin skin (Tyndall effect).
- Poor effect or weak filling effect.
- Rare but serious adverse events associated with intravascular injection of dermal fillers in the face and tissue compression have been reported and include temporary or permanent vision impairment, blindness, cerebral ischemia or cerebral hemorrhage, leading to stroke, skin necrosis and damage to underlying structures. Immediately stop the injection if a patient exhibits any of the following symptoms, including changes in the vision, signs of stroke, blanching of the skin or unusual pain during or shortly after the procedure. Patients should receive prompt medical attention and possibly evaluation by an appropriate medical practitioner specialist should an intravascular injection occur. Abscesses, granuloma and immediate or delayed hypersensitivity after hyaluronic acid and/or lidocaine injections have also been reported. It is therefore advisable to take these potential risks into account.
- Patients must report inflammatory reactions which persist for more than one week, or any other side effect which develops, to their medical practitioner as soon as possible. The medical practitioner should use an appropriate treatment.
- Any other undesirable side effects associated with injection of SKINVIVE™ by JUVÉDERM® must be reported to the distributor and/or to the manufacturer.

Caution should be used in patients with a history of herpes simplex (cold sores), bleeding disorders, autoimmune diseases, or those on immunosuppressive therapy.

All dermal fillers must be avoided if a patient is pregnant or breastfeeding or under the age of 18.

All dermal fillers must be avoided if a patient has a history of anaphylaxis or severe allergy / sensitivity to these products, including lidocaine.

Notify the doctor if you have had any dermal fillers in the past (including type and location). Adding different types of fillers to the same location can result in granulomas / scarring (this is not true when adding different Juvéderm® products together).

Small bumps may be noticed at each injection site certainly for the first 48 hours, and may even last (in part) up to 14 days. Scarring at injection sites, especially in those with a history of keloid scars, is also a rare risk.

Aftercare: Minor swelling may be improved with cool compresses. Patients should avoid the sun, smoking and alcohol for 2-3 days after having fillers injected. Full and final effects are expected around the 1-month mark and last up to 9 months.

Patients must consent to photos being taken before and after any filler procedure (this will only be used for medical documentation in the Electronic Medical Record).

Treatment Areas and Costs (not covered by MSP)

Cost is \$350 per syringe. Most facial treatments take 2-3 syringes. Neck treatments take 2 syringes.

The practice of medicine is not an exact science and as such, no guarantees can be given concerning the results of any cosmetic procedure performed in our clinics.

The clinic will not provide a refund or free re-injection of any filler service performed, regardless of the cosmetic or medical outcome.

Patient Signature

Date