

I'm not robot  reCAPTCHA

**I am not robot!**

## Cultivation of aerobic and anaerobic bacteria pdf. What is cultivation of bacteria.

Understanding the role of our microbiota in health and disease has long been hampered by the strict growth requirements of many of its constituent members. Underpinning modern day investigations into the vast complexity and functions of the human microbiota are fundamental methodologies to culture anaerobic bacteria outside their natural environment. From the rudimentary oxygen-free culture methods in the era of Pasteur, and subsequent advances in surface culture in the early twentieth century, the mid-1900s saw a substantial expansion and refinement of anaerobic culture techniques, largely due to the pioneering work of Robert. E. Hungate. In a 1944 study of cellulose-degrading microorganisms in the bovine rumen, his revolutionary roll-tube approach enabled the successful culture of *Clostridium cellobioparus* and, in 1950, he published a complete description of his technique. The protocol used rubber-stoppered tubes of boiled culture medium with cellulose agar, through which anoxic gas was bubbled to remove any remaining oxygen. Firstly, passing this gas through a column of hot, reduced copper wire excluded any oxygen from the gas itself, and the subsequent addition of a reducing agent to the medium removed residual traces of oxygen. Rolling tubes under cold water produced a thin layer of solid agarose medium, and for the first time, anaerobiosis was maintained throughout manipulations using a constant flow of anoxic gas. The method, now known as 'the Hungate technique', is still in use to this day. Diagram of a Hungate roll tube. Credit: S. Fenwick / Springer Nature Limited Several modifications later emerged, such as the VPI (Virginia Polytech Institute) method for larger-scale culture introduced by Moore in 1966, using pre-reduced medium and pre-hardened roll tubes. Hungate also made adaptations to culture methanogens, the strictest of anaerobes, reported in 1969. Others, such as Spears and Freter in 1967, similarly recognised the importance of continuously avoiding any exposure to oxygen, yet the Hungate technique was still more efficient and enabled a wealth of anaerobes that had not grown previously in surface cultures to be isolated for further study. Alternative approaches used today were also launched in the mid-late 1960s, namely the GasPak and the anaerobic glove-box. The former, a self-contained combustion jar system, quickly made surface culture of anaerobic microorganisms accessible to more laboratories. The glove-box, a sealed chamber with attached gloves, filled with anoxic gases, was also a popular choice, simplifying equipment and procedures for oxygen-free culture. As well as apparatus to create an oxygen-free environment, culturing anaerobes requires appropriate media, which must have a low oxidation-reduction potential, as well as the substrates obtained by microorganisms in their natural habitat. Many researchers working on *Bacteroides* species were instrumental in determining the requirements of specific anaerobic microorganisms, and a recent breakthrough in media composition (the inclusion of antioxidants) has since permitted the aerobic growth of anaerobic bacteria. Moving into the twenty-first century, the advent of metagenomics revealed that the majority of environmental microbial biodiversity remained uncultured, inspiring a rebirth of culture techniques. Recent culture-dependent efforts to characterize the human microbiota (see Milestone 19) utilised dilution culturing and culminated in the development of culturomics; a high-throughput methodology using hundreds of different culture conditions, prolonged incubations, and matrix-assisted laser desorption/ionization-time of flight (MALDI-TOF) spectrometry, combined with 16S ribosomal RNA gene sequencing for the rapid identification of a great number of previously uncultured gut bacteria. With a large proportion of the human microbiota requiring oxygen-free growth conditions, early breakthroughs in anaerobic culture were crucial in enabling more of our microbiota to be isolated and classified, and for their metabolism, distribution and roles within the microbiota to be studied.



**Aerobic and anaerobic enrichment cultures highlight the pivotal role of facultative anaerobes in soil hydrocarbon degradation**

Errol D. Cason, Jan-G Vermeulen, Walter J. Müller, Esta van Heerden & Angel Valverde

To cite this article: Errol D. Cason, Jan-G Vermeulen, Walter J. Müller, Esta van Heerden & Angel Valverde (2019) Aerobic and anaerobic enrichment cultures highlight the pivotal role of facultative anaerobes in soil hydrocarbon degradation, *Journal of Environmental Science and Health, Part A*, DOI: 10.1080/10934529.2018.1558902

To link to this article: <https://doi.org/10.1080/10934529.2018.1558902>

View supplementary material

Published online: 24 Jan 2019


Submit your article to this journal

View Crossmark data


Full Terms & Conditions of Access and Use can be found at <http://www.tandfonline.com/action/journalInformation?journalCode=hesa20>

The protocol used rubber-stoppered tubes of boiled culture medium with cellulose agar, through which anoxic gas was bubbled to remove any remaining oxygen. Firstly, passing this gas through a column of hot, reduced copper wire excluded any oxygen from the gas itself, and the subsequent addition of a reducing agent to the medium removed residual traces of oxygen. Rolling tubes under cold water produced a thin layer of solid agarose medium, and for the first time, anaerobiosis was maintained throughout manipulations using a constant flow of anoxic gas. The method, now known as 'the Hungate technique', is still in use to this day. Diagram of a Hungate roll tube. Credit: S. Fenwick / Springer Nature Limited Several modifications later emerged, such as the VPI (Virginia Polytech Institute) method for larger-scale culture introduced by Moore in 1966, using pre-reduced medium and pre-hardened roll tubes. Hungate also made adaptations to culture methanogens, the strictest of anaerobes, reported in 1969. Others, such as Spears and Freter in 1967, similarly recognised the importance of continuously avoiding any exposure to oxygen, yet the Hungate technique was still more efficient and enabled a wealth of anaerobes that had not grown previously in surface cultures to be isolated for further study. Alternative approaches used today were also launched in the mid-late 1960s, namely the GasPak and the anaerobic glove-box. The former, a self-contained combustion jar system, quickly made surface culture of anaerobic microorganisms accessible to more laboratories. The glove-box, a sealed chamber with attached gloves, filled with anoxic gases, was also a popular choice, simplifying equipment and procedures for oxygen-free culture. As well as apparatus to create an oxygen-free environment, culturing anaerobes requires appropriate media, which must have a low oxidation-reduction potential, as well as the substrates obtained by microorganisms in their natural habitat. Many researchers working on *Bacteroides* species were instrumental in determining the requirements of specific anaerobic microorganisms, and a recent breakthrough in media composition (the inclusion of antioxidants) has since permitted the aerobic growth of anaerobic bacteria. Moving into the twenty-first century, the advent of metagenomics revealed that the majority of environmental microbial biodiversity remained uncultured, inspiring a rebirth of culture techniques. Recent culture-dependent efforts to characterize the human microbiota (see Milestone 19) utilised dilution culturing and culminated in the development of culturomics; a high-throughput methodology using hundreds of different culture conditions, prolonged incubations, and matrix-assisted laser desorption/ionization-time of flight (MALDI-TOF) spectrometry, combined with 16S ribosomal RNA gene sequencing for the rapid identification of a great number of previously uncultured gut bacteria. With a large proportion of the human microbiota requiring oxygen-free growth conditions, early breakthroughs in anaerobic culture were crucial in enabling more of our microbiota to be isolated and classified, and for their metabolism, distribution and roles within the microbiota to be studied. Initial methodologies paved the way for higher-throughput technologies that provide vital insights about the functions of the bacteria inhabiting the human body, and their effects on the human host. Now, with our understanding of the importance of the gut microbiota in human health advancing by the day, we are even more indebted to these early researchers and their innovations enabling the culture of anaerobes. Hall, I. C. Practical methods in the purification of obligate anaerobes.

## Cultivation of Anaerobic Bacteria



Underpinning modern day investigations into the vast complexity and functions of the human microbiota are fundamental methodologies to culture anaerobic bacteria outside their natural environment. From the rudimentary oxygen-free culture methods in the era of Pasteur, and subsequent advances in surface culture in the early twentieth century, the mid-1900s saw a substantial expansion and refinement of anaerobic culture techniques, largely due to the pioneering work of Robert. E. Hungate. In a 1944 study of cellulose-degrading microorganisms in the bovine rumen, his revolutionary roll-tube approach enabled the successful culture of *Clostridium cellobioparus* and, in 1950, he published a complete description of his technique. The protocol used rubber-stoppered tubes of boiled culture medium with cellulose agar, through which anoxic gas was bubbled to remove any remaining oxygen. Rolling tubes under cold water produced a thin layer of solid agarose medium, and for the first time, anaerobiosis was maintained throughout manipulations using a constant flow of anoxic gas. The method, now known as 'the Hungate technique', is still in use to this day. Diagram of a Hungate roll tube. Credit: S. Fenwick / Springer Nature Limited Several modifications later emerged, such as the VPI (Virginia Polytech Institute) method for larger-scale culture introduced by Moore in 1966, using pre-reduced medium and pre-hardened roll tubes. Hungate also made adaptations to culture methanogens, the strictest of anaerobes, reported in 1969. Others, such as Spears and Freter in 1967, similarly recognised the importance of continuously avoiding any exposure to oxygen, yet the Hungate technique was still more efficient and enabled a wealth of anaerobes that had not grown previously in surface cultures to be isolated for further study. Alternative approaches used today were also launched in the mid-late 1960s, namely the GasPak and the anaerobic glove-box. The former, a self-contained combustion jar system, quickly made surface culture of anaerobic microorganisms accessible to more laboratories. The glove-box, a sealed chamber with attached gloves, filled with anoxic gases, was also a popular choice, simplifying equipment and procedures for oxygen-free culture. As well as apparatus to create an oxygen-free environment, culturing anaerobes requires appropriate media, which must have a low oxidation-reduction potential, as well as the substrates obtained by microorganisms in their natural habitat. Many researchers working on *Bacteroides* species were instrumental in determining the requirements of specific anaerobic microorganisms, and a recent breakthrough in media composition (the inclusion of antioxidants) has since permitted the aerobic growth of anaerobic bacteria. Moving into the twenty-first century, the advent of metagenomics revealed that the majority of environmental microbial biodiversity remained uncultured, inspiring a rebirth of culture techniques. Recent culture-dependent efforts to characterize the human microbiota (see Milestone 19) utilised dilution culturing and culminated in the development of culturomics; a high-throughput methodology using hundreds of different culture conditions, prolonged incubations, and matrix-assisted laser desorption/ionization-time of flight (MALDI-TOF) spectrometry, combined with 16S ribosomal RNA gene sequencing for the rapid identification of a great number of previously uncultured gut bacteria. With a large proportion of the human microbiota requiring oxygen-free growth conditions, early breakthroughs in anaerobic culture were crucial in enabling more of our microbiota to be isolated and classified, and for their metabolism, distribution and roles within the microbiota to be studied. Initial methodologies paved the way for higher-throughput technologies that provide vital insights about the functions of the bacteria inhabiting the human body, and their effects on the human host. Now, with our understanding of the importance of the gut microbiota in human health advancing by the day, we are even more indebted to these early researchers and their innovations enabling the culture of anaerobes. Hall, I. C. Practical methods in the purification of obligate anaerobes. J. Infect. Dis. 27, 576-590 (1920). Hall, I. C. Differentiation and identification of the sporulating anaerobes.




The protocol used rubber-stoppered tubes of boiled culture medium with cellulose agar, through which anoxic gas was bubbled to remove any remaining oxygen. Firstly, passing this gas through a column of hot, reduced copper wire excluded any oxygen from the gas itself, and the subsequent addition of a reducing agent to the medium removed residual traces of oxygen. Rolling tubes under cold water produced a thin layer of solid agarose medium, and for the first time, anaerobiosis was maintained throughout manipulations using a constant flow of anoxic gas. The method, now known as 'the Hungate technique', is still in use to this day. Diagram of a Hungate roll tube. Credit: S. Fenwick / Springer Nature Limited Several modifications later emerged, such as the VPI (Virginia Polytech Institute) method for larger-scale culture introduced by Moore in 1966, using pre-reduced medium and pre-hardened roll tubes. Hungate also made adaptations to culture methanogens, the strictest of anaerobes, reported in 1969. Others, such as Spears and Freter in 1967, similarly recognised the importance of continuously avoiding any exposure to oxygen, yet the Hungate technique was still more efficient and enabled a wealth of anaerobes that had not grown previously in surface cultures to be isolated for further study. Alternative approaches used today were also launched in the mid-late 1960s, namely the GasPak and the anaerobic glove-box. The former, a self-contained combustion jar system, quickly made surface culture of anaerobic microorganisms accessible to more laboratories. The glove-box, a sealed chamber with attached gloves, filled with anoxic gases, was also a popular choice, simplifying equipment and procedures for oxygen-

free culture.As well as apparatus to create an oxygen-free environment, culturing anaerobes requires appropriate media, which have a low oxidation-reduction potential, as well as the substrates obtained by microorganisms in their natural habitat. Many researchers working on Bacteroides species instrumental in determining the requirements of specific anaerobic microorganisms, and a recent breakthrough in media composition (the inclusion of antioxidants) has since permitted the aerobic growth of anaerobic bacteria.Moving into the twenty-first century, the advent of metagenomics revealed that the majority of environmental microbial biodiversity remained uncultured, inspiring a rebirth of culture techniques. Recent culture-dependent efforts to characterize the human microbiota (see Milestone 19) utilised dilution culturing and culminated in the development of culturomics; a high-throughput methodology using hundreds of different culture conditions, prolonged incubations, and matrix-assisted laser desorption/ionization-time-of flight (MALDI-TOF) spectrometry, combined with 16S ribosomal RNA gene sequencing for the rapid identification of a great number of previously uncultured gut bacteria.With a large proportion of the human microbiota requiring oxygen-free growth conditions, early breakthroughs in anaerobic culture were crucial in enabling more of our microbiota to be isolated and classified, and for their metabolism, distribution and roles within the microbiota to be studied. Initial methodologies paved the way for higher-throughput technologies that provide vital insights about the functions of the bacteria inhabiting the human body, and their effects on the human host. Now, with our understanding of the importance of the gut microbiota in human health advancing by the day, we are even more indebted to these early researchers and their innovations enabling the culture of anaerobes. Hall, I. C. Practical methods in the purification of obligate anaerobes. J. Infect. Dis. 27, 576–590 (1920). Hall, I. C. Differentiation and identification of the sporulating anaerobes. J. Infect.



The protocol used rubber-stoppered tubes of boiled culture medium with cellulose agar, through which anoxic gas was bubbled to remove any remaining oxygen. Firstly, passing this gas through a column of hot, reduced copper wire excluded any oxygen from the gas itself, and the subsequent addition of a reducing agent to the medium removed residual traces of oxygen. Rolling tubes under cold water produced a thin layer of solid agarose medium, and for the first time, anaerobiosis was maintained throughout manipulations using a constant flow of anoxic gas.

## Anaerobic Culture Methods



The protocol used rubber-stoppered tubes of boiled culture medium with cellulose agar, through which anoxic gas was bubbled to remove any remaining oxygen. Firstly, passing this gas through a column of hot, reduced copper wire excluded any oxygen from the gas itself, and the subsequent addition of a reducing agent to the medium removed residual traces of oxygen. Rolling tubes under cold water produced a thin layer of solid agarose medium, and for the first time, anaerobiosis was maintained throughout manipulations using a constant flow of anoxic gas. The method, now known as ‘the Hungate technique’, is still in use to this day. Diagram of a Hungate roll tube.Credit: S. Fenwick / Springer Nature Limited Several modifications later emerged, such as the VPI (Virginia Polytech Institute) method for larger-scale culture introduced by Moore in 1966, using prerduced medium and prehardened roll tubes. Hungate also made adaptations to culture methanogens, the strictest of anaerobes, reported in 1969. Others, such as Spears and Freter in 1967, similarly recognised the importance of continuously avoiding any exposure to oxygen, yet the Hungate technique was still more efficient and enabled a wealth of anaerobes that had not grown previously in surface cultures to be isolated for further study.Alternative approaches used today were also launched in the mid-late 1960s, namely the GasPak and the anaerobic glove-box. The former, a self-contained combustion jar system, quickly made surface culture of anaerobic microorganisms accessible to more laboratories. The glove-box, a sealed chamber with attached gloves, filled with anoxic gases, was also a popular choice, simplifying equipment and procedures for oxygen-free culture.As well as apparatus to create an oxygen-free environment, culturing anaerobes requires appropriate media, which must have a low oxidation-reduction potential, as well as the substrates obtained by microorganisms in their natural habitat. Many researchers working on Bacteroides species were instrumental in determining the requirements of specific anaerobic microorganisms, and a recent breakthrough in media composition (the inclusion of antioxidants) has since permitted the aerobic growth of anaerobic bacteria.Moving into the twenty-first century, the advent of metagenomics revealed that the majority of environmental microbial biodiversity remained uncultured, inspiring a rebirth of culture techniques. Recent culture-dependent efforts to characterize the human microbiota (see Milestone 19) utilised dilution culturing and culminated in the development of culturomics; a high-throughput methodology using hundreds of different culture conditions, prolonged incubations, and matrix-assisted laser desorption/ionization-time-of flight (MALDI-TOF) spectrometry, combined with 16S ribosomal RNA gene sequencing for the rapid identification of a great number of previously uncultured gut bacteria.With a large proportion of the human microbiota requiring oxygen-free growth conditions, early breakthroughs in anaerobic culture were crucial in enabling more of our microbiota to be isolated and classified, and for their metabolism, distribution and roles within the microbiota to be studied. Initial methodologies paved the way for higher-throughput technologies that provide vital insights about the functions of the bacteria inhabiting the human body, and their effects on the human host. Now, with our understanding of the importance of the gut microbiota in human health advancing by the day, we are even more indebted to these early researchers and their innovations enabling the culture of anaerobes. Hall, I. C. Practical methods in the purification of obligate anaerobes. J. Infect. Dis. 27, 576–590 (1920). Hall, I. C. Differentiation and identification of the sporulating anaerobes.

The anaerobic mesophilic cellulolytic bacteria. Bacteriol. Rev. 14, 1–49 (1950). Bryant, M. P. & Doetsch, R. N. Factors necessary for the growth of Bacteroides succinogenes in the volatile acid fraction of rumen fluid. Science 120, 944–945 (1954). Moore W. E. C. Techniques for routine culture of fastidious anaerobes. Intern. J. Syst. Bacteriol. 16, 173–190 (1966). Brewer, J. H. & Allgeier, D. L. Safe self-contained carbon dioxide-hydrogen anaerobic system. Appl. Microbiol. 14, 985–988 (1966). Spears R. W. & Freter, R. Improved isolation of anaerobic bacteria from the mouse cecum by maintaining continuous strict anaerobiosis.

Proc. Soc. Exp. Biol. Med. 124, 903–909 (1967). Drasar, B. S. Cultivation of anaerobic intestinal bacteria. J. Pathol. Bacteriol. 94, 417–427 (1967). Savage, D. C., Dubos, R. & Schaedler, R. W. The gastrointestinal epithelium and its autochthonous bacterial flora. J. Exp. Med. 127, 67–76 (1968). Aranki, A. et al. Isolation of anaerobic bacteria from human gingiva and mouse cecum by means of a simplified glove box procedure. Appl. Microbiol. 17, 568–576 (1969). Hungate R.

E. Chapter IV: A roll tube method for cultivation of strict anaerobes. Method. Microbiol. 3, 117–132 (1969). Sutter, V. L. & Finegold, S. M. Antibiotic disc susceptibility tests for rapid presumptive identification of gram-negative anaerobic bacteria. Appl. Microbiol. 21, 13–20 (1970). Sonnenwirth, A. C. Evolution of anaerobic methodology. Am. J. Clin. Nutr. 25, 1295–1298 (1972). Holdeman, L. V. & Moore, W. E. C. Roll-tube techniques for anaerobic bacteria. Am. J. Clin. Nutr. 25, 1314–1317 (1972). Salyers, A. A. Energy sources of major intestinal fermentative anaerobes. Am. J. Clin. Nutr. 32, 158–163 (1979). Goodman, A. L.

et al. Extensive personal human gut microbiota culture collections characterized and manipulated in gnotobiotic mice. Proc. Natl Acad. Sci. USA 108, 6252–6257 (2011). Lagier, J. C. et al. Microbial culturomics: paradigm shift in the human gut microbiome study. Clin. Microbiol. Infect. 18, 1185–1193 (2012). Dione, N. et al. A quasi-universal medium to break the aerobic/anaerobic bacterial culture dichotomy in clinical microbiology. Clin. Microbiol. Infect. 22, 53–58 (2016). Lagier, J. C. et al. Culturing the human microbiota and culturomics. Nat. Rev. Micro. 16, 540–550 (2018). Hungate, R. E. (1969) A roll tube method for cultivation of strict anaerobes, in Methods in Microbiology, Vol. 3b (Norris, J. R. and Ribbons, D. W., ed.), Academic, New York, pp. 117–132. Google Scholar

Bryant, M. P (1972) Commentary on the Hungate technique for culture of anaerobic bacteria. Am. J. Clin. Nutr. 25, 1324–1328.CAS Google Scholar Balch, W. E., Fox, G. E., Magrum, L. J., Woese, C. R., and Wolfe, R. S. (1979) Methanogens: reevaluation of a unique biological group. Microbiol. Mol. Biol. Rev. 43, 260–296.CAS Google Scholar Kaar W.E., Holtzapfle M.T. (2000) The anaerobic mesophilic cellulolytic bacteria. Bacteriol. Rev. 14, 1–49.CAS Google Scholar Monod, J. (1950) La technique de culture continue, theorie et applications. Ann. Inst. Pasteur 79, 390–410.CAS Google Scholar Novick, A. and Szilard, L (1950) Description of the chemostat. Science 112, 715–716.CrossRef CAS Google Scholar Herbert, D. R., Elsworth, R., and Telling R. C. (1956) The continuous culture of bacteria: a theoretical and experimental study. J. Gen. Microbiol.

14, 601–622.CrossRef CAS Google Scholar Pirt, S. J. (1975) Principles of Microbe and Cell Cultivation. Blackwell, Oxford. Google Scholar Kistner, A. and Kornelius, J. H. (1990) A small-scale, 3- vessel, continuous culture system for quantitative studies of plant fiber degradation by anaerobic bacteria. J. Microbiol. Meth. 12, 173–182.CrossRef Google Scholar Pavlostathis, S. G., Miller, T. L., and Wolin, M. J. (1988) Fermentation of insoluble cellulose by continuous cultures of Ruminococcus albus. Appl. Environ. Microbiol. 54, 2655–2659.CAS Google Scholar Fukushima, R. S., Weimer, P. J., and Kunz, D. A. (2002) Photocatalytic interaction of resazurin N-oxide with cysteine optimizes preparation of anaerobic culture media. Anaerobe 8, 29–34.CrossRef CAS Google Scholar Jones, G. A. and Pickard, M. D. (1980) Effect of titanium (III) citrate as reducing agent on growth of rumen bacteria. Appl. Environ. Microbiol. 39, 1144–1147.CAS Google Scholar Page 2 91k Accesses 736 Citations 20 Altmetric Page 3M Foster, N., Wyman, C., Dale, B., Elandor, R., Lee, Y.Y., Holtzapfle, M., Ladisch, M., (2005) Features of promising technologies for pretreatment of lignocellulosic biomass. Bioresource Technology 96, 673–686 Google Scholar Sun, Y., Cheng, J. (2002). Hydrolysis of lignocellulosic materials for ethanol production: a review. Bioresource Technology 83, 1–11CrossRef CAS Google Scholar O'Dwyer, J., Zhu, L., Granda, C.B., Holtzapfle, M.T. (2007). Enzymatic hydrolysis of lime-pretreated corn stover and investigation of the HCH-1 Model: Inhibition pattern, degree of inhibition, validity of simplified HCH-1 Model.

Bioresource Technology 98, 2969–2977CrossRef Google Scholar Sehoon, K., Holtzapfle, M.T. (2006). Effect of structural features on enzyme digestibility of corn stover. Bioresource Technology 97, 583–591CrossRef Google Scholar Chang, V.S., Holtzapfle, M.T. (2000). Fundamental factors affecting Biomass Enzymatic Reactivity. Applied Biochemistry and Biotechnology 84–86, 1–37 Google Scholar Sehoon, K., Holtzapfle, M.T. (2005). Lime pretreatment and enzymatic hydrolysis of corn stover. Bioresource Technology 96, 1994–2006CrossRef Google Scholar Chang, V.S., Nagwani, M., Holtzapfle, M.T. (1997). Lime pretreatment of switchgrass. Applied Biochemistry and Biotechnology 63–65, 3–19CrossRef Google Scholar Chang, V.S., Nagwani, M., Kim, C.H., Holtzapfle, M.T. (2001). Oxidative lime pretreatment of high-lignin biomass.

Applied Biochemistry and Biotechnology 94, 1–28CrossRef CAS Google Scholar Kaar W.E., Holtzapfle M.T. (2000). Using lime pretreatment to facilitate the enzyme hydrolysis of corn stover. Biomass and Bioenergy 18(3), 189–199CrossRef Google Scholar Granda, C.B. (2004). Sugarcane Juice Extraction and Preservation, and Long-Term Lime Pretreatment of Bagasse. PhD dissertation, Texas A&M University, College Station, TX Google Scholar Sierra, R. (2005). Long-Term Lime Pretreatment of Poplar Wood. Thesis. Texas A&M University.

Environ. Microbiol. 67:3077–3085. [PMC free article] [PubMed] [Google Scholar]13. Fisk, M. R., and S. J. Giovanni. 1999. Sources of nutrients and energy for a deep biosphere on Mars. J. Geophys. Res. 104:11805–11816. [Google Scholar]14. Hillmann, F., R. J. Fischer, F. Saint-Prix, L. Girbal, and H. Bahl. 2008. PerR acts as a switch for oxygen tolerance in the strict anaerobe Clostridium acetobutylicum. Mol. Microbiol. 68:848–860. [PubMed] [Google Scholar]15. Hino, S., and P. W. Wilson. 1957. Nitrogen fixation by a facultative Bacillus. J. Bacteriol. 57:403–408. [PMC free article] [PubMed] [Google Scholar]16. Horneck, G., H. Buckner, and G. Reitz. 1994. Long-term survival of bacterial spores in space. Adv. Space Res. 14:41–45. [PubMed] [Google Scholar]17. Huber, H., M. J. Hohn, R. Rachel, T. Fuchs, V. C. Wimmer, and K. O. Stetter. 2002. A new phylum of Archaea represented by a nanosized hyperthermophilic symbiont. Nature 417:63–67. [PubMed] [Google Scholar]18. Huber, R., T. A. Langworthy, H. König, M. Thomm, C. R. Woese, U. B. Sleytr, and K. O. Stetter. 1986. Thermotoga maritima sp. nov. represents a new genus of unique extremely thermophilic eubacteria growing up to 90°C. Arch. Microbiol. 144:324–333. [Google Scholar]19. Hungate, R. E. 1950. The anaerobic mesophilic cellulolytic bacteria. Bacteriol. Rev. 14:1–49. [PMC free article] [PubMed] [Google Scholar]20. Hungate, R. E. 1969. A roll tube method for cultivation of strict anaerobes, p. 117–132. In J. R. Norris and D. W. Ribbons (ed.), Methods in microbiology.

Academic Press, New York. NY.21. Kral, T. A., C. R. Bekkum, and C. P. McKay. 2004. Growth of methanogens on a Mars soil simulant. Orig. Life Evol. Biosph. 34:615–626. [PubMed] [Google Scholar]22. Küsel, K., C. Wagner, and H. L. Drake. 1999. Enumeration and metabolic product profiles of the anaerobic microflora in the mineral soil and litter of a beech forest. FEMS Microbiol. Ecol. 29:91–103. [Google Scholar]23. La Duc, M. T., W. Nicholson, R. Kern, and K. Venkateswaran. 2003. Microbial characterization of the Mars Odyssey spacecraft and its encapsulation facility. Environ. Microbiol. 67:977–985. [PMC free article] [PubMed] [Google Scholar]13. Fisk, M. R., and S. J. Giovanni. 1999. Sources of nutrients and energy for a deep biosphere on Mars. J. Geophys. Res. 104:11805–11816. [Google Scholar]14. Hillmann, F., R. J. Fischer, F. Saint-Prix, L. Girbal, and H. Bahl. 2008. PerR acts as a switch for oxygen tolerance in the strict anaerobe Clostridium acetobutylicum. Mol. Microbiol. 68:848–860. [PubMed] [Google Scholar]15. Hino, S., and P. W. Wilson. 1957. Nitrogen fixation by a facultative Bacillus. J. Bacteriol. 57:403–408. [PMC free article] [PubMed] [Google Scholar]16. Horneck, G., H. Buckner, and G. Reitz. 1994. Long-term survival of bacterial spores in space. Adv. Space Res. 14:41–45. [PubMed] [Google Scholar]17. Huber, H., M. J. Hohn, R. Rachel, T. Fuchs, V. C. Wimmer, and K. O. Stetter. 2002. A new phylum of Archaea represented by a nanosized hyperthermophilic symbiont. Nature 417:63–67. [PubMed] [Google Scholar]18. Huber, R., T. A. Langworthy, H. König, M. Thomm, C. R. Woese, U. B. Sleytr, and K. O. Stetter. 1986. Thermotoga maritima sp. nov. represents a new genus of unique extremely thermophilic eubacteria growing up to 90°C. Arch. Microbiol. 144:324–333. [Google Scholar]19. Hungate, R. E. 1950. The anaerobic mesophilic cellulolytic bacteria. Bacteriol. Rev. 14:1–49. [PMC free article] [PubMed] [Google Scholar]20. Hungate, R. E. 1969. A roll tube method for cultivation of strict anaerobes, p. 117–132. In J. R. Norris and D. W. Ribbons (ed.), Methods in microbiology.

Academic Press, New York. NY.21. Kral, T. A., C. R. Bekkum, and C. P. McKay. 2004. Growth of methanogens on a Mars soil simulant. Orig. Life Evol. Biosph. 34:615–626. [PubMed] [Google Scholar]22. Küsel, K., C. Wagner, and H. L. Drake. 1999. Enumeration and metabolic product profiles of the anaerobic microflora in the mineral soil and litter of a beech forest. FEMS Microbiol. Ecol. 29:91–103. [Google Scholar]23. La Duc, M. T., W. Nicholson, R. Kern, and K. Venkateswaran. 2003. Microbial characterization of the Mars Odyssey spacecraft and its encapsulation facility. Environ. Microbiol. 67:977–985. [PMC free article] [PubMed] [Google Scholar]13. Fisk, M. R., and S. J. Giovanni. 1999. Sources of nutrients and energy for a deep biosphere on Mars. J. Geophys. Res. 104:11805–11816. [Google Scholar]14. Hillmann, F., R. J. Fischer, F. Saint-Prix, L. Girbal, and H. Bahl. 2008. PerR acts as a switch for oxygen tolerance in the strict anaerobe Clostridium acetobutylicum. Mol. Microbiol. 68:848–860. [PubMed] [Google Scholar]15. Hino, S., and P. W. Wilson. 1957. Nitrogen fixation by a facultative Bacillus. J. Bacteriol. 57:403–408. [PMC free article] [PubMed] [Google Scholar]16. Horneck, G., H. Buckner, and G. Reitz. 1994. Long-term survival of bacterial spores in space. Adv. Space Res. 14:41–45. [PubMed] [Google Scholar]17. Huber, H., M. J. Hohn, R. Rachel, T. Fuchs, V. C. Wimmer, and K. O. Stetter. 2002. A new phylum of Archaea represented by a nanosized hyperthermophilic symbiont. Nature 417:63–67. [PubMed] [Google Scholar]18. Huber, R., T. A. Langworthy, H. König, M. Thomm, C. R. Woese, U. B. Sleytr, and K. O. Stetter. 1986. Thermotoga maritima sp. nov. represents a new genus of unique extremely thermophilic eubacteria growing up to 90°C. Arch. Microbiol. 144:324–333. [Google Scholar]19. Hungate, R. E. 1950. The anaerobic mesophilic cellulolytic bacteria. Bacteriol. Rev. 14:1–49. [PMC free article] [PubMed] [Google Scholar]20. Hungate, R. E. 1969. A roll tube method for cultivation of strict anaerobes, p. 117–132. In J. R. Norris and D. W. Ribbons (ed.), Methods in microbiology.

Academic Press, New York. NY.21. Kral, T. A., C. R. Bekkum, and C. P. McKay. 2004. Growth of methanogens on a Mars soil simulant. Orig. Life Evol. Biosph. 34:615–626. [PubMed] [Google Scholar]22. Küsel, K., C. Wagner, and H. L. Drake. 1999. Enumeration and metabolic product profiles of the anaerobic microflora in the mineral soil and litter of a beech forest. FEMS Microbiol. Ecol. 29:91–103. [Google Scholar]23. La Duc, M. T., W. Nicholson, R. Kern, and K. Venkateswaran. 2003. Microbial characterization of the Mars Odyssey spacecraft and its encapsulation facility. Environ. Microbiol. 67:977–985. [PMC free article] [PubMed] [Google Scholar]13. Fisk, M. R., and S. J. Giovanni. 1999. Sources of nutrients and energy for a deep biosphere on Mars. J. Geophys. Res. 104:11805–11816. [Google Scholar]14. Hillmann, F., R. J. Fischer, F. Saint-Prix, L. Girbal, and H. Bahl. 2008. PerR acts as a switch for oxygen tolerance in the strict anaerobe Clostridium acetobutylicum. Mol. Microbiol. 68:848–860. [PubMed] [Google Scholar]15. Hino, S., and P. W. Wilson. 1957. Nitrogen fixation by a facultative Bacillus. J. Bacteriol. 57:403–408. [PMC free article] [PubMed] [Google Scholar]16. Horneck, G., H. Buckner, and G. Reitz. 1994. Long-term survival of bacterial spores in space. Adv. Space Res. 14:41–45. [PubMed] [Google Scholar]17. Huber, H., M. J. Hohn, R. Rachel, T. Fuchs, V. C. Wimmer, and K. O. Stetter. 2002. A new phylum of Archaea represented by a nanosized hyperthermophilic symbiont. Nature 417:63–67. [PubMed] [Google Scholar]18. Huber, R., T. A. Langworthy, H. König, M. Thomm, C. R. Woese, U. B. Sleytr, and K. O. Stetter. 1986. Thermotoga maritima sp. nov. represents a new genus of unique extremely thermophilic eubacteria growing up to 90°C. Arch. Microbiol. 144:324–333. [Google Scholar]19. Hungate, R. E. 1950. The anaerobic mesophilic cellulolytic bacteria. Bacteriol. Rev. 14:1–49. [PMC free article] [PubMed] [Google Scholar]20. Hungate, R. E. 1969. A roll tube method for cultivation of strict anaerobes, p. 117–132. In J. R. Norris and D. W. Ribbons (ed.), Methods in microbiology.

Academic Press, New York. NY.21. Kral, T. A., C. R. Bekkum, and C. P. McKay. 2004. Growth of methanogens on a Mars soil simulant. Orig. Life Evol. Biosph. 34:615–626. [PubMed] [Google Scholar]22. Küsel, K., C. Wagner, and H. L. Drake. 1999. Enumeration and metabolic product profiles of the anaerobic microflora in the mineral soil and litter of a beech forest. FEMS Microbiol. Ecol. 29:91–103. [Google Scholar]23. La Duc, M. T., W. Nicholson, R. Kern, and K. Venkateswaran. 2003. Microbial characterization of the Mars Odyssey spacecraft and its encapsulation facility. Environ. Microbiol. 67:977–985. [PMC free article] [PubMed] [Google Scholar]13. Fisk, M. R., and S. J. Giovanni. 1999. Sources of nutrients and energy for a deep biosphere on Mars. J. Geophys. Res. 104:11805–11816. [Google Scholar]14. Hillmann, F., R. J. Fischer, F. Saint-Prix, L. Girbal, and H. Bahl. 2008. PerR acts as a switch for oxygen tolerance in the strict anaerobe Clostridium acetobutylicum. Mol. Microbiol. 68:848–860. [PubMed] [Google Scholar]15. Hino, S., and P. W. Wilson. 1957. Nitrogen fixation by a facultative Bacillus. J. Bacteriol. 57:403–408. [PMC free article] [PubMed] [Google Scholar]16. Horneck, G., H. Buckner, and G. Reitz. 1994. Long-term survival of bacterial spores in space. Adv. Space Res. 14:41–45. [PubMed] [Google Scholar]17. Huber, H., M. J. Hohn, R. Rachel, T. Fuchs, V. C. Wimmer, and K. O. Stetter. 2002. A new phylum of Archaea represented by a nanosized hyperthermophilic symbiont. Nature 417:63–67. [PubMed] [Google Scholar]18. Huber, R., T. A. Langworthy, H. König, M. Thomm, C. R. Woese, U. B. Sleytr, and K. O. Stetter. 1986. Thermotoga maritima sp. nov. represents a new genus of unique extremely thermophilic eubacteria growing up to 90°C. Arch. Microbiol. 144:324–333. [Google Scholar]19. Hungate, R. E. 1950. The anaerobic mesophilic cellulolytic bacteria. Bacteriol. Rev. 14:1–49. [PMC free article] [PubMed] [Google Scholar]20. Hungate, R. E. 1969. A roll tube method for cultivation of strict anaerobes, p. 117–132. In J. R. Norris and D. W. Ribbons (ed.), Methods in microbiology.

Academic Press, New York. NY.21. Kral, T. A., C. R. Bekkum, and C. P. McKay. 2004. Growth of methanogens on a Mars soil simulant. Orig. Life Evol. Biosph. 34:615–626. [PubMed] [Google Scholar]22. Küsel, K., C. Wagner, and H. L. Drake. 1999. Enumeration and metabolic product profiles of the anaerobic microflora in the mineral soil and litter of a beech forest. FEMS Microbiol. Ecol. 29:91–103. [Google Scholar]23. La Duc, M. T., W. Nicholson, R. Kern, and K. Venkateswaran. 2003. Microbial characterization of the Mars Odyssey spacecraft and its encapsulation facility. Environ. Microbiol. 67:977–985. [PMC free article] [PubMed] [Google Scholar]13. Fisk, M. R., and S. J. Giovanni. 1999. Sources of nutrients and energy for a deep biosphere on Mars. J. Geophys. Res. 104:11805–11816. [Google Scholar]14. Hillmann, F., R. J. Fischer, F. Saint-Prix, L. Girbal, and H. Bahl. 2008. PerR acts as a switch for oxygen tolerance in the strict anaerobe Clostridium acetobutylicum. Mol. Microbiol. 68:848–860. [PubMed] [Google Scholar]15. Hino, S., and P. W. Wilson. 1957. Nitrogen fixation by a facultative Bacillus. J. Bacteriol. 57:403–408. [PMC free article] [PubMed] [Google Scholar]16. Horneck, G., H. Buckner, and G. Reitz. 1994. Long-term survival of bacterial spores in space. Adv. Space Res. 14:41–45. [PubMed] [Google Scholar]17. Huber, H., M. J. Hohn, R. Rachel, T. Fuchs, V. C. Wimmer, and K. O. Stetter. 2002. A new phylum of Archaea represented by a nanosized hyperthermophilic symbiont. Nature 417:63–67. [PubMed] [Google Scholar]18. Huber, R., T. A. Langworthy, H. König, M. Thomm, C. R. Woese, U. B. Sleytr, and K. O. Stetter. 1986. Thermotoga maritima sp. nov. represents a new genus of unique extremely thermophilic eubacteria growing up to 90°C. Arch. Microbiol. 144:324–333. [Google Scholar]19. Hungate, R. E. 1950. The anaerobic mesophilic cellulolytic bacteria. Bacteriol. Rev. 14:1–49. [PMC free article] [PubMed] [Google Scholar]20. Hungate, R. E. 1969. A roll tube method for cultivation of strict anaerobes, p. 117–132. In J. R. Norris and D. W. Ribbons (ed.), Methods in microbiology.

Academic Press, New York. NY.21. Kral, T. A., C. R. Bekkum, and C. P. McKay. 2004. Growth of methanogens on a Mars soil simulant. Orig. Life Evol. Biosph. 34:615–626. [PubMed] [Google Scholar]22. Küsel, K., C. Wagner, and H. L. Drake. 1999. Enumeration and metabolic product profiles of the anaerobic microflora in the mineral soil and litter of a beech forest. FEMS Microbiol. Ecol. 29:91–103. [Google Scholar]23. La Duc, M. T., W. Nicholson, R. Kern, and K. Venkateswaran. 2003. Microbial characterization of the Mars Odyssey spacecraft and its encapsulation facility. Environ. Microbiol. 67:977–985. [PMC free article] [PubMed] [Google Scholar]13. Fisk, M. R., and S. J. Giovanni. 1999. Sources of nutrients and energy for a deep biosphere on Mars. J. Geophys. Res. 104:11805–11816. [Google Scholar]14. Hillmann, F., R. J. Fischer, F. Saint-Prix, L. Girbal, and H. Bahl. 2008. PerR acts as a switch for oxygen tolerance in the strict anaerobe Clostridium acetobutylicum. Mol. Microbiol. 68:848–860. [PubMed] [Google Scholar]15. Hino, S., and P. W. Wilson. 1957. Nitrogen fixation by a facultative Bacillus. J. Bacteriol. 57:403–408. [PMC free article] [PubMed] [Google Scholar]16. Horneck, G., H. Buckner, and G. Reitz. 1994. Long-term survival of bacterial spores in space. Adv. Space Res. 14:41–45. [PubMed] [Google Scholar]17. Huber, H., M. J. Hohn, R. Rachel, T. Fuchs, V. C. Wimmer, and K. O. Stetter. 2002. A new phylum of Archaea represented by a nanosized hyperthermophilic symbiont. Nature 417:63–67. [PubMed] [Google Scholar]18. Huber, R., T. A. Langworthy, H. König, M. Thomm, C. R. Woese, U. B. Sleytr, and K. O. Stetter. 1986. Thermotoga maritima sp. nov. represents a new genus of unique extremely thermophilic eubacteria growing up to 90°C. Arch. Microbiol. 144:324–333. [Google Scholar]19. Hungate, R. E. 1950. The anaerobic mesophilic cellulolytic bacteria. Bacteriol. Rev. 14:1–49. [PMC free article] [PubMed] [Google Scholar]20. Hungate, R. E. 1969. A roll tube method for cultivation of strict anaerobes, p. 117–132. In J. R. Norris and D. W. Ribbons (ed.), Methods in microbiology.

Academic Press, New York. NY.21. Kral, T. A., C. R. Bekkum, and C. P. McKay. 2004. Growth of methanogens on a Mars soil simulant. Orig. Life Evol. Biosph. 34:615–626. [PubMed] [Google Scholar]22. Küsel, K., C. Wagner, and H. L. Drake. 1999. Enumeration and metabolic product profiles of the anaerobic microflora in the mineral soil and litter of a beech forest. FEMS Microbiol. Ecol. 29:91–103. [Google Scholar]23. La Duc, M. T., W. Nicholson, R. Kern, and K. Venkateswaran. 2003. Microbial characterization of the Mars Odyssey spacecraft and its encapsulation facility. Environ. Microbiol. 67:977–985. [PMC free article] [PubMed] [Google Scholar]13. Fisk, M. R., and S. J. Giovanni. 1999. Sources of nutrients and energy for a deep biosphere on Mars. J. Geophys. Res. 104:11805–11816. [Google Scholar]14. Hillmann, F., R. J. Fischer, F. Saint-Prix, L. Girbal, and H. Bahl. 2008. PerR acts as a switch for oxygen tolerance in the strict anaerobe Clostridium acetobutylicum. Mol. Microbiol. 68:848–860. [PubMed] [Google Scholar]15. Hino, S., and P. W. Wilson. 1957. Nitrogen fixation by a facultative Bacillus. J. Bacteriol. 57:403–408. [PMC free article] [PubMed] [Google Scholar]16. Horneck, G., H. Buckner, and G. Reitz. 1994. Long-term survival of bacterial spores in space. Adv. Space Res. 14:41–45. [PubMed] [Google Scholar]17. Huber, H., M. J. Hohn, R. Rachel, T. Fuchs, V. C. Wimmer, and K. O. Stetter. 2002. A new phylum of Archaea represented by a nanosized hyperthermophilic symbiont. Nature 417:63–67. [PubMed] [Google Scholar]18. Huber, R., T. A. Langworthy, H. König, M. Thomm, C. R. Woese, U. B. Sleytr, and K. O. Stetter. 1986. Thermotoga maritima sp. nov. represents a new genus of unique extremely thermophilic eubacteria growing up to 90°C. Arch. Microbiol. 144:324–333. [Google Scholar]19. Hungate, R. E. 1950. The anaerobic mesophilic cellulolytic bacteria. Bacteriol. Rev. 14:1–49. [PMC free article] [PubMed] [Google Scholar]20. Hungate, R. E. 1969. A roll tube method for cultivation of strict anaerobes, p. 117–132. In J. R. Norris and D. W. Ribbons (ed.), Methods in microbiology.

Academic Press, New York. NY.21. Kral, T. A., C. R. Bekkum, and C. P. McKay. 2004. Growth of methanogens on a Mars soil simulant. Orig. Life Evol. Biosph. 34:615–626. [PubMed] [Google Scholar]22. Küsel, K., C. Wagner, and H. L. Drake. 1999. Enumeration and metabolic product profiles of the anaerobic microflora in the mineral soil and litter of a beech forest. FEMS Microbiol. Ecol. 29:91–103. [Google Scholar]23. La Duc, M. T., W. Nicholson, R. Kern, and K. Venkateswaran. 2003. Microbial characterization of the Mars Odyssey spacecraft and its encapsulation facility. Environ. Microbiol. 67:977–985. [PMC free article] [PubMed] [Google Scholar]13. Fisk, M. R., and S. J. Giovanni. 1999. Sources of nutrients and energy for a deep biosphere on Mars. J. Geophys. Res. 104:11805–11816. [Google Scholar]14. Hillmann, F., R. J. Fischer, F. Saint-Prix, L. Girbal, and H. Bahl. 2008. PerR acts as a switch for oxygen tolerance in the strict anaerobe Clostridium acetobutylicum. Mol. Microbiol. 68:848–860. [PubMed] [Google Scholar]15. Hino, S., and P. W. Wilson. 1957. Nitrogen fixation by a facultative Bacillus. J. Bacteriol. 57:403–408. [PMC free article] [PubMed] [Google Scholar]16. Horneck, G., H. Buckner, and G. Reitz. 1994. Long-term survival of bacterial spores in space. Adv. Space Res. 14:41–45. [PubMed] [Google Scholar]17. Huber, H., M. J. Hohn, R. Rachel, T. Fuchs, V. C. Wimmer, and K. O. Stetter. 2002. A new phylum of Archaea represented by a nanosized hyperthermophilic symbiont. Nature 417:63–67. [PubMed] [Google Scholar]18. Huber, R., T. A. Langworthy, H. König, M. Thomm, C. R. Woese, U. B. Sleytr, and K. O. Stetter. 1986. Thermotoga maritima sp. nov. represents a new genus of unique extremely thermophilic eubacteria growing up to 90°C. Arch. Microbiol. 144:324–333. [Google Scholar]19. Hungate, R. E. 1950. The anaerobic mesophilic cellulolytic bacteria. Bacteriol. Rev. 14:1–49. [PMC free article] [PubMed] [Google Scholar]20. Hungate, R. E. 1969. A roll tube method for cultivation of strict anaerobes, p. 117–132. In J. R. Norris and D. W. Ribbons (ed.), Methods in microbiology.

Academic Press, New York. NY.21. Kral, T. A., C. R. Bekkum, and C. P. McKay. 2004. Growth of methanogens on a Mars soil simulant. Orig. Life Evol. Biosph. 34:615–626. [PubMed] [Google Scholar]22. Küsel, K., C. Wagner, and H. L. Drake. 1999. Enumeration and metabolic product profiles of the anaerobic microflora in the mineral soil and litter of a beech forest. FEMS Microbiol. Ecol. 29:91–103. [Google Scholar]23. La Duc, M. T., W. Nicholson, R. Kern, and K. Venkateswaran. 2003. Microbial characterization of the Mars Odyssey spacecraft and its encapsulation facility. Environ. Microbiol. 67:977–985. [PMC free article] [PubMed] [Google Scholar]13. Fisk, M. R., and S. J. Giovanni. 1999. Sources of nutrients and energy for a deep biosphere on Mars. J. Geophys. Res. 104:11805–11816. [Google Scholar]14. Hillmann, F., R. J. Fischer, F. Saint-Prix, L. Girbal, and H. Bahl. 2008. PerR acts as a switch for oxygen tolerance in the strict anaerobe Clostridium acetobutylicum. Mol. Microbiol. 68:848–860. [PubMed] [Google Scholar]15. Hino, S., and P. W. Wilson. 1957. Nitrogen fixation by a facultative Bacillus. J. Bacteriol. 57:403–408. [PMC free article] [PubMed] [Google Scholar]16. Horneck, G., H. Buckner, and G. Reitz. 1994. Long-term survival of bacterial spores in space. Adv. Space Res. 14:41–45. [PubMed] [Google Scholar]17. Huber, H., M. J. Hohn, R. Rachel, T. Fuchs, V. C. Wimmer, and K. O. Stetter. 2002. A new phylum of Archaea represented by a nanosized hyperthermophilic symbiont. Nature 417:63–67. [PubMed] [Google Scholar]18. Huber, R., T. A. Langworthy, H. König, M. Thomm, C. R. Woese, U. B. Sleytr, and K. O. Stetter. 1986. Thermotoga maritima sp. nov. represents a new genus of unique extremely thermophilic eubacteria growing up to 90°C. Arch. Microbiol. 144:324–333. [Google Scholar]19. Hungate, R. E. 1950. The anaerobic mesophilic cellulolytic bacteria. Bacteriol. Rev. 14:1–49. [PMC free article] [PubMed] [Google Scholar]20. Hungate, R. E. 1969. A roll tube method for cultivation of strict anaerobes, p. 117–132. In J. R. Norris and D. W. Ribbons (ed.), Methods in microbiology.