

Before & After Instructions Form for Filler Before Filler Treatment Instructions

Please inform us if you are pregnant, breastfeeding or under 18 years of age as products used for the following treatments may not be appropriate

- Avoid alcoholic beverages and green tea.
- Avoid any substances that affect platelet function, such as aspirin, non steroidal anti-inflammatory drugs (e.g. Advil, Motrin), and supplements of fish oil, vitamin E and St John's Wort. These substances may increase the risk of bruising and bleeding at the injection site.
- Avoid waxing, bleaching tweezing, or hair removal creams in the area of treatment.
- Avoid makeup the day of treatment.
- Talk with your doctor if you have a history of herpes viral infection (e.g cold sores) as injection treatments can lead to the reactivation of herpes viral infections.

After Filler Treatment Instructions

- Avoid lying down for 4 hours following treatment.
- Avoid manipulation of the area the first 4 hours after the procedure.
- No Alcohol or ASA for 24 hours after.
- No makeup for 24 hours on treated area.
- No chemical solutions or facial cleansers to face for 24 hours.
- May apply ice for swelling if required.
- No taking a trip via plane for 2 weeks.
- No vaccination for 3 weeks.
- No dental work for 2 weeks.
- No sipping through a straw for 24 hours.
- Drink lots of water!

Additional Information

- A touch-up may be necessary in 2 weeks.
- Contact the practitioner as soon as possible if you are experiencing extreme pain around the injection site along with discolouration.