

Topics: [Logistic Differential Equations](#)

Goals: Students will be able to model problems with differential equations and use current data to write a logistic model for the data.

Daily Recommended Schedule:

[Monday:](#) Read summary and do Problem 1

[Tuesday:](#) Do Problems 2 and 3 and read Problem 4.

[Wednesday:](#) Do Problem 4a

[Thursday:](#) Do Problem 4b.

[Friday:](#) Do Differential Equation Review.

Optional:

If you have access to the internet and would like to help me test this out, please, click on the link below corresponding to each day and try the check for understandings.

[Monday](#)

[Tuesday](#)

[Wednesday](#)

Resources:

[Differential Equations Cheat Sheet](#) (end of packet)

[Review of Partial Fractions](http://www.baileyworldofmath.org) (www.baileyworldofmath.org)

[Review Lesson on Euler's Method](http://www.baileyworldofmath.org) (www.baileyworldofmath.org)

[Khan Academy Video on Separable Differential Equations](#)

[Khan Academy Videos on Logistic Differential Equations](#)

MONDAY: Although a population grows exponentially at first, it cannot outgrow its resources without consequences. For this reason, Verhust, a Belgian mathematician modeled population growth with a more accurate differential equation given by the theory that the rate of population growth is proportional to the population and its difference to the limiting population.

This is modeled by the differential equation:

$$\frac{dP}{dt} = kP(L - P)$$

Where the population function is given by $P(t)$ at time t with limiting population, L .

Example 1:

A Petri dish has a carrying capacity of 24 micrograms of bacteria. If a scientist initially introduced 6 micrograms of bacteria and the proportionality factor is 3 micrograms per hour.

Write the differential equation: (Use the limiting population (carrying capacity) and proportionality factor:

Separate the Variables and Integrate both Sides (You will need partial fractions):

Use the initial condition:

Solve for $P(t)$:

TUESDAY: Make sure to show your work

1.

A population of wolves is modeled by the function P and grows according to the logistic differential equation $\frac{dP}{dt} = 5P\left(1 - \frac{P}{5000}\right)$, where t is the time in years and $P(0) = 1000$. Which of the following statements are true?

- I. $\lim_{t \rightarrow \infty} P(t) = 5000$
- II. $\frac{dP}{dt}$ is positive for $t > 0$.
- III. $\frac{d^2P}{dt^2}$ is positive for $t > 0$.

- (A) I only
- (B) II only
- (C) I and II only
- (D) I and III only
- (E) I, II, and III

2. A population of javelinas is modeled by the function P and grows according to the logistic

differential equation $\frac{dP}{dt} = P\left(2 - \frac{P}{3000}\right)$, where t is the time in years and $P(0) = 1000$. Which of the following is equivalent to $\lim_{t \rightarrow \infty} P(t)$?


- A) 1500
- B) 3000
- C) 4500
- D) 6000

Wednesday:

4. The Coronavirus Covid 19 is infecting people exponentially right now, but the model is actually a logistic equation because the limiting population is the population in the area in which the virus resides.

a)

CHINA




<https://www.worldometers.info/coronavirus/country/china/>

The graph above is a link to the actual data for the cumulative Covid19 cases in China. You can hover your mouse over the graph and find the actual numbers. Use some data points to find a logarithmic function to model the data above.

Thursday:

b)

UNITED STATES



<https://www.worldometers.info/coronavirus/country/us/>

The graph above is a link to Covid19 data for the US. Write a logarithmic model for this data. You may have to guess the limiting capacity. How close is your model to the actual data? Explain.

Friday: Differential Equation Reviews

1) Consider the differential equation $\frac{dy}{dx} = y^2(2x+2)$. Let $y = f(x)$ be the particular solution to the differential equation with initial condition $f(0) = -1$.

(a) Use Euler's Method, starting at $x = 0$ with two steps of equal size, to approximate $f\left(\frac{1}{2}\right)$.

b) Find $y = f(x)$, the particular solution to the differential equation with initial condition $f(0) = -1$.


3) A certain rumor spreads through a community at the rate of $\frac{dy}{dt} = 2y(1-y)$, where y is the proportion of the population that has heard the rumor at time t . This is a logistic differential equation.

a) If at time $t = 0$ ten percent of the people have heard the rumor, find y as a function of t .

c) Find $\lim_{t \rightarrow \infty} y(t)$, and explain its meaning in the context of the problem.

Logistic Differential Equations

$$\frac{dy}{dx} = ay \left(1 - \frac{y}{L}\right) \quad \text{or} \quad \frac{dy}{dx} = ay \left(\frac{L-y}{L}\right)$$

<i>Partial Fraction Decomposition:</i>	<i>Information about y</i>
$\frac{dy}{dx} = ay \left(1 - \frac{y}{L}\right)$ $\frac{dy}{dx} = ay \left(\frac{L-y}{L}\right)$ $\frac{Ldy}{y(L-y)} = adx$ $\int \frac{1}{y} + \frac{1}{L-y} = \int adx$ $\ln(y) - \ln(L-y) = ax + C$ $\ln\left(\frac{y}{L-y}\right) = ax + C$ $\frac{y}{L-y} = e^{ax+C}$ $\frac{L-y}{y} = Pe^{-ax}$ $1 - \frac{L}{y} = -Pe^{-ax}$ $\frac{L}{y} = 1 + Pe^{-ax}$ $y = \frac{L}{1 + Pe^{-ax}}$	<p>Models Population Growth</p> $y = \frac{L}{1 - Pe^{-ax}}$ <p>L is called the limiting factor or limiting population or carrying capacity</p> $\lim_{x \rightarrow \infty} y(x) = L$  $C = \frac{L - y(0)}{y(0)}$ <p>$y(x)$ is growing x_1 fastest at where $y(x_1) = \frac{L}{2}$</p>

Differential Equations and Euler's Method

- Use Implicit Differentiation to find $\frac{d^2y}{dx^2}$. Substitute what you have as $\frac{dy}{dx}$.
- Euler's Method uses a Δx and some steps.
 - Use that $y_2 = y_1 + \frac{dy}{dx} \Big|_{(x_1, y_1)} \Delta x$.
 - Then use $y_3 = y_2 + \frac{dy}{dx} \Big|_{(x_2, y_2)} \Delta x$
 - Etc

(continued on next page)

- To solve Separable Differential Equations
 - Separate the variables
 - y 's and dy 's on one side of the equal sign
 - x 's and dx 's on the other side of the equal sign
 - dy 's and dx 's are NOT allowed to be in the denominator when integrating
 - Integrate both sides
 - Don't forget the C
 - C should only be on the side with the x .
 - Use Initial condition
 - This can be done before solving for y to find " C "
 - Solve for y
 - Sometimes y is called P and x is called t : don't let this confuse you