

POST-OPERATIVE INSTRUCTIONS PROCEDURE: DISTAL BICEPS TENDON REPAIR, ELBOW

- Pain Medication: You will be given a prescription for pain medication at your preoperative appointment. This should be filled and ready for use when you return home from the hospital/surgery center. You should not drive, operate heavy machinery or participate in activities that require concentration while taking narcotic pain medications such as Percocet, Oxycodone, Oxycontin, Norco, Vicodin or Tylenol with Codeine. You should take an over the counter stool softener while taking narcotic pain medications to prevent constipation.
- <u>Dressings</u>: Leave the dressing/splint intact until you return for your post-operative visit. Keep incision dry and clean until sutures are removed at your first post-op visit. Do not get the incisions or splint wet until you have bee seen by your surgeon and given approval for showering. You will not be able to soak the incision in a bath, pool, or body of water for 4 weeks.
- ➤ <u>Sling/Brace</u>: Use at all times (even when sleeping).
- ➤ <u>Post-Operative Appointment</u>: Please make a post-operative appointment to see your surgeon 10-14 days following surgery.
- ➤ <u>Prescription Refill Protocol</u>: You <u>MUST</u> give your physician <u>48 hours</u> notice for any medication refill. Many medications require a written prescription and cannot be called into a pharmacy. Prescriptions will not be filled over the weekend.
- When to call your surgeon:
 - o Complications after surgery are fortunately very rare. Call the office if you have any of the following symptoms:
 - Persistent fever >101, chills
 - Increasing pain at the surgical site
 - New onset numbness or tingling
 - Hives or new rashes
 - Shortness of breath of chest pain
 - Persistent nausea/vomiting
 - Drainage from your incision lasting >5 days