TAKE 5

CHECKING YOUR BLOOD GLUCOSE

Blood glucose tracking is an essential part of taking care of your diabetes.



How frequent blood glucose (blood sugar) checks can help you

Checking your blood glucose is an important part of managing diabetes. Checking often will tell you:

- How well your insulin and other diabetes medicines are working
- How physical activity and the foods you eat affect your blood glucose

You'll usually feel better and have more energy when your blood glucose stays at or near normal. Managing your blood glucose can also reduce your risk of developing problems from diabetes.



How to check your blood glucose

You can check your own blood glucose using a meter. There are two main types:

- Standard blood glucose meter checks your blood glucose levels at that moment using a drop of blood
- Continuous glucose monitor (CGM) checks your blood glucose regularly day or night through a sensor placed on your body

Your diabetes care team can help you choose one and show you how to use it.

When to check your blood glucose

You and your diabetes care team will decide when and how often to check your blood glucose. The table below shows some times when you might want to check and why.

When your team may want you to check	Why you should check
When you wake up	To see if your blood glucose is maintained at goal while you're asleep (called fasting blood glucose)
Before meals	To know what your blood glucose is before you eat and to know if you will need to adjust your mealtime (prandial) insulin
1 or 2 hours after you start your meal	To see how your blood glucose is affected by insulin and/or the food you eat
Before, during, and after physical activity	To see how being active affects your blood glucose
At bedtime	Depending on the medicine that you take

Keeping a blood glucose tracker

It's important to keep track of your blood glucose levels and what makes them go up or down. See the last page for a tracker that you can use to record your levels and other important information.

To track your readings in an app, download the FREE Cornerstones4Care® app to your mobile device.



CHECKING YOUR BLOOD GLUCOSE

Time	Goals for many nonpregnant adults with diabetes	Your goals
Before meals	80 to 130 mg/dL	
1 or 2 hours after the start of a meal	Less than 180 mg/dL	
A1C	Less than 7%	

Adapted from the American Diabetes Association. Standards of medical care in diabetes—2021. Diabetes Care. 2021 Jan;44(suppl1):S1-S232.

Setting your blood glucose goals

The table above lists blood glucose goals for many nonpregnant adults with diabetes. You and your diabetes care team will set the goals that are right for you.

Knowing your A1C

The A1C test measures your estimated average blood glucose levels over approximately 3 months. It's like a "memory" of your blood glucose levels over time. The table to the right shows how A1C relates to the estimated average blood glucose level.

Lowering your A1C to below 7% reduces your risk of problems from diabetes. Therefore, the A1C goal for most people is less than 7%.

It is recommended that you get an A1C test:

- At least 2 times a year if your blood glucose levels are at your goal
- 4 times a year if you are not meeting your goals or if your treatment has changed

Average blood glucose (blood sugar)				
97 mg/dL				
126 mg/dL				
154 mg/dL				
183 mg/dL				
212 mg/dL				
240 mg/dL				
269 mg/dL				
298 mg/dL				

Adapted from the American Diabetes Association. Standards of medical care in diabetes—2021. *Diabetes Care*. 2021 Jan;44(suppl1):S1-S232.



If you have any questions, be sure to talk with your diabetes care team. They are there to help!

Visit **Cornerstones4Care.com** for additional resources and to register for a FREE diabetes support program! Point your smartphone camera at the QR code for quick access to the website on your phone.













Novo Nordisk Inc. grants permission to reproduce this piece for nonprofit educational purposes only on condition that the piece is maintained in its original format and that the copyright notice is displayed. Novo Nordisk Inc. reserves the right to revoke this permission at any time.

Cornerstones4Care® is a registered trademark of Novo Nordisk A/S.

Novo Nordisk is a registered trademark of Novo Nordisk A/S.

© 2021 Novo Nordisk Printed in the U.S.A. US21DI00084 August 2021





HOW TO USE YOUR BLOOD GLUCOSE TRACKER

Your blood glucose tracker

Diabetes changes over time. That is why your treatment may also need to change over time.


The blood glucose tracker on the next page can help you keep track of your diabetes medicines and any changes in dosage and timing that your doctor may tell you to make. It can also help you keep track of how much mealtime insulin you took and when you took it.

Your diabetes care team can provide you with a blood glucose tracker and show you how to use it. Or call **1-800-727-6500** to have a tracker sent to you.

Here's how to use the blood glucose tracker

- Write down the date for the start of the week. (You can start tracking on any day of the week)
- Write the name(s) and dose(s) of your non-insulin diabetes medicine(s)
- Write the time and your blood glucose readings in the "before" and "after" spaces. After-meal readings are usually taken 1 to 2 hours after you start your meal. Nighttime readings may be taken during the night as needed
- If you are counting carbs, write how many grams of carbs you ate
- If your doctor has told you to use mealtime insulin when you eat, write your dose here
- If your doctor has told you to use long-acting insulin, write your dose and time(s) here. Longacting insulin is taken either once or twice a day
- Add notes on anything else you might want to track (such as blood pressure, activity, or weight)

EXAMPLE



YOUR BLOOD GLUCOSE TRACKER

						Date:	/		
		BLOOD GLUCOSE RESULTS*							
		kfast	Lui	nch	Din	ner	Bedtime	Night	
MONDAY	BEFORE			AFTER			Time	Time	
.⊑ ▶	Time	Time	Time	Time	Time	Time			
ine osce	ma/dl	mg/dL	ma/dl	ma/dl	ma/dl	ma/dl	mg/dL	ma/dl	
Non-insulin dose Carb intake Mealtime insulin dose		IIIg/uL	IIIg/uL	IIIg/uL	IIIg/uL	IIIg/uL	IIIg/uL	mg/dL	
Carb intake Mealtime insulin dose	☐ Breakfast _	units	Lunch	units	☐ Dinner	units	Other >		
Long-acting insulin dose			If needed at di	nner or bedtime					
		kfast		nch		ner	Bedtime	Night	
TUESDAY	BEFORE	AFTER	BEFORE		BEFORE	AFTER	Time	Time	
	Time	Time	Time	Time	Time	Time			
Non-insulin dose Carb intake Mealtime insulin dose Mealtime in	(-II			(-II			, , , , , , , , , , , , , , , , , , ,		
Carb intake ►	mg/aL	mg/dL	mg/aL	mg/dL	mg/aL	mg/dL	mg/dL	mg/dL	
Mealtime insulin dose	☐ Breakfast	units	□Lunch	units	□ Dinner	units	Other >		
Long-acting insulin dose		time					Other		
3 3	Brea	kfast	Lui	nch	Din	ner	Bedtime	Night	
WEDNESDAY	BEFORE	AFTER	BEFORE	AFTER	BEFORE	AFTER	Time	Time	
	Time	Time	Time	Time	Time	Time			
Monting in the property of the									
Non-insular description descri	mg/dL	mg/dL	mg/dL	mg/dL	mg/dL	mg/dL	mg/dL	mg/dL	
Carb intake Mealtime insulin dose		unite	□ Lunch	unite	□ Dinner	unite	Other >		
Long-acting insulin dose	units	time	If needed at di	nner or bedtime	units	time	Other		
		kfast	Lui		Din		Bedtime	Niaht	
THURSDAY	BEFORE		BEFORE	AFTER		AFTER	Time	Time	
	Time	Time	Time	Time	Time	Time			
suliins se									
Montaine insuling dose		mg/dL	mg/dL	mg/dL	mg/dL	mg/dL	mg/dL	mg/dL	
Carb intake Mealtime insulin dose		unite	□ Lunch	units	□ Dinner	unite	Other >		
Long-acting insulin dose		time		nner or bedtime			Other		
3		kfast		Lunch		Dinner		Night	
FRIDAY	BEFORE	AFTER	BEFORE		BEFORE	AFTER	Bedtime Time	Time	
	Time	Time	Time	Time	Time	Time			
Non-insuling dose Mealtime insulin dose Mealtime insulin dose									
dici:		mg/dL	mg/dL	mg/dL	mg/dL	mg/dL	mg/dL	mg/dL	
Carb intake Mealtime insulin dose		units	Lunch	units	☐ Dinner	units	Other >		
Long-acting insulin dose		time		nner or bedtime	units	time	Other		
		kfast		nch	Din	ner	Bedtime	Niaht	
SATURDAY	BEFORE	AFTER	BEFORE	AFTER	BEFORE	AFTER	Time	Time	
C •	Time	Time	Time	Time	Time	Time			
Non-insulin dose Mealtime insulin dose									
	mg/dL	mg/dL	mg/dL	mg/dL	mg/dL	mg/dL	mg/dL	mg/dL	
Carb intake Mealtime insulin dose	Rreakfast	units	Lunch	units	☐ Dinner	units	Other >		
Long-acting insulin dose		time		nner or bedtime	units	time	Other		
J 1 11 J 12 11 11 11 11 11 11 11 11 11 11 11 11		kfast		nch		ner	Bedtime	Night	
SUNDAY	BEFORE	AFTER	BEFORE	AFTER	BEFORE	AFTER	Time	Time	
	Time	Time	Time	Time	Time	Time			
suli									
Modring institution of the Marketine institut	mg/dL	mg/dL	mg/dL	mg/dL	mg/dL	mg/dL	mg/dL	mg/dL	
Non-insurer of documents of the control of the con		units	Lunch	units	☐ Dinner	units	Other >		
Long-acting insulin dose		time		nner or bedtime	units	time	Ould P		

