

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

PEST CONTROLLERS OWN

Rodent Soft Bait[®]

ACTIVE CONSTITUENT: 0.05 g/kg DIFENACOUM

READY TO USE PALATABLE BAIT FOR THE CONTROL OF NORWAY RATS AND MICE, including those resistant to other anti-coagulants.

FOR USE IN AND AROUND BUILDINGS, including industrial, commercial, agricultural, public services and domestic structures.

BEFORE USE READ ALL
DIRECTIONS ON THE
OUTER PACK



PCO
PEST CONTROLLERS OWN[®]

Manufactured by:

Amgrow Pty Ltd ABN 81 100 684 786
Unit B2A, 3-29 Birnie Avenue, Lidcombe NSW 2141
TELEPHONE: 1800 631 008

Amgrow

NET CONTENTS
2kg (CONTAINS 200 X 10G
BAIT SACHETS WHICH IT
IS ILLEGAL SEPARATELY)

DIRECTIONS FOR USE

Restrains: DO NOT place baits in the open unless in bait stations.
DO NOT place baits in locations accessible to children or pets.
DO NOT remove bait from paper wrapping; rats and mice will chew through the wrapping to consume the bait.

Pests	Situations	Rate	Critical Comments
Mice	For use in and around (within 2 metres), industrial, commercial, public services, domestic and agricultural buildings (machinery sheds etc)	Use 1 to 2 baits per bait point	Explore the site and locate runs, nests, droppings and signs of damage. Eliminate as far as practical, all alternate food sources. Place bait points wherever droppings, damage or other signs of activity are seen. As mice are sporadic feeders use many bait points, not more than 3 m apart.
Norway Rat (<i>Rattus norvegicus</i>)		Use up to 10 baits per bait point	Explore the site and locate runs, burrows, droppings and signs of damage. Eliminate as far as practical, all alternative food sources. Place a number of bait points in dry locations throughout the infested area, on runs, in burrows and where droppings are seen, but not more than 9 m apart. Use weather resistant bait stations when using bait outdoors.

General Information and Baiting Strategy for Rats and Mice: Eliminate as far as practicable all other food sources. Always place bait where it is inaccessible to children, animals and birds. Protect and cover bait. When using outdoors, use weather resistant bait stations. Alternatively place bait in suitable containers or use weather resistant bait stations constructed from drainpipes or boxes. Try not to touch the bait as this may deter mice and rats from eating it. DO NOT place bait where food, feed or water could become contaminated. Place baits only in situations where there is no ready access to occupants of the buildings. Record the number and locations of bait stations for the purpose of replacement and later removal. Inspect baits frequently, particularly in the first 10 days. Replace eaten bait. If bait is completely consumed, the quantity should be increased. Continue replenishing baits until they are left untouched. Baiting for at least two weeks will be necessary to reduce rat/mouse numbers. Two or more feeds are required to produce a delayed mortality. Most deaths occur 4 to 10 days after consumption of bait, but deaths may occur for up to 2 weeks. Following treatment remove all remains of bait and bait containers and look for and collect dead rats or mice. Burn or bury carcasses. DO NOT place in refuse bins. DO NOT let cats, dogs and non-target animals eat poisoned mice and rats. When handling remaining bait material/animal carcasses, wear rubber gloves.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

33166

GENERAL INSTRUCTIONS: Pest Controllers Own Rodent Soft Bait is a highly palatable cereal based all-purpose rodenticide based on the advanced anticoagulant difenacoum, which is equally as effective against rats and mice, including some strains resistant to other anticoagulants. In some circumstances, more than one feed may be required to produce mortality that generally occurs 4-10 days after consuming the bait. This product contains Bitrex (a human taste deterrent) to help prevent accidental human consumption.

PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS AND THE ENVIRONMENT:

This bait is hazardous to pigs, cats, dogs, poultry, birds and wildlife and care should be exercised in its use. DO NOT place baits in locations that are accessible to domestic animals, livestock, birds or other animals. DO NOT contaminate dams, waterways and drains with bait or its used containers. DO NOT use this product to control native animals. The product is restricted to use in and around buildings (within 2 metres), structures or within enclosed spaces e.g. drains. When using outdoors use weather resistant bait stations. In areas where possums and other non-target animals frequent, baits should be secured inside proper bait stations or bait stations constructed from drainpipes or boxes.

STORAGE AND DISPOSAL: Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Break, crush or puncture and deliver empty packaging (or unused baits) for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging or unused bait 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers and product. DO NOT use containers that have held bait for any other purpose. Inspect bait stations regularly and ensure baits are removed when rodent activity has ceased. Dispose of carcasses safely by burning or burying them.

SAFETY DIRECTIONS: Poisonous if swallowed. May irritate the eyes. Repeated minor exposure may have a cumulative poisoning effect. Avoid contact with eyes. Vitamin K1 (phytomenadione) is antidotal. Wash hands after use.

FIRST AID: If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. Vitamin K₁ (phytomenadione) is antidotal.

SAFETY DATA SHEET: Additional information is listed in the Safety Data Sheet which is available from the supplier.

NOTE TO PHYSICIANS AND VETERINARIANS: Difenacoum, the active ingredient in this product, is a long lasting coagulant chemical, which is ingested by humans, domestic animals, or pets, can reduce the clotting of the blood and haemorrhage may result. Vitamin K₁ (phytomenadione) should be administered urgently by medical, appropriate paramedical personnel or veterinarians. If administered intravenously the injection must be given slowly.

CONDITIONS OF SALE: The use of Pest Controllers Own Rodent Soft Bait being beyond the control of the manufacturer no warranty expressed or implied is given by Amgrow Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Amgrow Pty Ltd accepts no responsibility for any consequence whatsoever resulting from the use of this product.



Additional information required under the Globally Harmonised System (GHS) classification of the substance/mixture: Obtain special instructions before use. Do not handle until all safety precautions have been read and understood. Do not breathe fumes, mists, vapours or spray. Do not get in eyes, on skin, or on clothing. Wash contacted areas thoroughly after handling. Use only outdoors or in a well ventilated area. Contaminated work clothing should not be allowed out of the workplace. Avoid release to the environment. Wear protective gloves, protective clothing and eye or face protection. Wear respiratory protection.

APVMA Approval No: 85513/115837

PCO and Pest Controllers Own are registered trademarks

PCO
PEST CONTROLLERS OWN®

**IN A TRANSPORT
EMERGENCY DIAL 000
• POLICE OR FIRE
BRIGADE •**

Batch
No:

D.O.M.: