

## 2021/22 Vaccination Record

### All Bulls must have two rounds of vaccination prior to weaning

Please check and date vaccination use as applicable and/or indicate other vaccinations used

	<u>Pre-wean</u>	<u>Wean</u>	<u>Other</u>
<input type="checkbox"/> Bovishield Gold 5 or equivalent + Manneheimia Haemolytica:	_____ / _____	_____ / _____	_____ / _____
<input type="checkbox"/> Ultrabac 8 or equivalent	_____ / _____	_____ / _____	_____ / _____
<input type="checkbox"/> De-wormed: injectable required Brand: _____	_____ / _____	_____ / _____	_____ / _____
<input type="checkbox"/> Recommend injectable mineral be given. Multimin or a similar product requires a vet prescription	_____ / _____	_____ / _____	_____ / _____
Other vaccinations given (i.e.) Wart Vaccination			
— _____	_____ / _____	_____ / _____	_____ / _____
— _____	_____ / _____	_____ / _____	_____ / _____
— _____	_____ / _____	_____ / _____	_____ / _____
— _____	_____ / _____	_____ / _____	_____ / _____

Bulls must receive booster shots approximately four weeks prior to arrival at the test facilities. Bulls will be given another set of booster shots and dewormed at the test facility. Tassels on the sheath should be trimmed prior to delivery to lesson infections.

Bulls must be weaned at least 45 days prior to arrival at the test facility and it is recommended bulls are weaned for at least 60 days prior to arrival at the test facility.

I certify information provided above is accurate to the best of my knowledge:

Signed: \_\_\_\_\_ Date: \_\_\_\_\_