

Bioacoustic monitoring uncovers the breeding phenology of a critically endangered upland rainforest frog (Taudactylus pleione).

Natalie J. McLeod BSc(Hons)





Acknowledgement of Country

I would like to acknowledge the Gurang and Gooreng Gooreng people as the Traditional Custodians of the Country on which my research took place, the Turrbal/Jagera peoples where my thesis was written, and the Bundjalung people upon whose Country we meet today.


I recognise their continued connection to the land and waters and thank them for caring for its ecosystems for thousands of generations. I pay my respect to the Elders, past and present, and extend that respect to any First Nations people gathered here today.



Introduction

Amphibian extinction crisis

- Amphibians are the most endangered vertebrate class globally¹
- Disproportionate declines in undisturbed upland rainforests of the Americas and Australia^{2, 3}
- Patterns attributed to amphibian chytrid fungus (Bd), shared life-history traits and impacts of climate change




(Luedtke J. A., et al. 2023)


Species on the brink: Taudactylus pleione

- All six species in genus Taudactylus are highly threatened (two presumed extinct) by Bd and climate change⁴
- Taudactylus pleione most recently described member⁵
- Critically endangered under NCA 1992 and EPBC Act 1999
 - Extremely restricted and fragmented distribution, < 70 ha total
 - Low number of animals left in the wild
- Occurs only along headwater streams and seepages of Kroombit Tops NP
- Small and cryptic frog thar shelters under leaf litter and rocks





Calling for conservation: Taudactylus pleione



Male advertisement calls

- Circadian and circannual rhythms
- Environmental cues

Convey fitness

- Female partners
- Male competitors

Ecological information

- Breeding activity
- Population status

Species management

- Visual and aural surveys
- Bioacoustics




Aims and research questions

My research aimed to improve our understanding of the breeding phenology and detectability of a critically endangered upland rainforest frog, *Taudactylus pleione*, to contribute to their conservation.

My research questions:

- 1. Can an automated call recogniser be applied with a high-level of confidence?
- 2. What are the diel and seasonal patterns of calling activity?
- 3. How do abiotic environmental factors, such as temperature, humidity and rainfall, influence these patterns?

Methods






Results & Discussion

Recogniser performance





	Performance metric	Value (0-1)
Threshold 1 (≥ 1 matches/per min)	Accuracy (TP + TN) / (TP +TN + FP + FN)	0.85
	Precision (TP / TP + FP)	0.55
	Recall (TP / TP + FN)	0.60
Threshold 2 (≥ 21 matches/per min)	Accuracy (TP + TN) / (TP +TN + FP + FN)	0.89
	Precision (TP / TP + FP)	0.84
	Recall (TP / TP + FN)	0.43




General calling patterns










Diel calling activity











Seasonal calling activity



MaxT	MinT	. Hum	n BarP	Rain	Rain7	Rain30	Moon	Photo	MaxT×P	AICc	ΔΑΙСα	W
√	√	<i>✓</i>	·	√	· ✓	✓	Х	✓	✓	16,582.7	0.00	0.60

Fixed effects	Estimate	SE
Intercept	-0.86	0.42
Maximum temperature	0.08	0.04
Minimum temperature	0.17	0.04
Relative humidity	0.20	0.02
Barometric pressure	-0.10	0.03
Rainfall	-0.05	0.02
Rainfall 7-day	-0.16	0.03
Rainfall 30-day	0.11	0.02
Photoperiod	0.37	0.18
Max temperature × photoperiod	-0.21	0.04
Random effects	σ^2	
Site	0.65	
Frog season	0.16	
AR(1)	0.24	
Marginal / Conditional R ²	0.16 / 0.33	



^{*} Photoperiod levels are -1 standard deviation, +1 standard deviation and 0 equals the mean

Key findings & Future research

Summary

01

Can the call recogniser be applied with confidence?

The recogniser had high accuracy and precision, but a low recall rate may have influenced results

02

What are the diel and seasonal patterns of calling activity?

Calling was most likely in the summer from late-November to early-March at dawn and dusk

03

How do environmental factors influence these patterns?

Calling was associated with cloudy, warm and humid conditions and high monthly rainfall



Implications and future research

- Baseline data on calling, and thus detectability and breeding activity
- Optimal timing and conditions for active search surveys
- Bioacoustics better targeted to reduce costs and improve efficiency
- Sensitivity to accelerating climate change shift timing of calling and result in reduced breeding opportunities





Image: Harry Hine

Email: njanemcleod@gmail.com biodiversity-health.org

Special thanks to my Honours supervisors:

Harry Hines, Senior Conservation Officer Queensland Parks and Wildlife Service, Department of Environment and Science

Dr. Laura Grogan and Dr. Clare Morrison
School of Environment and Science, Griffith University