TIRZEPATIDE Bi-Monthly Kit #3 (BK3) Dosing



≈ MG per syringe

7.5mg

BK3 is for:

Weeks#

1-4

1.	Initial High-Dose (2.5 to 10mg):	BK3 is for 2 nd Bi-Monthly (after BK1)
2.	Initial Super-Dose (2.5 to 15mg):	BK3 is for 2 nd Bi-Monthly (after BK1)
3.	Transition Mid- to High-Dose	BK3 is for 1 st Bi-Monthly (before BK4)
4.	Transition Mid- to Super-Dose	BK3 is for 1 st Bi-Monthly (before BK5 & BK6)

Instructions. Fill each syringe to the unit line below.

If you need extra syringes, use 1ml Insulin Syringes, 5/16" (8mm), 31G (.25mm).

of syringes ≈ Units per syringe

75 units

Order extra syringes **HERE**. Order extra alcohol pads **HERE**.

4 syringes

BK3 transitions dosing from 7.5mg to 10mg. Includes 7ml of Tirzepatide.

If you would like to request a consultation, call us at (470) 474-1000.

5-8	4 syringes	100 units	10mg		
Remember: Months are 4-weeks. Bi-Monthly periods are 2-months or 8-weeks.					
Use MedPure's Dosing Calendar to write-down the units above on your inject-days.					
Your BK contains 2-Months (8 weeks) of Tirzepatide in vials and empty syringes.					
Check all foil & paper wrapping for your vials and empty syringes.					
To fill syringes, use <u>How to Fill Weight-Loss Syringes</u> and watch this <u>video</u> .					
Refrigerate your vials & filled-syringes & vial(s) in a sealed plastic bag or a hard-case.					
Immediately, or on your next inject-day, self-inject Week #1's syringe.					
Use <u>How to Se</u>	<u>lf-Inject</u> and watch this <u>vic</u>	<u>deo</u> .			
Then, every 7-days (weekly), on your "inject-day," self-inject your next correctly-dosed syringe.					
Properly discard any unused medication after 90 days. Do not reuse syringes.					
Do not flush medications or pour down a drain.					
Use a Sharps Container or a Bottle with Cap to dispose of used syringes.					
Keep all medic	ines out of the reach of ch	ildren.			
Do not share o	r take anyone else's medic	cine.			
Go to MedPure	e.com for side-effects, cau	tions, and <u>Terms & Conditio</u>	<u>ins</u> .		