

TIRZEPATIDE Bi-Monthly Kit #5 (BK5) Dosing

BK5 is used for these Tirzepatide Memberships:

1. Initial Super-Dose (2.5 to 15mg): **BK5** is for 3rd Bi-Monthly (after BK1 & BK3).
 2. Transition Mid- to Super-Dose (7.5 to 15mg): **BK5** is 2nd Bi-Monthly (after BK3).
-

Instructions. Fill each syringe to the unit line below.

If you need extra syringes, use 1ml Insulin Syringes, 5/16” (8mm), 31G (.25mm).

Order extra syringes [HERE](#). Order extra alcohol pads [HERE](#).

BK5 is for a transition from High-Dose to Super-Dose for 8 weeks. Includes 11ml of Tirzepatide.

<u>Weeks #</u>	<u># of syringes</u>	<u>≈ Units per syringe</u>	<u>≈ MG per syringe</u>
1-4	8 syringes*	2 X 62.5 units	12.5mg
5-8	8 syringes*	2 X 75 units	15mg

* 2 syringes are required for each weekly dose.

-
- Remember: Months are 4-weeks. Bi-Monthly periods are 2-months or 8-weeks.
 - Use MedPure’s [Dosing Calendar](#) to write-down the units above on your inject-days.
 - Your **BK** contains 2-Months (8 weeks) of Tirzepatide in vials and empty syringes.
 - Check all foil & paper wrapping for your vials and empty syringes.**
 - To fill syringes, use [How to Fill Weight-Loss Syringes](#) and watch this [video](#).
 - Refrigerate your vials & filled-syringes & vial(s) in a sealed plastic bag or a [hard-case](#).
 - Immediately, or on your next inject-day, self-inject Week #1’s syringe.
 - Use [How to Self-Inject](#) and watch this [video](#).
 - Then, every 7-days (weekly), on your “inject-day,” self-inject your next correctly-dosed syringe.
 - Properly discard any unused medication after 90 days. Do not reuse syringes.
 - Do not flush medications or pour down a drain.
 - Use a Sharps Container or a Bottle with Cap to dispose of used syringes.
 - Keep all medicines out of the reach of children.
 - Do not share or take anyone else’s medicine.
 - Go to MedPure.com for side-effects, cautions, and [Terms & Conditions](#).
 - If you would like to request a consultation, call us at (470) 474-1000.