

TIRZEPATIDE Bi-Monthly Kit Maintenance (BKM)

BKM is used for this one Membership:

1. Maintenance Fixed Dose (5mg):

BKM is used after goal-weight has been achieved to prevent weight-rebound and to keep health & medical benefits. **BKM** is only available to existing Members of MedPure.

Instructions. Fill each syringe to the unit line below.

If you need extra syringes, use 1ml Insulin Syringes, 5/16” (8mm), 31G (.25mm).

Order extra syringes [HERE](#). Order extra alcohol pads [HERE](#).

BKM is a fixed dose of 5mg for all 8 weeks (Bi-Monthly). Includes 4ml of Tirzepatide.

<u>Weeks #</u>	<u># of syringes</u>	<u>≈ Units per syringe</u>	<u>≈ MG per syringe</u>
1-12	8 syringes	50 units	5mg

- Remember: Months are 4-weeks. Bi-Monthly periods are 2-months or 8-weeks.
- Use MedPure’s [Dosing Calendar](#) to write-down the units above on your inject-days.
- Your **BK** contains 2-Months (8 weeks) of Tirzepatide in vials and empty syringes.
- Check all foil & paper wrapping for your vials and empty syringes.**
- To fill syringes, use [How to Fill Weight-Loss Syringes](#) and watch this [video](#).
- Refrigerate your vials & filled-syringes & vial(s) in a sealed plastic bag or a [hard travel-case](#).
- Immediately, or on your next inject-day, self-inject Week #1’s syringe.
- Use [How to Self-Inject](#) and watch this [video](#).
- Then, every 7-days (weekly), on your “inject-day,” self-inject your next correctly-dosed syringe.
- Properly discard any unused medication after 90 days. Do not reuse syringes.
- Do not flush medications or pour down a drain.
- Use a Sharps Container or a Bottle with Cap to dispose of used syringes.
- Keep all medicines out of the reach of children.
- Do not share or take anyone else’s medicine.
- Go to MedPure.com for side-effects, cautions, and [Terms & Conditions](#).
- If you would like to request a consultation, call us at (470) 474-1000.