

SEMAGLUTIDE Daily DROPS™ Super-Dose Dosing

Members are shipped 3-months worth of Semaglutide Daily DROPS for the doses below. Take the following amounts daily. There are 28 days a month. Quarters are 3-months or 12-weeks. The soft filler-syringe is marked in 1/10th of a ml up to 1ml.

<u>Week #</u>	<u># of Days</u>	<u>≈ ml per day</u>	<u>≈ MG per day</u>
1-12	84 days	.5 ml	2.5mg

- Refrigerate the tightly-closed Semaglutide DROPS bottle.
- Each day, place under your tongue the above ml per dose.
- Use the Shipped Bottle for your current Membership. There are 2 different Bottles (2mg/ml & 5mg/ml).
- IMPORTANT: For this Super-Dose Dosing use the bottle marked as 5mg/ml.**

How to take the correct dose every day

- Replace the shipped screw-on cap with the enclosed blue plastic cap. Tighten securely. Always keep on.
- Push the top small cap on the larger cap. Push down to seal. Store this way.
- Rigorously shake the bottle with your thumb on top before every use.
- To draw-out your dose:
 - Remove the top smaller cap.
 - Pull-in a bit of air into the soft filler-syringe (to later push-out all the medicine).
 - Push the soft filler-syringe into the opening in the cap on the bottle.
 - Turn the bottle upside down while keeping in the soft syringe.
 - Pull-out the desired dose of the medicine (measure just the medicine).
 - Push the small cap back on the larger cap.
 - Make sure both caps are tightly closed.
- Place the tip of the syringe under your tongue. Push the plunger all the way to insert all the medicine.
- Deliberately hold the medicine under your tongue for at least 1 to 2 minutes.
- Do not drink or eat for at least 5-minutes to allow the medicine to fully absorb under your tongue.
- Rinse out the soft filler-syringe in water. Shake out and let dry. When dry, replace cap on soft syringe.
- Store the bottle and filler-syringes in re-sealable plastic bags. Keep refrigerated.
- Properly discard unused medication after 90 days.
- Do not flush medications or pour down a drain. Keep all medicines out of the reach of children.
- Go to [MedPure.com](https://www.MedPure.com) for side-effects, cautions, and [Terms & Conditions](#).
- If you would like to request a consultation, call your Primary Consultant or (470) 474-1000.

Side-effects may include nausea, acid reflux, diarrhea, constipation, and vomiting but usually decrease over time. See [MedPure.com](https://www.MedPure.com) for a more complete list and how to reduce them. Seek medical attention if severe abdominal pain or allergic reaction. Rodent studies have shown risk of Thyroid C-Cell Tumors (but not in human-studies). Avoid if personal or family history of MTC or MEN 2).