SEMAGLUTIDE Quarterly Kit Maintenance (QKM)



QKM is used for this one Membership:

1. Maintenance Fixed Dose (.6mg):

QKM is used after goal-weight has been achieved to prevent weight-rebound and to keep health & medical benefits. **QKM** is only available to existing Members of MedPure.

Instructions. Fill each syringe to the unit line below.

If you need extra syringes, use 1ml Insulin Syringes, 5/16" (8mm), 31G (.25mm). Order extra syringes <u>HERE</u>. Order extra alcohol pads <u>HERE</u>.

QKM is a fixed dose of .6mg for all 12 weeks (Quarter). Includes 3ml of Semaglutide.

Weeks #	<u># of syringes</u>	<u>≈ Units per syringe</u>	<u>≈ MG per syringe</u>
1-12	12 syringes	25 units	.6mg

- Remember: Months are 4-weeks. Quarters are 3-months or 12-weeks.
- Use MedPure's <u>Dosing Calendar</u> to write-down the units above on your inject-days.
- □ Your **QK** contains 3-Months (12 weeks) of Semaglutide in vials and empty syringes.

Check all foil & paper wrapping for your vials and empty syringes.

- □ To fill syringes, use <u>How to Fill Weight-Loss Syringes</u> and watch this <u>video</u>.
- □ Refrigerate your vials & filled-syringes & vial(s) in a sealed plastic bag or a hard travel-case.
- □ Immediately, or on your next inject-day, self-inject Week #1's syringe.
- □ Use <u>How to Self-Inject</u> and watch this <u>video</u>.
- Then, every 7-days (weekly), on your "inject-day," self-inject your next correctly-dosed syringe.
- Derived any unused medication after 90 days. Do not reuse syringes.
- Do not flush medications or pour down a drain.
- Use a Sharps Container or a Bottle with Cap to dispose of used syringes.
- Keep all medicines out of the reach of children.
- Do not share or take anyone else's medicine.
- Go to MedPure.com for side-effects, cautions, and <u>Terms & Conditions</u>.
- □ If you would like to request a consultation, call us at (470) 474-1000.