

SEMAGLUTIDE w/B3 Super-Dose Dosing (2.5mg)

Who uses Semaglutide Super-Dose

1. Clients after taking High-Dose who wish to continue to maximize their weight-loss.
2. Clients with high weight-loss goals who wish to lose weight fast.
3. Clients who wish max weight-loss but do not want to pay Tirzepatide prices.
4. Clients who wish or need to add B6 to reduce side effects of regular Semaglutide.

Instructions. Fill each syringe to the unit line below. Note: 1ml = 100 units.

Clients are initially shipped 3-months supply of Semaglutide for the following doses.

| <u>Weeks #</u> | <u># of syringes</u> | <u>≈ Units per syringe</u> | <u>≈ MG per syringe</u> |
|----------------|----------------------|----------------------------|-------------------------|
| 1-12 | 12 syringes | 100 units | 2.5mg |

- ☐ There are no auto-renews or auto-ships. Clients need to re-order every 3 months.
- ☐ Remember: Months are 4-weeks. 3 Months is 12 weeks.
- ☐ If you need extra syringes, order 1ml or .5ml Insulin Syringes, 5/16" (8mm), 31G (.25mm) [HERE](#).
- ☐ Use MedPURE's [Dosing Calendar](#) to write-down the units above on your inject-days.
- ☐ Your **initial shipment** contains 3-Months (12 weeks) of Semaglutide in vials and empty syringes.
- ☐ **Check all foil & paper wrapping for your vials and empty syringes.**
- ☐ To fill syringes, use [How to Fill Weight-Loss Syringes](#) and watch this [video](#).
- ☐ Refrigerate your vials & filled-syringes & vial(s) in a sealed plastic bag or a [hard travel-case](#).
- ☐ Immediately, or on your next inject-day, self-inject Week #1's syringe.
- ☐ Use [How to Self-Inject](#) and watch this [video](#).
- ☐ Then, every 7-days (weekly), on your "inject-day," self-inject your next correctly-dosed syringe.
- ☐ Properly discard any unused medication after 90 days. Do not reuse syringes.
- ☐ Do not flush medications or pour down a drain.
- ☐ Use a Sharps Container or a Bottle with Cap to dispose of used syringes.
- ☐ Keep all medicines out of the reach of children.
- ☐ Do not share or take anyone else's medicine.
- ☐ Go to MedPURE.com for side-effects, cautions, and [Terms & Conditions](#).
- ☐ If you would like to request a consultation, call your Primary Consultant or (470) 474-1000.

Side-effects may include nausea, acid reflux, diarrhea, constipation, and vomiting but usually decrease over time. See [MedPURE.com](#) for a more complete list and how to reduce them. Seek medical attention if severe abdominal pain or allergic reaction. Rodent studies have shown risk of Thyroid C-Cell Tumors (but not in human-studies). Avoid if personal or family history of MTC or MEN 2).