

Tirzepatide PLUS B12 & Glycine High-Dose Dosing

Who uses Tirzepatide PLUS™ High-Dose

1. Members who, after taking Tirzepatide PLUS Mid-Dose, want to lose a great deal more weight.
2. Members seeking to improve their shape, skin, hair, & bone and lower side-effects as they lose weight.
3. Members who may be on their way to taking Tirzepatide PLUS Super-Dose (max FDA-allowed).

Instructions. Fill each syringe to the unit line below. Note: 1ml = 100 units.

Members are initially shipped 2-months worth of Tirzepatide PLUS for the following doses.

If using Tirzepatide High-Dose the first time after Mid-Dose, use the following dosing:

<u>Weeks #</u>	<u># of syringes</u>	<u>≈ Units per syringe</u>	<u>≈ MG per syringe</u>
1-4	4 syringes	75 units	7.5mg
4-8	4 syringes	100 units	10mg

If already transitioned to High-Dose, use the following dosing:

1-8	8 syringes	100 units	10mg
-----	------------	-----------	------

- ☐ Remember: Months are 4-weeks. Bi-Monthly is 2-months or 8-weeks.
- ☐ If you need extra syringes, order 1ml Insulin Syringes, 5/16" (8mm), 31G (.25mm) [HERE](#).
- ☐ Use MedPure's [Dosing Calendar](#) to write-down the units above on your inject-days.
- ☐ Your **initial shipment** contains 2-Months (8 weeks) of Tirzepatide PLUS in vials and empty syringes.
- ☐ **Check all foil & paper wrapping for your vials and empty syringes.**
- ☐ To fill syringes, use [How to Fill Weight-Loss Syringes](#) and watch this [video](#).
- ☐ Refrigerate your vials & filled-syringes & vial(s) in a sealed plastic bag or a [hard travel-case](#).
- ☐ Immediately, or on your next inject-day, self-inject Week #1's syringe.
- ☐ Use [How to Self-Inject](#) and watch this [video](#).
- ☐ Then, every 7-days (weekly), on your "inject-day," self-inject your next correctly-dosed syringe.
- ☐ Properly discard any unused medication after 90 days. Do not reuse syringes.
- ☐ Do not flush medications or pour down a drain.
- ☐ Use a Sharps Container or a Bottle with Cap to dispose of used syringes.
- ☐ Keep all medicines out of the reach of children. Do not share or take anyone else's medicine.
- ☐ Go to MedPure.com for side-effects, cautions, and [Terms & Conditions](#).

Side-effects may include nausea, acid reflux, diarrhea, constipation, and vomiting but usually decrease over time. See [MedPure.com](#) for a more complete list and how to reduce them. Seek medical attention if severe abdominal pain or allergic reaction. Rodent studies have shown risk of Thyroid C-Cell Tumors (but not in human-studies). Avoid if personal or family history of MTC or MEN 2).