

Tirzepatide PLUS B12 & Glycine Maintenance Dosing

Who uses PLUS Maintenance

1. Members who have already achieved their weight-loss goals and want to keep it off at a low-cost.
2. Members who wish to keep the many medical benefits of Tirzepatide, B12, & Glycine.
3. Members with low weight-loss goals who wish to have a little more weight-loss at lower-cost.

Note: To be on Maintenance, Members must take at least 4-months of any MedPure Memberships.

Instructions. Fill each syringe to the unit line below. Note: 1ml = 100 units.

Members are initially shipped 2-months worth of Tirzepatide PLUS for the following doses.

| <u>Weeks #</u> | <u># of syringes</u> | <u>≈ Units per syringe</u> | <u>≈ MG per syringe</u> |
|----------------|----------------------|----------------------------|-------------------------|
| 1-8 | 8 syringes | 25 units | 2.5mg |

- ☐ Remember: Months are 4-weeks. Bi-Monthly is 2-months or 8-weeks.
- ☐ If you need extra syringes, order 1ml Insulin Syringes, 5/16" (8mm), 31G (.25mm) [HERE](#).
- ☐ Use MedPure's [Dosing Calendar](#) to write-down the units above on your inject-days.
- ☐ Your **shipment** contains 2-Months (8 weeks) of Tirzepatide PLUS in vials and empty syringes.
- ☐ **Check all foil & paper wrapping for your vials and empty syringes.**
- ☐ To fill syringes, use [How to Fill Weight-Loss Syringes](#) and watch this [video](#).
- ☐ Refrigerate your vials & filled-syringes & vial(s) in a sealed plastic bag or a [hard travel-case](#).
- ☐ Immediately, or on your next inject-day, self-inject Week #1's syringe.
- ☐ Use [How to Self-Inject](#) and watch this [video](#).
- ☐ Then, every 7-days (weekly), on your "inject-day," self-inject your next correctly-dosed syringe.
- ☐ Properly discard any unused medication after 90 days. Do not reuse syringes.
- ☐ Do not flush medications or pour down a drain.
- ☐ Use a Sharps Container or a Bottle with Cap to dispose of used syringes.
- ☐ Keep all medicines out of the reach of children.
- ☐ Do not share or take anyone else's medicine.
- ☐ Go to MedPure.com for side-effects, cautions, and [Terms & Conditions](#).
- ☐ If you would like to request a consultation, call your Primary Consultant or (470) 474-1000.

Side-effects may include nausea, acid reflux, diarrhea, constipation, and vomiting but usually decrease over time. See [MedPure.com](#) for a more complete list and how to reduce them. Seek medical attention if severe abdominal pain or allergic reaction. Rodent studies have shown risk of Thyroid C-Cell Tumors (but not in human-studies). Avoid if personal or family history of MTC or MEN 2).