

TIRZEPATIDE (New with B6) MID-DOSE Dosing (5mg)

Note: New Tirzepatide is 2X the concentration as before and requires only ½ the units for the same mgs.

Who uses Tirzepatide Mid-Dose

1. Clients who complete 3-months of Tirzepatide FAST-START Dosing may renew to Mid-Dose at \$195.
2. Clients with lower weight-loss goals may wish to stay at Mid-Dose (5mg) for lower cost (\$195).
3. Clients with higher weight-loss goals may wish to increase to High-Dose (10mg) at \$395.
4. After 6-months, Clients may choose Super-Dose (12.5mg) at \$595/mo.

Instructions. Fill each syringe to the unit line below. 1ml = 100 units.

Clients are initially shipped 3-months worth of Tirzepatide for the following dose:

<u>Weeks #</u>	<u># of syringes</u>	<u>≈ Units per syringe</u>	<u>≈ MG per syringe</u>
1-12	12 syringes	25 units	5mg

Remember: The new units are ½ prior units due to 2X concentration.

- There are no auto-renews or auto-ships. Clients need to re-order every 3 months.
- Remember: Months are 4-weeks. 3 Months is 12 weeks.
- If you need extra syringes, order 1ml Insulin Syringes, 5/16" (8mm), 31G (.25mm) [HERE](#).
- Use MedPure's [Dosing Calendar](#) to write-down the units above on your inject-days.
- Your **initial shipment** contains 3-Months (12 weeks) of Tirzepatide in vials and empty syringes.
- Check all foil & paper wrapping for your vials and empty syringes.**
- To fill syringes, use [How to Fill Weight-Loss Syringes](#) and watch this [video](#).
- Refrigerate your vials & filled-syringes & vial(s) in a sealed plastic bag or a [hard travel-case](#).
- Immediately, or on your next inject-day, self-inject Week #1's syringe.
- Use [How to Self-Inject](#) and watch this [video](#).
- Then, every 7-days (weekly), on your "inject-day," self-inject your next correctly-dosed syringe.
- Properly discard any unused medication after 90 days. Do not reuse syringes.
- Do not flush medications or pour down a drain.
- Use a Sharps Container or a Bottle with Cap to dispose of used syringes.
- Keep all medicines out of the reach of children.
- Do not share or take anyone else's medicine.
- Go to [MedPure.com](#) for side-effects, cautions, and [Terms & Conditions](#).
- If you would like to request a consultation, call your Consultant or (470) 474-1000.

Side-effects may include nausea, acid reflux, diarrhea, constipation, and vomiting but usually decrease over time. See [MedPure.com](#) for a more complete list and how to reduce them. Seek medical attention if severe abdominal pain or allergic reaction. Rodent studies have shown risk of Thyroid C-Cell Tumors (but not in human-studies). Avoid if personal or family history of MTC or MEN 2).