



TIRZEPATIDE MID-DOSE SPECIAL Dosing (5mg)

Note: New Tirzepatide is 2X the concentration as before and has B6 and requires only ½ the units for the same mgs.

Who uses Tirzepatide Mid-Dose SPECIAL

1. Tirzep Mid-Dose SPECIAL is offered 1-time to prior MedPure members at the very low price of \$195/month.
2. Unlike prior Tirzepatide Memberships, new Tirzepatide Memberships are for 3-Months.
3. After this SPECIAL, Members may wish to stay at Mid-Dose (5mg) for the regular cost of \$295/month.
4. Members with higher weight-loss goals may wish to increase to High-Dose (10mg) at \$495.
5. After HIGH-DOSE, Members may choose Super-Dose (12.5mg) at \$695/mo.

Instructions. Fill each syringe to the unit line below. 1ml = 100 units.

Members are initially shipped 3-months supply of Tirzepatide for the following dose:

<u>Weeks #</u>	<u># of syringes</u>	<u>≈ Units per syringe</u>	<u>≈ MG per syringe</u>
1-12	12 syringes	50 units*	5mg

Remember: The new units are ½ prior units due to 2X concentration.

-
- Remember: Months are 4-weeks. Three months are 12 weeks.
 - If you need extra syringes, order 1ml Insulin Syringes, 5/16" (8mm), 31G (.25mm) [HERE](#).
 - Use MedPure's [Dosing Calendar](#) to write-down the units above on your inject-days.
 - Your **initial shipment** contains 3-Months (12 weeks) of Tirzepatide in vials and empty syringes.
 - Check all foil & paper wrapping for your vials and empty syringes.**
 - To fill syringes, use [How to Fill Weight-Loss Syringes](#) and watch this [video](#).
 - Refrigerate your vials & filled-syringes & vial(s) in a sealed plastic bag or a [hard travel-case](#).
 - Immediately, or on your next inject-day, self-inject Week #1's syringe.
 - Use [How to Self-Inject](#) and watch this [video](#).
 - Then, every 7-days (weekly), on your "inject-day," self-inject your next correctly-dosed syringe.
 - Properly discard any unused medication after 90 days. Do not reuse syringes.
 - Do not flush medications or pour down a drain.
 - Use a Sharps Container or a Bottle with Cap to dispose of used syringes.
 - Keep all medicines out of the reach of children.
 - Do not share or take anyone else's medicine.
 - Go to MedPure.com for side-effects, cautions, and [Terms & Conditions](#).
 - If you would like to request a consultation, call your Consultant or (470) 474-1000.

Side-effects may include nausea, acid reflux, diarrhea, constipation, and vomiting but usually decrease over time. See [MedPure.com](#) for a more complete list and how to reduce them. Seek medical attention if severe abdominal pain or allergic reaction. Rodent studies have shown risk of Thyroid C-Cell Tumors (but not in human-studies). Avoid if personal or family history of MTC or MEN 2).