

## TIRZEPATIDE (New with B6) FAST-START Dosing

Note: New Tirzepatide is 2X the concentration as before and requires only ½ the units for the same mgs.

### Who uses Tirzepatide FAST-START Dosing

1. Clients starting dual GLP-1 & GIP certified Tirzepatide to lose the most weight quickly.
2. Clients seeking to minimize their side-effects as they lose weight.
3. In 3-months, Clients titrate (increase dose) from start-dose of 2.5mg to 7.5mg at only \$195 a month.
4. After 3-months, Clients may renew to Mid-Dose (5mg) at \$195, or increase to High-Dose (10mg) at \$395/mo.
5. After that, Clients may renew to Super-Dose (12.5mg) at \$595/mo for fastest & greatest weight-loss.

**Instructions.** Fill each syringe to the unit line below. Note: 1ml = 100 units.

Clients are initially shipped 3-months supply of Tirzepatide for the following doses:

<b>Weeks #</b>	<b># of syringes</b>	<b>≈ Units per syringe</b>	<b>≈ MG per syringe</b>
1-4	4 syringes	12 units	2.5mg
5-8	4 syringes	25 units	5mg
9-12	4 syringes	37 units	7.5mg

Remember: The new units are ½ prior units due to 2X concentration.

- There are no auto-renews or auto-ships. Clients need to re-order every 3 months.
- Remember: Months are 4-weeks. 3 Months is 12 weeks.
- Use MedPure's [Dosing Calendar](#) to write-down the units above on your inject-days.
- Your **initial shipment** contains 3-Months (12 weeks) of Tirzepatide in vials and empty syringes.
- Check all foil & paper wrapping for your vials and empty syringes.**
- To fill syringes, use [How to Fill Weight-Loss Syringes](#) and watch this [video](#).
- Refrigerate your vials & filled-syringes & vial(s) in a sealed plastic bag or a [hard travel-case](#).
- Immediately, or on your next inject-day, self-inject Week #1's syringe.
- Use [How to Self-Inject](#) and watch this [video](#).
- Then, every 7-days (weekly), on your "inject-day," self-inject your next correctly-dosed syringe.
- Properly discard any unused medication after 90 days. Do not reuse syringes.
- Do not flush medications or pour down a drain.
- Use a Sharps Container or a Bottle with Cap to dispose of used syringes.
- Keep all medicines out of the reach of children.
- Do not share or take anyone else's medicine.
- Go to MedPure.com for side-effects, cautions, and [Terms & Conditions](#).
- If you would like to request a consultation, call your Primary Consultant or (470) 474-1000.

Side-effects may include nausea, acid reflux, diarrhea, constipation, and vomiting but usually decrease over time. See [MedPure.com](#) for a more complete list and how to reduce them. Seek medical attention if severe abdominal pain or allergic reaction. Rodent studies have shown risk of Thyroid C-Cell Tumors (but not in human-studies). Avoid if personal or family history of MTC or MEN 2).