

TIRZEPATIDE (New with B6) FAST-START Dosing

Note: New Tirzepatide is 2X the concentration as before and requires only ½ the units for the same mgs.

Who uses Tirzepatide FAST-START Dosing

1. Members starting dual GLP-1 & GIP certified Tirzepatide to lose the most weight quickly.
2. Members seeking to minimize their side-effects as they lose weight.
3. In 3-months, Members titrate (increase dose) from start-dose of 2.5mg to 7.5mg at only \$195 a month.
4. After 3-months, Members may renew to Mid-Dose (5mg) at \$295, or increase to High-Dose (10mg) at \$495/mo.
5. After that, Members may renew to Super-Dose (12.5mg) at \$695/mo for fastest & greatest weight-loss.

Instructions. Fill each syringe to the unit line below. Note: 1ml = 100 units.

Members are initially shipped 3-months supply of Tirzepatide for the following doses:

<u>Weeks #</u>	<u># of syringes</u>	<u>≈ Units per syringe</u>	<u>≈ MG per syringe</u>
1-4	4 syringes	12 units	2.5mg
5-8	4 syringes	25 units	5mg
9-12	4 syringes	37 units	7.5mg

Remember: The new units are ½ prior units due to 2X concentration.

- ☐ Remember: Months are 4-weeks. 3 Months is 12 weeks.
- ☐ Use MedPure's [Dosing Calendar](#) to write-down the units above on your inject-days.
- ☐ Your **initial shipment** contains 3-Months (12 weeks) of Tirzepatide in vials and empty syringes.
- ☐ **Check all foil & paper wrapping for your vials and empty syringes.**
- ☐ To fill syringes, use [How to Fill Weight-Loss Syringes](#) and watch this [video](#).
- ☐ Refrigerate your vials & filled-syringes & vial(s) in a sealed plastic bag or a [hard travel-case](#).
- ☐ Immediately, or on your next inject-day, self-inject Week #1's syringe.
- ☐ Use [How to Self-Inject](#) and watch this [video](#).
- ☐ Then, every 7-days (weekly), on your "inject-day," self-inject your next correctly-dosed syringe.
- ☐ Properly discard any unused medication after 90 days. Do not reuse syringes.
- ☐ Do not flush medications or pour down a drain.
- ☐ Use a Sharps Container or a Bottle with Cap to dispose of used syringes.
- ☐ Keep all medicines out of the reach of children.
- ☐ Do not share or take anyone else's medicine.
- ☐ Go to MedPure.com for side-effects, cautions, and [Terms & Conditions](#).
- ☐ If you would like to request a consultation, call your Primary Consultant or (470) 474-1000.

Side-effects may include nausea, acid reflux, diarrhea, constipation, and vomiting but usually decrease over time. See [MedPure.com](#) for a more complete list and how to reduce them. Seek medical attention if severe abdominal pain or allergic reaction. Rodent studies have shown risk of Thyroid C-Cell Tumors (but not in human-studies). Avoid if personal or family history of MTC or MEN 2).