

Reviewer Disclosures

Reviewer	Employment	Research Grant	Other Research Support	Speakers' Bureau/Honoraria	Expert Witness	Ownership Interest	Consultant/Advisory Board	Other
Joshua Beckman	Brigham and Women's Hospital	None	None	None	None	None	Novartis*	None
Joanne Murabito	Boston University Medical Center and Framingham Heart Study	NIH/NHLBI†	None	None	None	None	None	None
Roberta Oka	VA Palo Alto HCS	None	None	None	None	None	None	None

This table represents the relationships of reviewer that may be perceived as actual or reasonably perceived conflicts of interest as reported on the Disclosure Questionnaire, which all reviewers are required to complete and submit. A relationship is considered to be "significant" if (a) the person receives \$10 000 or more during any 12-month period, or 5% or more of the person's gross income; or (b) the person owns 5% or more of the voting stock or share of the entity, or owns \$10 000 or more of the fair market value of the entity. A relationship is considered to be "modest" if it is less than "significant" under the preceding definition.

*Modest.

†Significant.

References

- Winsor T. Influence of arterial disease on the systolic blood pressure gradients of the extremity. *Am J Med Sci.* 1950;220:117–126.
- Carter SA. Indirect systolic pressures and pulse waves in arterial occlusive diseases of the lower extremities. *Circulation.* 1968;37:624–637.
- Yao ST, Hobbs JT, Irvine WT. Ankle systolic pressure measurements in arterial disease affecting the lower extremities. *Br J Surg.* 1969;56:676–679.
- Criqui MH, Langer RD, Fronek A, Feigelson HS, Klauber MR, McCann TJ, Browner D. Mortality over a period of 10 years in patients with peripheral arterial disease. *N Engl J Med.* 1992;326:381–386.
- McDermott MM, Guralnik JM, Tian L, Liu K, Ferrucci L, Liao Y, Sharma L, Criqui MH. Associations of borderline and low normal ankle-brachial index values with functional decline at 5-year follow-up: the WALCS (Walking and Leg Circulation Study). *J Am Coll Cardiol.* 2009;53:1056–1062.
- Ankle Brachial Index Collaboration, Fowkes FG, Murray GD, Butcher I, Heald CL, Lee RJ, Chambless LE, Folsom AR, Hirsch AT, Dramaix M, deBacker G, Wautrecht JC, Kornitzer M, Newman AB, Cushman M, Sutton-Tyrrell K, Lee AJ, Price JF, d'Agostino RB, Murabito JM, Norman PE, Jamrozik K, Curb JD, Masaki KH, Rodriguez BL, Dekker JM, Bouter LM, Heine RJ, Nijpels G, Stehouwer CD, Ferrucci L, McDermott MM, Stoffers HE, Hooi JD, Knottnerus JA, Ogren M, Hedblad B, Witteman JC, Breteler MM, Hunink MG, Hofman A, Criqui MH, Langer RD, Fronek A, Hiatt WR, Hamman R, Resnick HE, Guralnik J. Ankle brachial index combined with Framingham risk score to predict cardiovascular events and mortality: a meta-analysis. *JAMA.* 2008;300:197–208.
- Lange SF, Trampisch HJ, Pittrow D, Darius H, Mahn M, Allenberg JR, Tepohl G, Haberl RL, Diehm C; getABI Study Group. Profound influence of different methods for determination of the ankle brachial index on the prevalence estimate of peripheral arterial disease. *BMC Public Health.* 2007;147.
- Aboyans V, Lacroix P, Preux PM, Vergnenegre A, Ferrieres J, Laskar M. Variability of ankle-arm index in general population according to its mode of calculation. *Int Angiol.* 2002;21:237–243.
- Allison MA, Aboyans V, Granston T, McDermott MM, Kamineni A, Ni H, Criqui MH. The relevance of different methods of calculating the ankle-brachial index: the Multi-Ethnic Study of Atherosclerosis. *Am J Epidemiol.* 2010;171:368–376.
- Klein S, Hage JJ. Measurement, calculation, and normal range of the ankle-arm index: a bibliometric analysis and recommendation for standardization. *Ann Vasc Surg.* 2006;20:282–292.
- Dachun X, Jue L, Liling Z, Yawei X, Dayi H, Pagoto SL, Yunsheng M. Sensitivity and specificity of the ankle-brachial index to diagnose peripheral artery disease: a structured review. *Vasc Med.* 2010;15:361–369.
- Stein JH, Korcarz CE, Hurst RT, Lonn E, Kendall CB, Mohler ER, Najjar SS, Rembold CM, Post WS; American Society of Echocardiography Carotid Intima-Media Thickness Task Force. Use of carotid ultrasound to identify subclinical vascular disease and evaluate cardiovascular disease risk: a consensus statement from the American Society of Echocardiography Carotid Intima-Media Thickness Task Force: endorsed by the Society for Vascular Medicine. *J Am Soc Echocardiogr.* 2008;21:93–111; quiz 189–190.
- Greenland P, Bonow RO, Brundage BH, Budoff MJ, Eisenberg MJ, Grundy SM, Lauer MS, Post WS, Raggi P, Redberg RF, Rodgers GP, Shaw LJ, Taylor AJ, Weintraub WS. ACCF/AHA 2007 clinical expert consensus document on coronary artery calcium scoring by computed tomography in global cardiovascular risk assessment and in evaluation of patients with chest pain: a report of the American College of Cardiology Foundation Clinical Expert Consensus Task Force (ACCF/AHA Writing Committee to Update the 2000 Expert Consensus Document on Electron Beam Computed Tomography) developed in collaboration with the Society of Atherosclerosis Imaging and Prevention and the Society of Cardiovascular Computed Tomography. *Circulation.* 2007;115:402–426.
- Hiatt WR, Goldstone J, Smith SC Jr, McDermott M, Moneta G, Oka R, Newman AB, Pearce WH; American Heart Association Writing Group 1. Atherosclerotic peripheral vascular disease symposium II: nomenclature for vascular diseases. *Circulation.* 2008;118:2826–2829.
- Safar ME, Protogerou AD, Blacher J, Statins, central blood pressure, and blood pressure amplification. *Circulation.* 2009;119:9–12.
- Murgo JP, Westerhof N, Giolma JP, Altabelli SA. Aortic input impedance in normal man: relationship to pressure wave forms. *Circulation.* 1980;62:105–116.
- Latham RD, Westerhof N, Sipkema P, Rubal BJ, Reuderink P, Murgo JP. Regional wave travel and reflections along the human aorta: a study with six simultaneous micromanometric pressures. *Circulation.* 1985;72:1257–1269.
- Hope SA, Tay DB, Meredith IT, Cameron JD. Waveform dispersion, not reflection, may be the major determinant of aortic pressure wave morphology. *Am J Physiol Heart Circ Physiol.* 2005;289:H2497–H2502.
- Wang JJ, Parker KH. Wave propagation in a model of the arterial circulation. *J Biomech.* 2004;37:457–470.
- Tsami A, Stergiopoulos N. Arterial remodeling in response to hypertension using a constituent-based model. *Am J Physiol Heart Circ Physiol.* 2007;293:H3130–H3139.
- Humphrey JD. Mechanisms of arterial remodeling in hypertension: coupled roles of wall shear and intramural stress. *Hypertension.* 2008;52:195–200.
- Katz S, Globerman A, Avitzour M, Dolfin T. The ankle-brachial index in normal neonates and infants is significantly lower than in older children and adults. *J Pediatr Surg.* 1997;32:269–271.
- Smith FB, Lee AJ, Price JF, van Wijk MC, Fowkes FG. Changes in ankle brachial index in symptomatic and asymptomatic subjects in the general population. *J Vasc Med.* 2003;38:1323–1330.
- Hiatt WR, Hoag S, Hamman RF. Effect of diagnostic criteria on the prevalence of peripheral arterial disease: the San Luis Valley Diabetes Study. *Circulation.* 1995;91:1472–1479.
- Bird CE, Criqui MH, Fronek A, Denenberg JO, Klauber MR, Langer RD. Quantitative and qualitative progression of peripheral arterial disease by non-invasive testing. *Vasc Med.* 1999;4:15–21.
- London GM, Guerin AP, Pannier B, Marchais SJ, Stimpel M. Influence of sex on arterial hemodynamics and blood pressure: role of body height. *Hypertension.* 1995;26:514–519.
- Aboyans V, Criqui MH, McClelland RL, Allison MA, McDermott MM, Goff DC Jr, Manolio TA. Intrinsic contribution of gender and ethnicity to normal ankle-brachial index values: the Multi-Ethnic Study of Atherosclerosis (MESA). *J Vasc Med.* 2007;45:319–327.
- Stoffers HE, Kester AD, Kaiser V, Rinkens PE, Kitslaar PJ, Knottnerus JA. The diagnostic value of the measurement of the ankle-brachial

- systolic pressure index in primary health care. *J Clin Epidemiol*. 1996; 49:1401–1405.
29. Zheng ZJ, Sharrett AR, Chambless LE, Rosamond WD, Nieto FJ, Sheps DS, Dobs A, Evans GW, Heiss G. Associations of ankle-brachial index with clinical coronary heart disease, stroke and preclinical carotid and popliteal atherosclerosis: the Atherosclerosis Risk in Communities (ARIC) Study. *Atherosclerosis*. 1997;131:115–125.
 30. Zheng ZJ, Rosamond WD, Chambless LE, Nieto FJ, Barnes RW, Hutchinson RG, Tyroler HA, Heiss G; ARIC Investigators. Lower extremity arterial disease assessed by ankle-brachial index in a middle-aged population of African Americans and whites: the Atherosclerosis Risk in Communities (ARIC) Study. *Am J Prev Med*. 2005; 29(suppl 1):42–49.
 31. Carmelli D, Fabsitz RR, Swan GE, Reed T, Miller B, Wolf PA. Contribution of genetic and environmental influences to ankle-brachial blood pressure index in the NHLBI Twin Study: National Heart, Lung, and Blood Institute. *Am J Epidemiol*. 2000;151:452–458.
 32. Allison MA, Peralta CA, Wassel CL, Aboyans V, Arnett DK, Cushman M, Eng J, Ix J, Rich SS, Criqui MH. Genetic ancestry and lower extremity peripheral artery disease in the Multi-Ethnic Study of Atherosclerosis. *Vasc Med*. 2010;15:351–359.
 33. Su HM, Lee KT, Chu CS, Lee MY, Lin TH, Voon WC, Sheu SH, Lai WT. Effects of heart rate on brachial-ankle pulse wave velocity and ankle-brachial pressure index in patients without significant organic heart disease. *Angiology*. 2007;58:67–74.
 34. Wilkinson IB, MacCallum H, Flint L, Cockcroft JR, Newby DE, Webb DJ. The influence of heart rate on augmentation index and central arterial pressure in humans. *J Physiol*. 2000;525(pt 1):263–270.
 35. Abraham P, Desvaux B, Colin D, Leftheriotis G, Saumet JL. Heart rate-corrected ankle-to-arm index in the diagnosis of moderate lower extremity arterial disease. *Angiology*. 1995;46:673–677.
 36. Su HM, Chang JM, Lin FH, Chen SC, Voon WC, Cheng KH, Wang CS, Lin TH, Lai WT, Sheu SH. Influence of different measurement time points on brachial-ankle pulse wave velocity and ankle-brachial index in hemodialysis patients. *Hypertens Res*. 2007;30:965–970.
 37. Allen J, Oates CP, Henderson J, Jago J, Whittingham TA, Chamberlain J, Jones NA, Murray A. Comparison of lower limb arterial assessments using color-duplex ultrasound and ankle/brachial pressure index measurements. *Angiology*. 1996;47:225–232.
 38. Lijmer JG, Hunink MG, van den Dungen JJ, Loostra J, Smit AJ. ROC analysis of noninvasive tests for peripheral arterial disease. *Ultrasound Med Biol*. 1996;22:391–398.
 39. Niazi K, Khan TH, Easley KA. Diagnostic utility of the two methods of ankle brachial index in the detection of peripheral arterial disease of lower extremities. *Catheter Cardiovasc Interv*. 2006;68:788–792.
 40. Ouriel K, McDonnell AE, Metz CE, Zarins CK. Critical evaluation of stress testing in the diagnosis of peripheral vascular disease. *Surgery*. 1982;91:686–693.
 41. Ouriel K, Zarins CK. Doppler ankle pressure: an evaluation of three methods of expression. *Arch Surg*. 1982;117:1297–1300.
 42. Parameswaran GI, Brand K, Dolan J. Pulse oximetry as a potential screening tool for lower extremity arterial disease in asymptomatic patients with diabetes mellitus. *Arch Intern Med*. 2005;165:442–446.
 43. Premalatha G, Ravikumar R, Sanjay R, Deepa R, Mohan V. Comparison of colour duplex ultrasound and ankle-brachial pressure index measurements in peripheral vascular disease in type 2 diabetic patients with foot infections. *J Assoc Physicians India*. 2002;50:1240–1244.
 44. Schroder F, Diehm N, Kareem S, Ames M, Pira A, Zwettler U, Lawall H, Diehm C. A modified calculation of ankle-brachial pressure index is far more sensitive in the detection of peripheral arterial disease. *J Vasc Surg*. 2006;44:531–536.
 45. Sumner DS, Strandness DE Jr. The relationship between calf blood flow and ankle blood pressure in patients with intermittent claudication. *Surgery*. 1969;65:763–771.
 46. Williams DT, Harding KG, Price P. An evaluation of the efficacy of methods used in screening for lower-limb arterial disease in diabetes. *Diabetes Care*. 2005;28:2206–2210.
 47. Alnaeb ME, Crabtree VP, Boutin A, Mikhailidis DP, Seifalian AM, Hamilton G. Prospective assessment of lower-extremity peripheral arterial disease in diabetic patients using a novel automated optical device. *Angiology*. 2007;58:579–585.
 48. Clairrotte C, Retout S, Potier L, Roussel R, Escoubet B. Automated ankle-brachial pressure index measurement by clinical staff for peripheral arterial disease diagnosis in nondiabetic and diabetic patients. *Diabetes Care*. 2009;32:1231–1236.
 49. Feigelson HS, Criqui MH, Fronek A, Langer RD, Molgaard CA. Screening for peripheral arterial disease: the sensitivity, specificity, and predictive value of noninvasive tests in a defined population. *Am J Epidemiol*. 1994;140:526–534.
 50. Guo X, Li J, Pang W, Zhao M, Luo Y, Sun Y, Hu D. Sensitivity and specificity of ankle-brachial index for detecting angiographic stenosis of peripheral arteries. *Circ J*. 2008;72:605–610.
 51. Wikstrom J, Hansen T, Johansson L, Lind L, Ahlstrom H. Ankle brachial index <0.9 underestimates the prevalence of peripheral artery occlusive disease assessed with whole-body magnetic resonance angiography in the elderly. *Acta Radiol*. 2008;49:143–149.
 52. Baxter GM, Polak JF. Lower limb colour flow imaging: a comparison with ankle:brachial measurements and angiography. *Clin Radiol*. 1993; 47:91–95.
 53. de Groot P, Millaire A, Deklunder G, Marache P, Decouls E, Ducloux G. Comparative diagnostic value of ankle-to-brachial index and transcutaneous oxygen tension at rest and after exercise in patients with intermittent claudication. *Angiology*. 1995;46:115–122.
 54. Flanagan DP, Ballard JL, Robinson D, Galliano M, Blecker G, Harward TR. Duplex ultrasound of the superficial femoral artery is a better screening tool than ankle-brachial index to identify at risk patients with lower extremity atherosclerosis. *J Vasc Surg*. 2008;47:789–792.
 55. Alnaeb ME, Boutin A, Crabtree VP, Mikhailidis DP, Seifalian AM, Hamilton G. Assessment of lower extremity peripheral arterial disease using a novel automated optical device. *Vasc Endovascular Surg*. 2007; 41:522–527.
 56. Carter SA. Clinical measurement of systolic pressures in limbs with arterial occlusive disease. *JAMA*. 1969;207:1869–1874.
 57. Bernstein EF, Fronek A. Current status of noninvasive tests in the diagnosis of peripheral arterial disease. *Surg Clin North Am*. 1982;62: 473–487.
 58. Carter SA. Response of ankle systolic pressure to leg exercise in mild or questionable arterial disease. *N Engl J Med*. 1972;287:578–582.
 59. Winsor T. Conditioned vasoconstrictive responses of digital vessels. *AMA Arch Surg*. 1958;76:193–199.
 60. Laing S, Greenhalgh RM. The detection and progression of asymptomatic peripheral arterial disease. *Br J Surg*. 1983;70:628–630.
 61. Sakurai T, Matsushita M, Nishikimi N, Nimura Y. Effect of walking distance on the change in ankle-brachial pressure index in patients with intermittent claudication. *Eur J Vasc Endovasc Surg*. 1997;13:486–490.
 62. Hoogeveen EK, Mackaay AJ, Beks PJ, Kostense PJ, Dekker JM, Heine RJ, Nijpels G, Rauwerda JA, Stehouwer CD. Evaluation of the one-minute exercise test to detect peripheral arterial disease. *Eur J Clin Invest*. 2008;38:290–295.
 63. McPhail IR, Spittel PC, Weston SA, Bailey KR. Intermittent claudication: an objective office-based assessment. *J Am Coll Cardiol*. 2001; 37:1381–1385.
 64. Amirhamzeh MM, Chant HJ, Rees JL, Powel RJ, Campbell WB. A comparative study of treadmill tests and heel raising exercise for peripheral arterial disease. *Eur J Vasc Endovasc Surg*. 1997;13:301–305.
 65. Suominen V, Rantanen T, Venermo M, Saarinen J, Salenius J. Prevalence and risk factors of PAD among patients with elevated ABI. *Eur J Vasc Endovasc Surg*. 2008;35:709–714.
 66. Aboyans V, Ho E, Denenberg JO, Ho LA, Natarajan L, Criqui MH. The association between elevated ankle systolic pressures and peripheral occlusive arterial disease in diabetic and nondiabetic subjects. *J Vasc Surg*. 2008;48:1197–1203.
 67. Aboyans V, Criqui MH, Denenberg JO, Knoke JD, Ridker PM, Fronek A. Risk factors for progression of peripheral arterial disease in large and small vessels. *Circulation*. 2006;113:2623–2629.
 68. Nicoloff AD, Taylor LM Jr, Sexton GJ, Schuff RA, Edwards JM, Yeager RA, Landry GJ, Moneta GL, Porter JM; Homocysteine and Progression of Atherosclerosis Study Investigators. Relationship between site of initial symptoms and subsequent progression of disease in a prospective study of atherosclerosis progression in patients receiving long-term treatment for symptomatic peripheral arterial disease. *J Vasc Surg*. 2002;35:38–46.
 69. Cronenwett JL, Warner KG, Zelenock GB, Whitehouse WM Jr, Graham LM, Lindenauer M, Stanley JC. Intermittent claudication: current results of nonoperative management. *Arch Surg*. 1984;119:430–436.
 70. Amighi J, Sabeti S, Schlager O, Francesconi M, Ahmadi R, Minar E, Schillinger M. Outcome of conservative therapy of patients with severe intermittent claudication. *Eur J Vasc Endovasc Surg*. 2004;27:254–258.
 71. Norgren L, Hiatt WR, Dormandy JA, Nehler MR, Harris KA, Fowkes FG; TASC II Working Group. Inter-society consensus for the man-

- agement of peripheral arterial disease (TASC II). *J Vasc Surg.* 2007; 45(suppl S):S5–S67.
72. Marston WA, Davies SW, Armstrong B, Farber MA, Mendes RC, Fulton JJ, Keagy BA. Natural history of limbs with arterial insufficiency and chronic ulceration treated without revascularization. *J Vasc Surg.* 2006;44:108–114.
 73. Hamalainen H, Ronnema T, Halonen JP, Toikka T. Factors predicting lower extremity amputations in patients with type 1 or type 2 diabetes mellitus: a population-based 7-year follow-up study. *J Intern Med.* 1999;246:97–103.
 74. Brothers TE, Esteban R, Robison JG, Elliott BM. Symptoms of chronic arterial insufficiency correlate with absolute ankle pressure better than with ankle:brachial index. *Minerva Cardioangiol.* 2000;48:103–109.
 75. Matzke S, Ollgren J, Lepantalo M. Predictive value of distal pressure measurements in critical leg ischaemia. *Ann Chir Gynaecol.* 1996;85: 316–321.
 76. Fowl RJ, Gewirtz RJ, Love MC, Kempczinski RF. Natural history of claudicants with critical hemodynamic indices. *Ann Vasc Surg.* 1992;6: 31–33.
 77. Decrinis M, Doder S, Stark G, Pilger E. A prospective evaluation of sensitivity and specificity of the ankle/brachial index in the follow-up of superficial femoral artery occlusions treated by angioplasty. *Clin Investig.* 1994;72:592–597.
 78. Motukuru V, Suresh KR, Vivekanand V, Raj S, Giriya KR. Therapeutic angiogenesis in Buerger's disease (thromboangiitis obliterans) patients with critical limb ischemia by autologous transplantation of bone marrow mononuclear cells. *J Vasc Surg.* 2008;48(suppl):53S–60S.
 79. Allouche-Cometto L, Leger P, Rousseau H, Lefebvre D, Bendayan P, Elefterion P, Boccalon H. Comparative of blood flow to the ankle-brachial index after iliac angioplasty. *Int Angiol.* 1999;18:154–157.
 80. Matoba S, Tatsumi T, Murohara T, Imaizumi T, Katsuda Y, Ito M, Saito Y, Uemura S, Suzuki H, Fukumoto S, Yamamoto Y, Onodera R, Teramukai S, Fukushima M, Matsubara H; TACT Follow-Up Study Investigators. Long-term clinical outcome after intramuscular implantation of bone marrow mononuclear cells (Therapeutic Angiogenesis by Cell Transplantation [TACT] trial) in patients with chronic limb ischemia. *Am Heart J.* 2008;156:1010–1018.
 81. Barnes RW, Thompson BW, MacDonald CM, Nix ML, Lambeth A, Nix AD, Johnson DW, Wallace BH. Serial noninvasive studies do not herald postoperative failure of femoropopliteal or femorotibial bypass grafts. *Ann Surg.* 1989;210:486–493.
 82. Stierli P, Aeberhard P, Livers M. The role of colour flow duplex screening in infra-inguinal vein grafts. *Eur J Vasc Surg.* 1992;6: 293–298.
 83. Laborde AL, Synn AY, Worsey MJ, Bower TR, Hoballah JJ, Sharp WJ, Kresowik TF, Corson JD. A prospective comparison of ankle/brachial indices and color duplex imaging in surveillance of the in situ saphenous vein bypass. *J Cardiovasc Surg (Torino).* 1992;33:420–425.
 84. Idu MM, Blankenstein JD, de Gier P, Truyen E, Buth J. Impact of a color-flow duplex surveillance program on infrainguinal vein graft patency: a five-year experience. *J Vasc Surg.* 1993;17:42–52.
 85. Dalsing MC, Cikrit DF, Lalka SG, Sawchuk AP, Schulz C. Femorodistal vein grafts: the utility of graft surveillance criteria. *J Vasc Surg.* 1995; 21:127–134.
 86. Lundell A, Lindblad B, Bergqvist D, Hansen F. Femoropopliteal-crural graft patency is improved by an intensive surveillance program: a prospective randomized study. *J Vasc Surg.* 1995;21:26–33.
 87. Radak D, Labs KH, Jager KA, Bojic M, Popovic AD. Doppler-based diagnosis of restenosis after femoropopliteal percutaneous transluminal angioplasty: sensitivity and specificity of the ankle/brachial pressure index versus changes in absolute pressure values. *Angiology.* 1999;50: 111–122.
 88. McDermott MM, Greenland P, Liu K, Guralnik JM, Criqui MH, Dolan NC, Chan C, Celic L, Pearce WH, Schneider JR, Sharma L, Clark E, Gibson D, Martin GJ. Leg symptoms in peripheral arterial disease: associated clinical characteristics and functional impairment. *JAMA.* 2001;286:1599–1606.
 89. McDermott MM, Fried L, Simonsick E, Ling S, Guralnik JM. Asymptomatic peripheral arterial disease is independently associated with impaired lower extremity functioning: the Women's Health and Aging Study. *Circulation.* 2000;101:1007–1012.
 90. McDermott MM, Greenland P, Liu K, Guralnik JM, Celic L, Criqui MH, Chan C, Martin GJ, Schneider J, Pearce WH, Taylor LM, Clark E. The ankle brachial index is associated with leg function and physical activity: the Walking and Leg Circulation Study. *Ann Intern Med.* 2002;136: 873–883.
 91. McDermott MM, Ohlmler SM, Liu K, Guralnik JM, Martin GJ, Pearce WH, Greenland P. Gait alterations associated with walking impairment in people with peripheral arterial disease with and without intermittent claudication. *J Am Geriatr Soc.* 2001;49:747–754.
 92. McDermott MM, Liu K, Greenland P, Guralnik JM, Criqui MH, Chan C, Pearce WH, Schneider JR, Ferrucci L, Celic L, Taylor LM, Vonesh E, Martin GJ, Clark E. Functional decline in peripheral arterial disease: associations with the ankle brachial index and leg symptoms. *JAMA.* 2004;292:453–461.
 93. Szuba A, Oka RK, Harada R, Cooke JP. Limb hemodynamics are not predictive of functional capacity in patients with PAD. *Vasc Med.* 2006;11:155–163.
 94. Gardner AW, Skinner JS, Cantwell BW, Smith LK. Prediction of claudication pain from clinical measurements obtained at rest. *Med Sci Sports Exerc.* 1992;24:163–170.
 95. Parr B, Noakes TD, Derman EW. Factors predicting walking intolerance in patients with peripheral arterial disease and intermittent claudication. *S Afr Med J.* 2008;98:958–962.
 96. McDermott MM, Criqui MH, Liu K, Guralnik JM, Greenland P, Martin GJ, Pearce W. Lower ankle/brachial index, as calculated by averaging the dorsalis pedis and posterior tibial arterial pressures, and association with leg functioning in peripheral arterial disease. *J Vasc Surg.* 2000; 32:1164–1171.
 97. McDermott MM, Ferrucci L, Guralnik JM, Dyer AR, Liu K, Pearce WH, Clark E, Liao Y, Criqui MH. The ankle-brachial index is associated with the magnitude of impaired walking endurance among men and women with peripheral arterial disease. *Vasc Med.* 2010;15:251–257.
 98. McDermott MM, Liu K, Ferrucci L, Tian L, Guralnik JM, Green D, Tan J, Liao Y, Pearce WH, Schneider JR, McCue K, Ridker P, Rifai N, Criqui MH. Circulating blood markers and functional impairment in peripheral arterial disease. *J Am Geriatr Soc.* 2008;56:1504–1510.
 99. Herman SD, Liu K, Tian L, Guralnik JM, Ferrucci L, Criqui MH, Liao Y, McDermott MM. Baseline lower extremity strength and subsequent decline in functional performance at 6-year follow-up in persons with lower extremity peripheral arterial disease. *J Am Geriatr Soc.* 2009;57: 2246–2252.
 100. Anderson JD, Epstein FH, Meyer CH, Hagstiel KD, Wang H, Berr SS, Harthun NL, Weltman A, Dimaria JM, West AM, Kramer CM. Multifactorial determinants of functional capacity in peripheral arterial disease: uncoupling of calf muscle perfusion and metabolism. *J Am Coll Cardiol.* 2009;54:628–635.
 101. McDermott MM, Liu K, Ferrucci L, Tian L, Guralnik JM, Liao Y, Criqui MH. Greater sedentary hours and slower walking speed outside the home predict faster declines in functioning and adverse calf muscle changes in peripheral arterial disease. *J Am Coll Cardiol.* 2011;57: 2356–2364.
 102. McDermott MM, Liu K, Ferrucci L, Criqui MH, Greenland P, Guralnik JM, Tian L, Schneider JR, Pearce WH, Tan J, Martin GJ. Physical performance in peripheral arterial disease: a slower rate of decline in patients who walk more. *Ann Intern Med.* 2006;144:10–20.
 103. Selvin E, Erlinger TP. Prevalence of and risk factors for peripheral arterial disease in the United States: results from the National Health and Nutrition Examination Survey, 1999–2000. *Circulation.* 2004;110: 738–743.
 104. Newman AB, Siscovick DS, Manolio TA, Polak J, Fried LP, Borhani NO, Wolfson SK. Ankle-arm index as a marker of atherosclerosis in the Cardiovascular Health Study: Cardiovascular Health Study (CHS) Collaborative Research Group. *Circulation.* 1993;88:837–845.
 105. Allison MA, Criqui MH, McClelland RL, Scott JM, McDermott MM, Liu K, Folsom AR, Bertoni AG, Sharrett AR, Homma S, Kori S. The effect of novel cardiovascular risk factors on the ethnic-specific odds for peripheral arterial disease in the Multi-Ethnic Study of Atherosclerosis (MESA). *J Am Coll Cardiol.* 2006;48:1190–1197.
 106. Weatherley BD, Nelson JJ, Heiss G, Chambless LE, Sharrett AR, Nieto FJ, Folsom AR, Rosamond WD. The association of the ankle-brachial index with incident coronary heart disease: the Atherosclerosis Risk in Communities (ARIC) study, 1987–2001. *BMC Cardiovasc Disord.* 2007;3.
 107. Newman AB, Shemanski L, Manolio TA, Cushman M, Mittelmark M, Polak JF, Powe NR, Siscovick D. Ankle-arm index as a predictor of cardiovascular disease and mortality in the Cardiovascular Health Study: the Cardiovascular Health Study Group. *Arterioscler Thromb Vasc Biol.* 1999;19:538–545.

108. Hirsch AT, Criqui MH, Treat-Jacobson D, Regensteiner JG, Creager MA, Olin JW, Krook SH, Hunninghake DB, Comerota AJ, Walsh ME, McDermott MM, Hiatt WR. Peripheral arterial disease detection, awareness, and treatment in primary care. *JAMA*. 2001;286:1317–1324.
109. Murabito JM, Evans JC, Nieto K, Larson MG, Levy D, Wilson PW. Prevalence and clinical correlates of peripheral arterial disease in the Framingham Offspring Study. *Am Heart J*. 2002;143:961–965.
110. Zander E, Heinke P, Reindel J, Kohnert KD, Kairies U, Braun J, Eckel L, Kerner W. Peripheral arterial disease in diabetes mellitus type 1 and type 2: are there different risk factors? *Vasa*. 2002;31:249–254.
111. Hayashi C, Ogawa O, Kubo S, Mitsuhashi N, Onuma T, Kawamori R. Ankle brachial pressure index and carotid intima-media thickness as atherosclerosis markers in Japanese diabetics. *Diabetes Res Clin Pract*. 2004;66:269–275.
112. Yang X, Sun K, Zhang W, Wu H, Zhang H, Hui R. Prevalence of and risk factors for peripheral arterial disease in the patients with hypertension among Han Chinese. *J Vasc Surg*. 2007;46:296–302.
113. Ovbiagele B. Association of ankle-brachial index level with stroke. *J Neurol Sci*. 2009;276:14–17.
114. Ramos R, Quesada M, Solanas P, Subirana I, Sala J, Vila J, Masia R, Cerezo C, Elosua R, Grau M, Cordon F, Juvinya D, Fito M, Isabel Covas M, Clara A, Angel Munoz M, Marrugat J; REGICOR Investigators. Prevalence of symptomatic and asymptomatic peripheral arterial disease and the value of the ankle-brachial index to stratify cardiovascular risk. *Eur J Vasc Endovasc Surg*. 2009;38:305–311.
115. Allison MA, Hiatt WR, Hirsch AT, Coll JR, Criqui MH. A high ankle-brachial index is associated with increased cardiovascular disease morbidity and lower quality of life. *J Am Coll Cardiol*. 2008;51:1292–1298.
116. Criqui MH, McClelland RL, McDermott MM, Allison MA, Blumenthal RS, Aboyans V, Ix JH, Burke GL, Liu K, Shea S. The ankle-brachial index and incident cardiovascular events in the MESA (Multi-Ethnic Study of Atherosclerosis). *J Am Coll Cardiol*. 2010;56:1506–1512.
117. Sutton-Tyrrell K, Venkitchalam L, Kanaya AM, Boudreau R, Harris T, Thompson T, Mackey RH, Visser M, Vaidean GD, Newman AB. Relationship of ankle blood pressures to cardiovascular events in older adults. *Stroke*. 2008;39:863–869.
118. Wattanakit K, Folsom AR, Duprez DA, Weatherley BD, Hirsch AT. Clinical significance of a high ankle-brachial index: insights from the Atherosclerosis Risk in Communities (ARIC) Study. *Atherosclerosis*. 2007;190:459–464.
119. Resnick HE, Foster GL. Prevalence of elevated ankle-brachial index in the United States 1999 to 2002. *Am J Med*. 2005;118:676–679.
120. Greenland P, Smith SC Jr, Grundy SM. Improving coronary heart disease risk assessment in asymptomatic people: role of traditional risk factors and noninvasive cardiovascular tests. *Circulation*. 2001;104:1863–1867.
121. Brindle P, Beswick A, Fahey T, Ebrahim S. Accuracy and impact of risk assessment in the primary prevention of cardiovascular disease: a systematic review. *Heart*. 2006;92:1752–1759.
122. Tsimikas S, Willerson JT, Ridker PM. C-reactive protein and other emerging blood biomarkers to optimize risk stratification of vulnerable patients. *J Am Coll Cardiol*. 2006;47(suppl):C19–C31.
123. Greenland P, LaBree L, Azen SP, Doherty TM, Detrano RC. Coronary artery calcium score combined with Framingham score for risk prediction in asymptomatic individuals. *JAMA*. 2004;291:210–215.
124. Leng GC, Fowkes FG, Lee AJ, Dunbar J, Housley E, Ruckley CV. Use of ankle brachial pressure index to predict cardiovascular events and death: a cohort study. *BMJ*. 1996;313:1440–1444.
125. Hooi JD, Kester AD, Stoffers HE, Rinkens PE, Knottnerus JA, van Ree JW. Asymptomatic peripheral arterial occlusive disease predicted cardiovascular morbidity and mortality in a 7-year follow-up study. *J Clin Epidemiol*. 2004;57:294–300.
126. Ogren M, Hedblad B, Isacson SO, Janzon L, Jungquist G, Lindell SE. Non-invasively detected carotid stenosis and ischaemic heart disease in men with leg arteriosclerosis. *Lancet*. 1993;342:1138–1141.
127. van der Meer IM, Bots ML, Hofman A, del Sol AI, van der Kuip DA, Witteman JC. Predictive value of noninvasive measures of atherosclerosis for incident myocardial infarction: the Rotterdam Study. *Circulation*. 2004;109:1089–1094.
128. Kornitzer M, Dramaix M, Sobolski J, Degre S, De Backer G. Ankle/arm pressure index in asymptomatic middle-aged males: an independent predictor of ten-year coronary heart disease mortality. *Angiology*. 1995;46:211–219.
129. Abbott RD, Petrovitch H, Rodriguez BL, Yano K, Schatz IJ, Popper JS, Masaki KH, Ross GW, Curb JD. Ankle/brachial blood pressure in men >70 years of age and the risk of coronary heart disease. *Am J Cardiol*. 2000;86:280–284.
130. Resnick HE, Lindsay RS, McDermott MM, Devereux RB, Jones KL, Fabsitz RR, Howard BV. Relationship of high and low ankle brachial index to all-cause and cardiovascular disease mortality: the Strong Heart Study. *Circulation*. 2004;109:733–739.
131. Aboyans V, Lacroix P, Tran MH, Salamaigne C, Galinat S, Archambeaud F, Criqui MH, Laskar M. The prognosis of diabetic patients with high ankle-brachial index depends on the coexistence of occlusive peripheral artery disease. *J Vasc Surg*. 2011;53:984–991.
132. Aboyans V, Lacroix P, Postil A, Guilloux J, Rolle F, Cornu E, Laskar M. Subclinical peripheral arterial disease and incompressible ankle arteries are both long-term prognostic factors in patients undergoing coronary artery bypass grafting. *J Am Coll Cardiol*. 2005;46:815–820.
133. Agnelli G, Cimminiello C, Meneghetti G, Urbini S; Polyvascular Atherothrombosis Observational Survey (PATHOS) Investigators. Low ankle-brachial index predicts an adverse 1-year outcome after acute coronary and cerebrovascular events. *J Thromb Haemost*. 2006;4:2599–2606.
134. Purroy F, Coll B, Oro M, Seto E, Pinol-Ripoll G, Plana A, Quilez A, Sanahuja J, Brieve L, Vega L, Fernandez E. Predictive value of ankle brachial index in patients with acute ischaemic stroke. *Eur J Neurol*. 2010;17:602–606.
135. Alberts MJ, Bhatt DL, Mas JL, Ohman EM, Hirsch AT, Rother J, Salette G, Goto S, Smith SC Jr, Liao CS, Wilson PW, Steg PG; Reduction of Atherothrombosis for Continued Health Registry Investigators. Three-year follow-up and event rates in the international Reduction of Atherothrombosis for Continued Health Registry. *Eur Heart J*. 2009;30:2318–2326.
136. Criqui MH, Ninomiya JK, Wingard DL, Ji M, Fronck A. Progression of peripheral arterial disease predicts cardiovascular disease morbidity and mortality. *J Am Coll Cardiol*. 2008;52:1736–1742.
137. Sheikh MA, Bhatt DL, Li J, Lin S, Bartholomew JR. Usefulness of postexercise ankle-brachial index to predict all-cause mortality. *Am J Cardiol*. 2011;107:778–782.
138. Mohler ER 3rd, Treat-Jacobson D, Reilly MP, Cunningham KE, Miani M, Criqui MH, Hiatt WR, Hirsch AT. Utility and barriers to performance of the ankle-brachial index in primary care practice. *Vasc Med*. 2004;9:253–260.
139. Bendermacher BLW, Teijink JAW, Willigendael EM. Applicability of the ankle brachial index measurement as screening device in general practice for high cardiovascular risk. In: Bendermacher B. *Peripheral Arterial Disease. Screening, Diagnosis and Conservative Treatment* [dissertation]. Maastricht, Netherlands: Maastricht University; 2007.
140. Pollak EW, Chavis P, Wolfman EF. The effect of postural changes upon the ankle arterial perfusion pressure. *Vasc Surg*. 1976;10:219–222.
141. Gornik HL, Garcia B, Wolski K, Jones DC, Macdonald KA, Fronck A. Validation of a method for determination of the ankle-brachial index in the seated position. *J Vasc Surg*. 2008;48:1204–1210.
142. Yataco AR, Gardner AW. Acute reduction in ankle/brachial index following smoking in chronic smokers with peripheral arterial occlusive disease. *Angiology*. 1999;50:355–360.
143. Manning DM, Kuchirka C, Kaminski J. Miscuffing: inappropriate blood pressure cuff application. *Circulation*. 1983;68:763–766.
144. Pickering TG, Hall JE, Appel LJ, Falkner BE, Graves J, Hill MN, Jones DW, Kurtz T, Sheps SG, Roccella EJ. Recommendations for blood pressure measurement in humans and experimental animals, part 1: blood pressure measurement in humans: a statement for professionals from the Subcommittee of Professional and Public Education of the American Heart Association Council on High Blood Pressure Research. *Circulation*. 2005;111:697–716.
145. Mundt KA, Chambless LE, Burnham CB, Heiss G. Measuring ankle systolic blood pressure: validation of the Dinamap 1846 SX. *Angiology*. 1992;43:555–566.
146. Takahashi O, Shimbo T, Rahman M, Musa R, Kurokawa W, Yoshinaka T, Fukui T. Validation of the auscultatory method for diagnosing peripheral arterial disease. *Fam Pract*. 2006;23:10–14.
147. Aboyans V, Lacroix P, Doucet S, Preux PM, Criqui MH, Laskar M. Diagnosis of peripheral arterial disease in general practice: can the ankle-brachial index be measured either by pulse palpation or an automatic blood pressure device? *Int J Clin Pract*. 2008;62:1001–1007.
148. Adiseshiah M, Cross FW, Belsham PA. Ankle blood pressure measured by automatic oscillometry: a comparison with Doppler pressure measurements. *Ann R Coll Surg Engl*. 1987;69:271–273.

149. Beckman JA, Higgins CO, Gerhard-Herman M. Automated oscillometric determination of the ankle-brachial index provides accuracy necessary for office practice. *Hypertension*. 2006;47:35–38.
150. Benchimol A, Bernard V, Pillois X, Hong NT, Benchimol D, Bonnet J. Validation of a new method of detecting peripheral artery disease by determination of ankle-brachial index using an automatic blood pressure device. *Angiology*. 2004;55:127–134.
151. Blebea J, Ali MK, Love M, Bodenham R, Bacik B. Automatic postoperative monitoring of infrainguinal bypass procedures. *Arch Surg*. 1997;132:286–291.
152. Cortez-Cooper MY, Supak JA, Tanaka H. A new device for automatic measurements of arterial stiffness and ankle-brachial index. *Am J Cardiol*. 2003;91:1519–1522, A9.
153. Diehm N, Dick F, Czuprin C, Lawall H, Baumgartner I, Diehm C. Oscillometric measurement of ankle-brachial index in patients with suspected peripheral disease: comparison with Doppler method. *Swiss Med Wkly*. 2009;139:357–363.
154. Ena J, Lozano T, Verdú G, Argente CR, González VL. Accuracy of ankle-brachial index obtained by automated blood pressure measuring devices in patients with diabetes mellitus. *Diabetes Res Clin Pract*. 2011;92:329–336.
155. Jonsson B, Lindberg LG, Skau T, Thulesius O. Is oscillometric ankle pressure reliable in leg vascular disease? *Clin Physiol*. 2001;21:155–163.
156. Korno M, Eldrup N, Sillesen H. Comparison of ankle-brachial index measured by an automated oscillometric apparatus with that by standard Doppler technique in vascular patients. *Eur J Vasc Endovasc Surg*. 2009;38:610–615.
157. Lee BY, Campbell JS, Berkowitz P. The correlation of ankle oscillometric blood pressures and segmental pulse volumes to Doppler systolic pressures in arterial occlusive disease. *J Vasc Surg*. 1996;23:116–122.
158. MacDonald E, Froggatt P, Lawrence G, Blair S. Are automated blood pressure monitors accurate enough to calculate the ankle brachial pressure index? *J Clin Monit Comput*. 2008;22:381–384.
159. MacDougall AM, Tandon V, Wilson MP, Wilson TW. Oscillometric measurement of ankle-brachial index. *Can J Cardiol*. 2008;24:49–51.
160. Mehlsen J, Wiinberg N, Bruce C. Oscillometric blood pressure measurement: a simple method in screening for peripheral arterial disease. *Clin Physiol Funct Imaging*. 2008;28:426–429.
161. Nukumizu Y, Matsushita M, Sakurai T, Kobayashi M, Nishikimi N, Komori K. Comparison of Doppler and oscillometric ankle blood pressure measurement in patients with angiographically documented lower extremity arterial occlusive disease. *Angiology*. 2007;58:303–308.
162. Pan CR, Staessen JA, Li Y, Wang JG. Comparison of three measures of the ankle-brachial blood pressure index in a general population. *Hypertens Res*. 2007;30:555–561.
163. Raines JK, Farrar J, Noicey K, Pena J, Davis WW, Willens HJ, Wallace DD. Ankle-brachial index in the primary care setting. *Vasc Endovascular Surg*. 2004;38:131–136.
164. Ramanathan A, Conaghan PJ, Jenkinson AD, Bishop CR. Comparison of ankle-brachial pressure index measurements using an automated oscillometric device with the standard Doppler ultrasound technique. *ANZ J Surg*. 2003;73:105–108.
165. Richart T, Kuznetsova T, Wizner B, Struijker-Boudier HA, Staessen JA. Validation of automated oscillometric versus manual measurement of the ankle-brachial index. *Hypertens Res*. 2009;32:884–888.
166. Salles-Cunha SX, Vincent DG, Towne JB, Bernhard VM. Noninvasive ankle pressure measurements by oscillometry. *Tex Heart Inst J*. 1982;9:349–357.
167. Bonham PA, Cappuccio M, Hulsey T, Michel Y, Kelechi T, Jenkins C, Robison J. Are ankle and toe brachial indices (ABI-TBI) obtained by a pocket Doppler interchangeable with those obtained by standard laboratory equipment? *J Wound Ostomy Continence Nurs*. 2007;34:35–44.
168. Carmo GA, Mandil A, Nascimento BR, Arantes BD, Bittencourt JC, Falqueto EB, Ribeiro AL. Can we measure the ankle-brachial index using only a stethoscope? A pilot study. *Fam Pract*. 2009;26:22–26.
169. Khandanpour N, Armon MP, Jennings B, Clark A, Meyer FJ. Photoplethysmography, an easy and accurate method for measuring ankle brachial pressure index: can photoplethysmography replace Doppler? *Vasc Endovascular Surg*. 2009;43:578–582.
170. Ludyga T, Kuczmik WB, Kazibudski M, Nowakowski P, Orawczyk T, Glanowski M, Kucharzewski M, Ziaja D, Szaniewski K, Ziaja K. Ankle-brachial pressure index estimated by laser Doppler in patients suffering from peripheral arterial obstructive disease. *Ann Vasc Surg*. 2007;21:452–457.
171. Miglicci R, Nasorri R, Ricciarini P, Gresele P. Ankle-brachial index measured by palpation for the diagnosis of peripheral arterial disease. *Fam Pract*. 2008;25:228–232.
172. Nicolai SP, Kruidenier LM, Rouwet EV, Wetzels-Gulpers L, Rozeman CA, Prins MH, Teijink JA. Pocket Doppler and vascular laboratory equipment yield comparable results for ankle brachial index measurement. *BMC Cardiovasc Disord*. 2008;26.
173. Sadiq S, Chithriki M. Arterial pressure measurements using infrared photosensors: comparison with CW Doppler. *Clin Physiol*. 2001;21:129–132.
174. Whiteley MS, Fox AD, Horrocks M. Photoplethysmography can replace hand-held Doppler in the measurement of ankle/brachial indices. *Ann R Coll Surg Engl*. 1998;80:96–98.
175. Fowkes FG, Housley E, Macintyre CC, Prescott RJ, Ruckley CV. Variability of ankle and brachial systolic pressures in the measurement of atherosclerotic peripheral arterial disease. *J Epidemiol Community Health*. 1988;42:128–133.
176. Stoffers J, Kaiser V, Kester A, Schouten H, Knottnerus A. Peripheral arterial occlusive disease in general practice: the reproducibility of the ankle-arm systolic pressure ratio. *Scand J Prim Health Care*. 1991;9:109–114.
177. Kaiser V, Kester AD, Stoffers HE, Kitslaar PJ, Knottnerus JA. The influence of experience on the reproducibility of the ankle-brachial systolic pressure ratio in peripheral arterial occlusive disease. *Eur J Vasc Endovasc Surg*. 1999;18:25–29.
178. Aboyans V, Lacroix P, Lebourdon A, Preux PM, Ferrieres J, Laskar M. The intra- and interobserver variability of ankle-arm blood pressure index according to its mode of calculation. *J Clin Epidemiol*. 2003;56:215–220.
179. Baker JD, Dix DE. Variability of Doppler ankle pressures with arterial occlusive disease: an evaluation of ankle index and brachial-ankle pressure gradient. *Surgery*. 1981;89:134–137.
180. Johnston KW, Hosang MY, Andrews DF. Reproducibility of noninvasive vascular laboratory measurements of the peripheral circulation. *J Vasc Surg*. 1987;6:147–151.
181. de Graaff JC, Ubbink DT, Legemate DA, de Haan RJ, Jacobs MJ. Inter-observer and intraobserver reproducibility of peripheral blood and oxygen pressure measurements in the assessment of lower extremity arterial disease. *J Vasc Surg*. 2001;33:1033–1040.
182. Holland-Letz T, Endres HG, Biedermann S, Mahn M, Kunert J, Groh S, Pittrow D, von Bilderling P, Sternitzky R, Diehm C. Reproducibility and reliability of the ankle-brachial index as assessed by vascular experts, family physicians and nurses. *Vasc Med*. 2007;12:105–112.
183. Espeland MA, Regensteiner JG, Jaramillo SA, Gregg E, Knowler WC, Wagenknecht LE, Bahnson J, Haffner S, Hill J, Hiatt WR; Look AHEAD Study Group. Measurement characteristics of the ankle-brachial index: results from the Action for Health in Diabetes study. *Vasc Med*. 2008;13:225–233.
184. Weatherley BD, Chambless LE, Heiss G, Catellier DJ, Ellison CR. The reliability of the ankle-brachial index in the Atherosclerosis Risk in Communities (ARIC) study and the NHLBI Family Heart Study (FHS). *BMC Cardiovasc Disord*. 2006;7.
185. Forster FK, Turney D. Oscillometric determination of diastolic, mean and systolic blood pressure: a numerical model. *J Biomech Eng*. 1986;108:359–364.
186. Hamel JF, Foucaud D, Fanello S. Comparison of the automated oscillometric method with the gold standard Doppler ultrasound method to access the ankle-brachial pressure index. *Angiology*. 2010;61:487–491.
187. Ursino M, Cristalli C. A mathematical study of some biomechanical factors affecting the oscillometric blood pressure measurement. *IEEE Trans Biomed Eng*. 1996;43:761–778.
188. van Montfrans GA. Oscillometric blood pressure measurement: progress and problems. *Blood Press Monit*. 2001;6:287–290.
189. Benchimol D, Pillois X, Benchimol A, Houitte A, Sagardiluz P, Torteliet L, Bonnet J. Accuracy of ankle-brachial index using an automatic blood pressure device to detect peripheral artery disease in preventive medicine. *Arch Cardiovasc Dis*. 2009;102:519–524.
190. Kawamura T. Assessing ankle-brachial index (ABI) by using automated oscillometric devices. *Arq Bras Cardiol*. 2008;90:294–298.
191. Arveschoug AK, Revsbech P, Brochner-Mortensen J. Sources of variation in the determination of distal blood pressure measured using the strain gauge technique. *Clin Physiol*. 1998;18:361–368.
192. Brown J, Asongwed E, Chesbro S, John E. Inter-rater and intra-rater reliability of ankle brachial index (ABI) measurements using a stethoscope and Doppler. Paper presented at: American Physical

- Therapy Association Meeting; 2008. http://apps.Apta.Org/custom/abstracts/pt2008/abstractspt.Cfm?M_id_17314. Accessed February 10, 2011.
193. Clyne CA, Jones T, Moss S, Ensell J. The use of radioactive oxygen to study muscle function in peripheral vascular disease. *Surg Gynecol Obstet*. 1979;149:225–228.
 194. Osmundson PJ, O'Fallon WM, Clements IP, Kazmier FJ, Zimmerman BR, Palumbo PJ. Reproducibility of noninvasive tests of peripheral occlusive arterial disease. *J Vasc Surg*. 1985;2:678–683.
 195. Fisher CM, Burnett A, Makeham V, Kidd J, Glasson M, Harris JP. Variation in measurement of ankle-brachial pressure index in routine clinical practice. *J Vasc Surg*. 1996;24:871–875.
 196. Jeelani NU, Braithwaite BD, Tomlin C, MacSweeney ST. Variation of method for measurement of brachial artery pressure significantly affects ankle-brachial pressure index values. *Eur J Vasc Endovasc Surg*. 2000;20:25–28.
 197. Atsma F, Bartelink ML, Grobbee DE, van der Schouw YT. Best reproducibility of the ankle-arm index was calculated using Doppler and dividing highest ankle pressure by highest arm pressure. *J Clin Epidemiol*. 2005;58:1282–1288.
 198. van Langen H, van Gorp J, Rubbens L. Interobserver variability of ankle-brachial index measurements at rest and post exercise in patients with intermittent claudication. *Vasc Med*. 2009;14:221–226.
 199. Ray SA, Srodon PD, Taylor RS, Dormandy JA. Reliability of ankle: brachial pressure index measurement by junior doctors. *Br J Surg*. 1994;81:188–190.
 200. Endres HG, Huckle C, Holland-Letz T, Trampisch HJ. A new efficient trial design for assessing reliability of ankle-brachial index measures by three different observer groups. *BMC Cardiovasc Disord*. 2006;33.
 201. Osborn LA, Vernon SM, Reynolds B, Timm TC, Allen K. Screening for subclavian artery stenosis in patients who are candidates for coronary bypass surgery. *Catheter Cardiovasc Interv*. 2002;56:162–165.
 202. Vierron E, Halimi JM, Tichet J, Balkau B, Cogneau J, Giraudeau B; DESIR Study Group. Center effect on ankle-brachial index measurement when using the reference method (Doppler and manometer): results from a large cohort study. *Am J Hypertens*. 2009;22:718–722.
 203. Aboyans V, Criqui MH, McDermott MM, Allison MA, Denenberg JO, Shadman R, Fronek A. The vital prognosis of subclavian stenosis. *J Am Coll Cardiol*. 2007;49:1540–1545.
 204. Shadman R, Criqui MH, Bundens WP, Fronek A, Denenberg JO, Gamst AC, McDermott MM. Subclavian artery stenosis: prevalence, risk factors, and association with cardiovascular diseases. *J Am Coll Cardiol*. 2004;44:618–623.
 205. Clark CE, Campbell JL, Powell RJ. The interarm blood pressure difference as predictor of cardiovascular events in patients with hypertension in primary care: cohort study. *J Hum Hypertens*. 2007;21:633–638.
 206. Orme S, Ralph SG, Birchall A, Lawson-Matthew P, McLean K, Channer KS. The normal range for inter-arm differences in blood pressure. *Age Ageing*. 1999;28:537–542.
 207. Aboyans V, Kamineni A, Allison MA, McDermott MM, Crouse JR, Ni H, Szklo M, Criqui MH. The epidemiology of subclavian stenosis and its association with markers of subclinical atherosclerosis: the Multi-Ethnic Study of Atherosclerosis (MESA). *Atherosclerosis*. 2010;211:266–270.
 208. Espinola-Klein C, Rupprecht HJ, Bickel C, Lackner K, Savvidis S, Messow CM, Munzel T, Blankenberg S; AtheroGene Investigators. Different calculations of ankle-brachial index and their impact on cardiovascular risk prediction. *Circulation*. 2008;118:961–967.
 209. O'Hare AM, Katz R, Shlipak MG, Cushman M, Newman AB. Mortality and cardiovascular risk across the ankle-arm index spectrum: results from the Cardiovascular Health Study. *Circulation*. 2006;113:388–393.
 210. Fowkes FG, Price JF, Stewart MC, Butcher I, Leng GC, Pell AC, Sandercock PA, Fox KA, Lowe GD, Murray GD; Aspirin for Asymptomatic Atherosclerosis Trialists. Aspirin for prevention of cardiovascular events in a general population screened for a low ankle brachial index: a randomized controlled trial. *JAMA*. 2010;303:841–848.
 211. Hoogeveen EK, Kostense PJ, Beks PJ, Mackaay AJ, Jakobs C, Bouter LM, Heine RJ, Stehouwer CD. Hyperhomocysteinemia is associated with an increased risk of cardiovascular disease, especially in non-insulin-dependent diabetes mellitus: a population-based study. *Arterioscler Thromb Vasc Biol*. 1998;18:133–138.
 212. Fowler B, Jamrozik K, Norman P, Allen Y. Prevalence of peripheral arterial disease: persistence of excess risk in former smokers. *Aust N Z J Public Health*. 2002;26:219–224.
 213. Jager A, Kostense PJ, Ruhe HG, Heine RJ, Nijpels G, Dekker JM, Bouter LM, Stehouwer CD. Microalbuminuria and peripheral arterial disease are independent predictors of cardiovascular and all-cause mortality, especially among hypertensive subjects: five-year follow-up of the Hoorn Study. *Arterioscler Thromb Vasc Biol*. 1999;19:617–624.
 214. McDermott MM, Guralnik JM, Albay M, Bandinelli S, Miniati B, Ferrucci L. Impairments of muscles and nerves associated with peripheral arterial disease and their relationship with lower extremity functioning: the INCHIANTI Study. *J Am Geriatr Soc*. 2004;52:405–410.
 215. Rooke TW, Hirsch AT, Misra S, Sidawy AN, Beckman JA, Findeiss LK, Golzarian J, Gornik HL, Halperin JL, Jaff MR, Moneta GL, Olin JW, Stanley JC, White CJ, White JV, Zierler RE. ACCF/AHA focused update of the guideline for the management of patients with peripheral artery disease (updating the 2005 guideline): a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *Circulation*. 2011;124:2020–2045.
 216. Diehm C, Allenberg JR, Pittrow D, Mahn M, Tepohl G, Haberl RL, Darius H, Burghaus I, Trampisch HJ; German Epidemiological Trial on Ankle Brachial Index Study Group. Mortality and vascular morbidity in older adults with asymptomatic versus symptomatic peripheral artery disease. *Circulation*. 2009;120:2053–61.
 217. American Diabetes Association. Peripheral arterial disease in people with diabetes. *Diabetes Care*. 2003;26:3333–3341.
 218. Kennedy M, Solomon C, Manolio TA, Criqui MH, Newman AB, Polak JF, Burke GL, Enright P, Cushman M. Risk factors for declining ankle-brachial index in men and women 65 years or older: the Cardiovascular Health Study. *Arch Intern Med*. 2005;165:1896–1902.

KEY WORDS: AHA Scientific Statements ■ ankle brachial index