


Calculus: Limits and Asymptotes

Notes, examples, & practice quiz (with solutions)



Topics include definitions, greatest integer function, strategies, infinity, slant asymptote, squeeze theorem, and more.

Definition of a Limit

If $f(x)$ gets arbitrarily close to a single number L as x approaches c , then we write

$$\lim_{x \rightarrow c} f(x) = L$$

and say that "the limit of $f(x)$, as x approaches c , is L ."

Also, in order for the limit to exist, the values of f must tend to the same number L from the left or the right.

$$\lim_{x \rightarrow c^-} f(x) = L$$

("left-hand limit of $f(x)$ " or "limit from the left")


$$\lim_{x \rightarrow c^+} f(x) = L$$

("right-hand limit of $f(x)$ " or "limit from the right")

Note -- from the definition:

- 1) The limit is *unique* if it exists.
(limit from the left = limit from the right)
- 2) The limit does not depend on the actual value of $f(x)$ at c . Instead, it is determined by values of $f(x)$ when x is near c


Illustrations and Examples:



$$f(x) = \frac{x^3 - 1}{x - 1}$$

$$f(1) = \frac{0}{0} \text{ undefined}$$


$$\lim_{x \rightarrow 1} f(x) = 3$$



$$g(x) = \begin{cases} \frac{x^3 - 1}{x - 1} & \text{when } x \neq 1 \\ 0 & \text{when } x = 1 \end{cases}$$

$$g(1) = 0$$

$$\lim_{x \rightarrow 1} g(x) = 3$$




$$h(x) = x^2 + x + 1$$

$$h(1) = 1 + 1 + 1 = 3$$

$$\lim_{x \rightarrow 1} h(x) = 3$$

Note: The values at 1 are all different, but the limits are all the same, because the values of the functions as x gets near 1, approach 3.

$g(x) = [x]$
"greatest integer function"




$$g(1.5) = 1$$

$$g(-2.3) = -3$$

$$g(2) = 2$$

$$g(1.00034) = 1$$

approaching 2 from the right...



approaching 2 from the left...

$$\lim_{x \rightarrow 1.5} [x] = 1$$

$$\lim_{x \rightarrow -2.3} [x] = -3$$

$$\lim_{x \rightarrow 2} [x] \neq 2$$

why is it not equal to 2?

Note: For the 'greatest integer function', the limit exists for all values of x that are NOT integers!

$$\lim_{x \rightarrow 2^+} [x] = 2$$

If you look at the graph, you can see when approaching 2 FROM THE RIGHT, the values are 2...

$$\lim_{x \rightarrow 2^-} [x] = 1$$

But, when approaching 2 FROM THE LEFT, the values are 1..

Since the limits are different, the limit does not exist at 2!!

Finding Limits: Examples

1) "Plug in the Number" (Direct Substitution)

$$\begin{aligned} \lim_{x \rightarrow 3} x^2 + 3x - 1 &= \lim_{x \rightarrow 3} x^2 + \lim_{x \rightarrow 3} 3x - \lim_{x \rightarrow 3} 1 \\ &= 9 + 9 - 1 = 17 \end{aligned}$$


$$\begin{aligned} \lim_{x \rightarrow 0} 1 + x \cos(2x) &= \lim_{x \rightarrow 0} 1 + \lim_{x \rightarrow 0} x \cdot \lim_{x \rightarrow 0} \cos(2x) \\ &= 1 + 0 \cdot 1 = 1 \end{aligned}$$

$$\lim_{x \rightarrow 8} \frac{x}{x-8} = \frac{8}{0} \text{ undefined (or, does not exist)}$$

2) "Eliminate the Problem"

$$\begin{aligned} \lim_{x \rightarrow 2} \frac{(x^2-4)}{(x-2)} &= \frac{0}{0} \quad \text{However, we can factor the numerator..} \\ &\quad \text{then, cancel the denominator...} \\ &\quad \text{then, solve...} \end{aligned}$$

$$\begin{aligned} \lim_{x \rightarrow 2} \frac{(x+2)(x-2)}{(x-2)} \\ \lim_{x \rightarrow 2} \frac{(x+2)(\cancel{x-2})}{(\cancel{x-2})} \\ \lim_{x \rightarrow 2} x+2 = \boxed{4} \end{aligned}$$




$$\begin{aligned} \lim_{x \rightarrow 2} \frac{x^3-8}{x^2-4} \quad \text{Again, at } x=2, \text{ there is a 0 in the denominator.} \\ \text{So, we factor (with difference of squares and difference of cubes)} \\ \text{Simplify and plug in 2...} \end{aligned}$$

$$\begin{aligned} \lim_{x \rightarrow 2} \frac{(x-2)(x^2+2x+4)}{(x+2)(x-2)} \\ \lim_{x \rightarrow 2} \frac{(\cancel{x-2})(x^2+2x+4)}{(x+2)(\cancel{x-2})} \\ \frac{(2)^2+2(2)+4}{(2)+2} = \frac{12}{4} = \boxed{3} \end{aligned}$$

3) "Extrapolate the Limit"

$$\lim_{x \rightarrow \infty} 1 + \frac{2}{x^3} = 1$$

x	1/2	1	2	4	10	approaches ∞
f(x)	17	3	1.25	1.031	1.002	approaches 1



$$\lim_{x \rightarrow \infty} \frac{2^{-x}}{2^x} = 0$$

As X increases, the numerator is getting smaller and the denominator is getting larger..
Therefore, the function is decreasing toward 0.

"End Behavior of Polynomial"

For polynomials whose (largest) degree ≥ 1

if leading coefficient is positive, then it becomes infinite as x does..

if leading coefficient is negative, then it becomes negatively infinite as x increases to ∞

$$\lim_{x \rightarrow \infty} x^2 - 23x + 2 = \infty$$

$$\lim_{x \rightarrow \infty} 8x - 5x^2 = -\infty$$

$$\lim_{x \rightarrow \infty} .00004x - 10^5 = \infty$$

4) "Utilizing the Conjugate"

$$\lim_{h \rightarrow 0} \frac{\sqrt{1+h} - 1}{h} \quad \text{"plug in the number"} \quad \frac{\sqrt{1+0} - 1}{0} = \frac{0}{0} \quad (\text{cannot determine yet})$$

$$\lim_{h \rightarrow 0} \frac{(\sqrt{1+h} - 1)(\sqrt{1+h} + 1)}{h(\sqrt{1+h} + 1)} = \frac{\sqrt{1+h}^2 - 1^2}{h(\sqrt{1+h} + 1)}$$

**Important observation: we multiplied the numerator terms, but did not combine the denominator terms

$$= \frac{\cancel{h}}{h(\sqrt{1+h} + 1)} = \lim_{h \rightarrow 0} \frac{1}{\sqrt{1+h} + 1} \quad \text{Try again...} \quad \boxed{\frac{1}{2}}$$

5) "Expand or Rewrite"

$$\lim_{h \rightarrow 0} \frac{(h+2)^3 - 8}{h} \quad \text{"substitute the 0"} \quad \frac{(0+2)^3 - 8}{0} = \frac{0}{0} \quad (\text{cannot determine yet})$$

Expand the numerator:

$$(h+2)(h+2) = h^2 + 4h + 4$$

$$(h^2 + 4h + 4)(h+2) = \frac{h^3 + 4h^2 + 4h}{2h^2 + 8h + 8}$$

$$h^3 + 6h^2 + 12h + 8$$

$$\lim_{h \rightarrow 0} \frac{h^3 + 6h^2 + 12h + 8 - 8}{h} =$$

$$\lim_{h \rightarrow 0} h^2 + 6h + 12 = \boxed{12}$$

$$\lim_{x \rightarrow -4} \frac{\frac{1}{4} + \frac{1}{x}}{4+x} = \lim_{x \rightarrow -4} \frac{\frac{x}{4x} + \frac{4}{4x}}{4+x} = \lim_{x \rightarrow -4} \frac{\frac{x+4}{4x}}{x+4} = \lim_{x \rightarrow -4} \frac{1}{4x} = \boxed{\frac{-1}{16}}$$

Example: Using 2 methods, find $\lim_{x \rightarrow 9} \frac{\sqrt{x} - 3}{x - 9}$

method 1: multiply by the conjugate

method 2: extrapolate / use a chart

$$\lim_{x \rightarrow 9} \frac{\sqrt{x} - 3}{x - 9} \cdot \frac{(\sqrt{x} + 3)}{(\sqrt{x} + 3)} =$$

$$\lim_{x \rightarrow 9} \frac{x - 9}{x - 9} \cdot \frac{1}{(\sqrt{x} + 3)} =$$

$$\lim_{x \rightarrow 9} \frac{1}{(\sqrt{x} + 3)} = \boxed{\frac{1}{6}}$$

x	8	8.5	8.9	9	9.1	9.5	10
f(x)	.171	.169	.167		.166	.164	.162



Example: $\lim_{x \rightarrow 0} \frac{1 - \cos x}{\sin x}$ substitute $x = 0$, $\frac{1 - \cos(0)}{\sin(0)} = \frac{0}{0}$ indeterminate...

(Use conjugate) $\frac{1 - \cos x}{\sin x} \cdot \frac{(1 + \cos x)}{(1 + \cos x)} =$

(Trig Identity) $\frac{1 - \cos^2 x}{\sin x(1 + \cos x)} =$

$$\frac{\sin^2 x}{\sin x(1 + \cos x)} = \frac{\sin x}{(1 + \cos x)}$$

substitute $x = 0$ $\frac{\sin(0)}{(1 + \cos(0))} = \frac{0}{2} = 0$

Example: $\lim_{x \rightarrow 0} \frac{\frac{x-1}{5x-3} - \frac{1}{3}}{x}$ substitute $x = 0$, and the result is $\frac{0}{0}$ indeterminate...

(Combine numerator terms and simplify) $\frac{\frac{3(x-1)}{3(5x-3)} - \frac{(5x-3)(1)}{(5x-3)(3)}}{x} = \frac{\frac{3x-3 - (5x-3)}{(5x-3)(3)}}{x} = \frac{-2x}{15x-9}$

$$= \frac{-2}{15x-9} = \frac{2}{9}$$

Example: $\lim_{x \rightarrow 2^-} \frac{x^2(x-2)(x+3)}{|x-2|}$ limit as x approaches 2 from the left...
at $x = 2$, the equation is $0/0$

$$\frac{x^2 \cancel{(x-2)}(x+3)}{|x-2|}$$

put in $x = 2$, and the solution is 20...

-20 from the left!

(20 from the right)..

x	1	1.5	1.9	1.95	2.05	2.1	2.5	3
f(x)	-4	-10.1	-17.7	-18.8	21.2	22.5	34.4	54

Example: $\lim_{x \rightarrow 4} \frac{\frac{3}{x+2} - \frac{1}{x-2}}{x-4}$ $\frac{\frac{3(x-2) - (x+2)}{(x-2)(x+2)}}{x-4} = \frac{3x-8-x}{(x-2)(x+2)}$

$$\frac{2(x-4)}{(x-2)(x+2)} \cdot \frac{1}{x-4} = \frac{2}{(x-2)(x+2)}$$

$$\lim_{x \rightarrow 4} \frac{2}{(x-2)(x+2)} = \frac{1}{6}$$

Using a graph to determine the limit: The absolute value example

$$h(x) = \frac{x - 3}{|x - 3|}$$

Find $\lim_{x \rightarrow 3} h(x)$

Step 1: Use direct substitution


$$\lim_{x \rightarrow 3} \frac{x - 3}{|x - 3|} = \frac{(3) - 3}{|(3) - 3|} = \frac{0}{0} \quad \text{Inconclusive!}$$

Step 2: Try to factor or use conjugates

These techniques will not work in this problem...

Step 3: Make tables of values and graph

approaching from the left		approaching from the right	
x	h(x)	x	h(x)
2	-1	4	1
2.5	-1	3.5	1
2.8	-1	3.2	1
2.9	-1	3.1	1
2.99	-1	3.01	1



Conclusion:

$$\lim_{x \rightarrow 3^-} h(x) = -1$$

$$\lim_{x \rightarrow 3^+} h(x) = 1$$

$$\lim_{x \rightarrow 3} h(x) = \text{Does Not Exist (DNE)}$$

(limit from the left and limit from the right differ)

"Squeeze" Theorem

(Also, called "pinching theorem" or "sandwich theorem")

Brief Summary: If $f(x) \leq g(x) \leq h(x)$ and $\lim_{x \rightarrow a} f(x) = \lim_{x \rightarrow a} h(x) = L$, then $\lim_{x \rightarrow a} g(x) = L$

Example: $\lim_{x \rightarrow 0} x^2 \cos\left(\frac{1}{x}\right)$

Try direct substitution: $(0)^2 \cos\left(\frac{1}{0}\right)$????

But, we know $-1 \leq \cos(x) \leq 1$ (range of a cosine function)

So, $-1 \leq \cos\left(\frac{1}{x}\right) \leq 1$

multiply each term by x^2


$$-x^2 \leq x^2 \cos\left(\frac{1}{x}\right) \leq x^2$$

then, find the limit of the lower and upper terms.....

$$\lim_{x \rightarrow 0} -x^2 = 0$$

$$\lim_{x \rightarrow 0} x^2 = 0$$

$$\lim_{x \rightarrow 0} x^2 \cos\left(\frac{1}{x}\right)$$



Finding the "limit as X approaches infinity"

General Rule: ("The Rational function Theorem")

Determining the limits at ∞ for functions expressed as a ratio of two polynomials:

$$f(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_0$$

$$g(x) = b_m x^m + b_{m-1} x^{m-1} + \dots + b_0$$

$$\lim_{x \rightarrow \pm \infty} \frac{f(x)}{g(x)} = \begin{cases} \pm \infty & \text{if } n > m & \text{("top heavy")} \\ 0 & \text{if } n < m & \text{("bottom heavy")} \\ \frac{a_n}{b_m} & \text{if } n = m & \text{("lead exponents equal"; look at the lead coefficients)} \end{cases}$$

Examples:

$$\lim_{x \rightarrow \infty} \frac{x}{x^2 - 1} = \frac{x^1}{x^2 - 1} = 0 \quad \text{("bottom heavy")}$$

degree of numerator < degree of denominator
degree of denominator > degree of numerator

(1)
(2)

$$\lim_{x \rightarrow \infty} \frac{5x^3 + 2}{(2x + 3)^3} = \lim_{x \rightarrow \infty} \frac{5x^3 + 2}{8x^3 + 36x^2 + 54x + 27} = \frac{5}{8}$$

highest degree of numerator = highest degree of denominator

(3)
(3)

$$\lim_{x \rightarrow \infty} \frac{7 + 3x^2 - 5x^4}{10x^3 - 17} = \lim_{x \rightarrow \infty} \frac{-5x^4 + 3x^2 + 7}{10x^3 - 17} = -\infty$$

rewrite polynomial in descending order
degree of numerator > degree of denominator

(4)
(3)

("top heavy")

(*Note: a simple graph showing end behavior confirms the limit is negative infinity, rather than positive infinity)

Comparison/Verification:

$$\lim_{x \rightarrow \infty} \frac{4 - x^2}{4x^2 - x - 2} = \lim_{x \rightarrow \infty} \frac{-x^2 + 4}{4x^2 - x - 2} = \frac{-1}{4}$$

(use "rational function theorem" above)


$$\lim_{x \rightarrow \infty} \frac{4 - x^2}{4x^2 - x - 2} = \lim_{x \rightarrow \infty} \frac{4/x^2 - 1}{4 - 1/x - 2/x^2} = \frac{0 - 1}{4 - 0 - 0} = \frac{-1}{4}$$

(rewrite and solve)
(multiply throughout by $\frac{1}{x^2}$)

Steps to find a limit (that approaches infinity)

- 1) (If necessary), expand the equation to reveal the degrees of the polynomials.
- 2) Arrange polynomials with highest degree first.
- 3a) If the numerator has a higher degree, then the limit is ∞
- b) If the denominator has a higher degree, then the limit is 0
- c) If the highest degree of the numerator polynomial is the same as the highest degree of the denominator polynomial, then the limit is

$$\frac{\text{lead coefficient of numerator}}{\text{lead coefficient of denominator}}$$



Asymptotes: Definitions and Example

The line $y = b$ is a *Horizontal Asymptote* of the graph of $y = f(x)$ if

$$\lim_{x \rightarrow \infty} f(x) = b \quad \text{or} \quad \lim_{x \rightarrow -\infty} f(x) = b$$

The line $x = a$ is a *Vertical Asymptote* of the graph of $y = f(x)$ if one or more of the following occur:


$$\lim_{x \rightarrow a^-} f(x) = +\infty \quad \text{or} \quad \lim_{x \rightarrow a^+} f(x) = -\infty$$

or

$$\lim_{x \rightarrow a^-} f(x) = -\infty \quad \text{or} \quad \lim_{x \rightarrow a^+} f(x) = +\infty$$

$$f(x) = \frac{2x+3}{x-1}$$

x	f(x)
-50	1.90
-20	1.76
-5	1.17
-2	0.33
-1	-5
-1/2	-1.3
0	-3
1/2	-8
1	undefined
2	7
5	3.25
10	2.55
20	2.26
50	2.10



horizontal asymptote:

$$\lim_{x \rightarrow \infty} \frac{2x+3}{x-1} = 2$$

also,

$$\lim_{x \rightarrow -\infty} \frac{2x+3}{x-1} = 2$$

vertical asymptote:

$$\lim_{x \rightarrow 1^+} \frac{2x+3}{x-1} = +\infty$$

$$\lim_{x \rightarrow 1^-} \frac{2x+3}{x-1} = -\infty$$


Graphs and the *Slant Asymptote*

Asymptotes are very useful when graphing functions.

Examples:

$$y = \frac{x}{x-4}$$

Vertical Asymptote: y is undefined at $x = 4$
 Horizontal Asymptote: degree of numerator: 1 (Since the degrees are equal, look at the coefficients of the lead terms)
 degree of denominator: 1
 therefore $y = 1$



Since (0, 0) is below the horizontal asymptote and to the left of the vertical asymptote, sketch the corresponding end behavior.

Then, select a point on the other side of the vertical asymptote. Examples: (5, 5) or (10, 5/3)

Since (5, 5) is above the horizontal asymptote and to the right of the vertical asymptote, sketch the corresponding end behavior.

(note: to check solutions, plug in random values)

$$y = \frac{x^2 + 1}{x^2 - 1}$$

Vertical Asymptote: $\frac{x^2 + 1}{x^2 - 1} = \frac{x^2 + 1}{(x+1)(x-1)}$


y is undefined at $x = -1$ and $x = 1$

Horizontal Asymptote: $\lim_{x \rightarrow \infty} \frac{x^2 + 1}{x^2 - 1} = 1$

therefore $y = 1$

y-intercept: (0, -1)

x-intercept: none because there is no x value that satisfies $\frac{x^2 + 1}{x^2 - 1} = 0$



x	-1.1	-1.8	-2	-3	-5	-10
y	221/21	53/28	5/3	5/4	13/12	101/99

x	2	3	5	10
y	5/3	5/4	13/12	101/99

Slant Asymptote: If the highest degree of the numerator is one more than the highest degree of the denominator, then there is a slant asymptote
 That asymptote is the quotient without the remainder.

Example:

$$f(x) = \frac{x^2 - 2x - 3}{x + 4}$$

$$= \frac{(x-3)(x+1)}{x+4}$$

Vertical Asymptote: $x = -4$

y-intercept: (0, -3/4)
 x-intercepts: (3, 0) (-1, 0)

Horizontal Asymptote: None


degree of numerator: 2
 degree of denominator: 1

slant asymptote

The slant asymptote is $y = x - 6$

As x gets larger and larger, the remainder gets closer and closer to zero. And, the function resembles $x - 6$ more and more.

$$x + 4 \overline{) \begin{array}{r} x - 6 + \frac{21}{x + 4} \\ x^2 - 2x - 3 \\ \underline{x^2 + 4x} \\ -6x - 3 \\ \underline{-6x - 24} \\ 21 \end{array}}$$




$$f(x) = \frac{(x-3)(x+1)}{x+4}$$

$$f(-5) = -32$$

Asymptote Note: The limit as x goes to infinity describes *end behavior* of the function.
 Although a graph cannot touch the vertical asymptote, it may cross over the horizontal asymptote.

Example:



The horizontal asymptote is $y = 2$
 There is no vertical asymptote...

Example: $f(x) = \frac{1}{x} - \frac{1}{|x|}$

Find $\lim_{x \rightarrow 0^+} f(x)$ $\lim_{x \rightarrow 0^-} f(x)$ $\lim_{x \rightarrow 0} f(x)$

Utilizing the graph of the function at the right,
 we can easily determine the limits!

$$\lim_{x \rightarrow 0^+} f(x) = 0$$


$$\lim_{x \rightarrow 0^-} f(x) = -\infty$$

$$\lim_{x \rightarrow 0} f(x) \text{ DNE}$$

does not exist...



Ultra-Marathon




Testing the limits of endurance,
these math figures will run on and on...

LanceAF #87 5-24-13
www.mathplane.com

PRACTICE QUIZ

Limits Quick Quiz

Part I: Identifying limits and values on a graph



1) $f(0) =$

2) $\lim_{x \rightarrow 0} =$

3) $\lim_{x \rightarrow -4} =$

4) $\lim_{x \rightarrow 3} =$

5) $\lim_{x \rightarrow -3^+} =$

6) $f(4) =$

7) $f(-3) =$

Part II: Finding Limits

1) $\lim_{x \rightarrow 3} \frac{2x - 6}{x^2 - 9}$

2) $\lim_{x \rightarrow 4} \frac{x^2 + 6x - 40}{3x + 6}$

3) $\lim_{x \rightarrow 9} \frac{\sqrt{x} - 3}{x - 9}$

4) $\lim_{x \rightarrow \infty} \frac{3 - 4x^2}{x^2 + 3x + 2}$

5) $\lim_{x \rightarrow 0} \frac{|x|}{3x} =$

6) $\lim_{x \rightarrow \infty} \frac{235}{3x + 2}$

7) $\lim_{x \rightarrow 2^+} \frac{4}{x - 2}$

8) $\lim_{x \rightarrow 2^-} \frac{4}{x - 2}$

9) $\lim_{x \rightarrow 2} \frac{4}{x - 2}$

10) $\lim_{x \rightarrow 2} \frac{x^2 - 4}{x^2 - 7x + 10}$

11) $\lim_{x \rightarrow 0} \frac{x^2 + 3x}{x}$

12) $\lim_{x \rightarrow \infty} \frac{2x^3 + 5x}{-3x^2 + 6}$

Part III: Graphing


1) Graph $f(x) = \begin{cases} 2x + 1 & \text{if } x > 1 \\ 2 - x & \text{if } x \leq 1 \end{cases}$

Then, identify: $f(1) =$

$$\lim_{x \rightarrow 1^+} f(x) =$$

$$\lim_{x \rightarrow 1^-} f(x) =$$

$$\lim_{x \rightarrow 1} f(x) =$$



2) Sketch a function with the following properties:


$$g(0) = 2$$

$$\lim_{x \rightarrow 2^+} g(x) = \infty$$



$$\lim_{x \rightarrow 2^-} g(x) = \infty$$

$$\lim_{x \rightarrow -\infty} g(x) = -1$$

$$\lim_{x \rightarrow +\infty} g(x) = -1$$



3) **Challenge:



Answer:

a) $\lim_{x \rightarrow 2} f(x) =$

b) $\lim_{x \rightarrow 2} g(x) =$

c) $\lim_{x \rightarrow 2} (f(x) + g(x)) =$

d) $\lim_{x \rightarrow 2} (f(x)g(x)) =$

IV. Miscellaneous Multiple Choice

1) As x increases to infinity, the function $f(x) = 2e^{-x}$ gets closer to

- a) 0
- b) $1/2$
- c) 2
- d) e
- e) infinity

2) A rational function of the form $y = \frac{ax}{x+b}$ has a vertical asymptote at $x = 5$ and a horizontal asymptote at $y = -3$

Which is a possible function?

- a) $\frac{5x}{x-3}$
- b) $\frac{3x}{x+5}$
- c) $\frac{-3x}{x-5}$
- d) $\frac{-5x}{x+3}$
- e) $\frac{-3x}{x+5}$

3) Let $p(x)$ be a cubic polynomial function, where $p(3) < 0$, $p(7) > 0$, and $p(9) < 0$,
Which statements are true?

- statement I: there are 3 zeros
- statement II: a zero exists at $x < 3$ OR $x > 9$
- statement III: for $p(x) = 0$, there are 2 solutions between 3 and 9

- a) I
- b) I and II
- c) I and III
- d) II
- e) I, II, and III

4) $\lim_{x \rightarrow 3} 9 =$

- a) 3
- b) 9
- c) Does not exist
- d) 0
- e) 27

5) Find the value of k so $g(x)$ is continuous:

$$g(x) = \begin{cases} k + x & x < 10 \\ xk & x \geq 10 \end{cases}$$

- a) 10
- b) 0
- c) $10/9$
- d) 1
- e) no solution

$$6) \lim_{t \rightarrow 4} \frac{t^2 - 16}{\frac{1}{4} - \frac{1}{t}}$$

- a) 4
- b) 16
- c) 64
- d) 128
- e) undefined

$$7) \lim_{x \rightarrow 0} \frac{(x+1)^2 - 1}{x}$$

- a) -1
- b) 0
- c) 1
- d) 2
- e) Does not exist

$$8) \lim_{x \rightarrow 3} \frac{x}{x^2 - 9}$$


- a) 3
- b) 9
- c) positive infinity
- d) negative infinity
- e) does not exist

$$9) \lim_{x \rightarrow 3} \frac{1}{(x-3)^2}$$

- a) 3
- b) 9
- c) positive infinity
- d) negative infinity
- e) does not exist


$$10) \lim_{x \rightarrow 4^+} \frac{x}{x^2 - 16}$$

- a) -16
- b) 0
- c) 4
- d) positive infinity
- e) does not exist



SOLUTIONS-→

Part I: Identifying limits and values on a graph



- 1) $f(0) = 4$
- 2) $\lim_{x \rightarrow 0} = 0$
- 3) $\lim_{x \rightarrow -4} = 2$
- 4) $\lim_{x \rightarrow 3} = \text{DNE}$ (Does Not Exist, because limit from the left (4) is not the same as limit from the right (-3))
- 5) $\lim_{x \rightarrow -3^+} = -\infty$
- 6) $f(4) = -4$
- 7) $f(-3) = \text{Undefined}$

Part II: Finding Limits

1) $\lim_{x \rightarrow 3} \frac{2x - 6}{x^2 - 9}$
 plug in value: $\frac{2(3) - 6}{(3)^2 - 9} = \frac{0}{0}$

factor and cancel: $\frac{2(x-3)}{(x+3)(x-3)}$
 plug in value again: $\frac{2}{(3+3)} = \frac{1}{3}$

4) $\lim_{x \rightarrow \infty} \frac{3 - 4x^2}{x^2 + 3x + 2}$

lead term of numerator: $-4x^2$ (degree: 2)
 lead term of denominator: x^2 (degree: 2)

Since the degrees are the same, use the coefficients:
 $\frac{-4}{1} = -4$

7) $\lim_{x \rightarrow 2^+} \frac{4}{x - 2}$

x	3	2.5	2.1	2.01	2.001
f(x)	4	8	40	400	4000

As x gets closer to 2, f(x) gets larger and larger ∞

10) $\lim_{x \rightarrow 2} \frac{x^2 - 4}{x^2 - 7x + 10}$

$f(2) = \frac{0}{0}$

factor the quadratics:

$\frac{(x+2)(x-2)}{(x-2)(x-5)}$ and plug in 2
 $\frac{4}{-3}$

2) $\lim_{x \rightarrow 4} \frac{x^2 + 6x - 40}{3x + 6}$
 plug in value: $\frac{(4)^2 + 6(4) - 40}{3(4) + 6} = \frac{0}{18} = 0$

5) $\lim_{x \rightarrow 0} \frac{|x|}{3x} = \text{DNE}$

$\lim_{x \rightarrow 0^-} = \frac{-1}{3}$
 $\lim_{x \rightarrow 0^+} = \frac{1}{3}$
 Since limits are not equal, then limit does not exist

8) $\lim_{x \rightarrow 2^-} \frac{4}{x - 2}$

x	1	1.5	1.9	1.99	1.999
f(x)	-4	-8	-40	-400	-4000

$-\infty$

11) $\lim_{x \rightarrow 0} \frac{x^2 + 3x}{x}$

$\lim_{x \rightarrow 0} \frac{x(x+3)}{x} = ((0) + 3) = 3$

3) $\lim_{x \rightarrow 9} \frac{\sqrt{x} - 3}{x - 9} \cdot \frac{\sqrt{x} + 3}{\sqrt{x} + 3}$ (multiply by the conjugate)
 $\frac{x-9}{(x-9)(\sqrt{x}+3)} = \frac{1}{\sqrt{x}+3}$
 limit is $\frac{1}{6}$

6) $\lim_{x \rightarrow \infty} \frac{235}{3x + 2}$

The degree of the denominator is greater than the degree of the numerator. Since it is "bottom heavy", the limit is 0 (as x gets larger and larger, the function decreases)

9) $\lim_{x \rightarrow 2} \frac{4}{x - 2}$

Does not exist.


limit from the left is negative ∞
 limit from the right is positive ∞
 Since they are not equal, limit DNE

12) $\lim_{x \rightarrow \infty} \frac{2x^3 + 5x}{-3x^2 + 6}$

Degree of numerator is greater than degree of denominator, so the equation would go to ∞

("top heavy")... Then, since there is a negative sign, equation goes to $-\infty$

(Note: sometimes, a quick sketch of the function can be helpful; or, it's a nice way to check your answers!)




1) Graph $f(x) = \begin{cases} 2x + 1 & \text{if } x > 1 \\ 2 - x & \text{if } x \leq 1 \end{cases}$

Then, identify: $f(1) = 1$

$\lim_{x \rightarrow 1^+} f(x) = 3$

$\lim_{x \rightarrow 1^-} f(x) = 1$

$\lim_{x \rightarrow 1} f(x) = \text{DNE}$ (Limit from the left and from the right are different, so limit does not exist)



2) Sketch a function with the following properties:


$g(0) = 2$ plot the point (0, 2)

$\lim_{x \rightarrow 2^+} g(x) = \infty$ vertical asymptote at 2, and function "goes up" on both sides of the asymptote...



$\lim_{x \rightarrow 2^-} g(x) = \infty$

$\lim_{x \rightarrow -\infty} g(x) = -1$ horizontal asymptote at $y = -1$

$\lim_{x \rightarrow +\infty} g(x) = -1$



3) **Challenge:



Answer:

a) $\lim_{x \rightarrow 2} f(x) = \text{DNE}$

b) $\lim_{x \rightarrow 2} g(x) = \text{DNE}$

c) $\lim_{x \rightarrow 2} (f(x) + g(x)) = 1$

d) $\lim_{x \rightarrow 2} (f(x)g(x)) = 0$

Note: according to limit theorems:
 $\lim (f + g) = \lim(f) + \lim(g)$
 $\lim(fg) = \lim(f)\lim(g)$
 But, that assumes the $\lim(f)$ and $\lim(g)$ exist. Since they do not, the theorems cannot be applied!

IV. Miscellaneous Multiple Choice

1) As x increases to infinity, the function $f(x) = 2e^{-x}$ gets closer to **SOLUTIONS**

- a) 0
- b) 1/2
- c) 2
- d) e
- e) infinity

rewrite function as $\frac{2}{e^x}$


as x gets infinitely larger, e^x goes to infinity...

therefore, $\frac{2}{e^x}$ gets smaller and smaller, approaching 0

2) A rational function of the form $y = \frac{ax}{x+b}$ has a vertical asymptote at $x = 5$ and a horizontal asymptote at $y = -3$

Which is a possible function?

- a) $\frac{5x}{x-3}$
- b) $\frac{3x}{x+5}$
- c) $\frac{-3x}{x-5}$
- d) $\frac{-5x}{x+3}$
- e) $\frac{-3x}{x+5}$




3) Let $p(x)$ be a cubic polynomial function, where $p(3) < 0$, $p(7) > 0$, and $p(9) < 0$. Which statements are true?

- statement I: there are 3 zeros
- statement II: a zero exists at $x < 3$ OR $x > 9$
- statement III: for $p(x) = 0$, there are 2 solutions between 3 and 9

- a) I
- b) I and II
- c) I and III
- d) II
- e) I, II, and III


(possible sketch)



polynomial function is continuous...

4) $\lim_{x \rightarrow 3} 9 = 9$

- a) 3
- b) 9
- c) Does not exist
- d) 0
- e) 27



5) Find the value of k so $g(x)$ is continuous:

$$g(x) = \begin{cases} k+x & x < 10 \\ xk & x \geq 10 \end{cases}$$

- a) 10
- b) 0
- c) 10/9
- d) 1
- e) no solution

to be continuous, each part of the piecewise function must meet:

$$k+x = xk$$

at $x = 10$:

$$\begin{aligned} 10+k &= 10k \\ 10 &= 9k \\ k &= 10/9 \end{aligned}$$

6) $\lim_{t \rightarrow 4} \frac{t^2 - 16}{\frac{1}{4} - \frac{1}{t}}$

- a) 4
- b) 16
- c) 64
- d) 128**
- e) undefined

substitute $t = 4$,
and the result is $\frac{0}{0}$ indeterminate

SOLUTIONS

$$\lim_{t \rightarrow 4} \frac{(t+4)(t-4)}{\frac{t}{4} - \frac{4}{4t}} = \lim_{t \rightarrow 4} \frac{(t+4)(t-4)}{\frac{(t-4)}{4t}} = \lim_{t \rightarrow 4} (t+4)(4t) = 128$$

7) $\lim_{x \rightarrow 0} \frac{(x+1)^2 - 1}{x}$

- a) -1
- b) 0
- c) 1
- d) 2**
- e) Does not exist

Using substitution, we see
the result is $0/0$ indeterminate

$$\lim_{x \rightarrow 0} \frac{x^2 + 2x + 1 - 1}{x}$$

so, we'll try expanding the numerator


$$\lim_{x \rightarrow 0} \frac{x^2 + 2x}{x} = \lim_{x \rightarrow 0} \frac{x(x+2)}{x} = 2$$

8) $\lim_{x \rightarrow 3} \frac{x}{x^2 - 9}$

- a) 3
- b) 9
- c) positive infinity
- d) **negative infinity**
- e) does not exist

Limit from the left is negative infinity
Limit from the right is positive infinity
Therefore, limit does not exist (DNE)

if $x = 3.1$, then numerator is positive
and denominator is positive..
if $x = 2.9$, then numerator is positive
and denominator is negative...



Note: These are limits;
approaching vertical asymptotes


9) $\lim_{x \rightarrow 3} \frac{1}{(x-3)^2}$

- a) 3
- b) 9
- c) positive infinity**
- d) negative infinity
- e) does not exist

Limit approaching 3 from the right is infinity
Limit approaching 3 from the left is infinity
Therefore, limit is positive infinity

$$\lim_{x \rightarrow 3^+} \frac{1}{(x-3)^2} = +\infty$$

$$\lim_{x \rightarrow 3^-} \frac{1}{(x-3)^2} = +\infty$$




10) $\lim_{x \rightarrow 4^+} \frac{x}{x^2 - 16}$

- a) -16
- b) 0
- c) 4
- d) positive infinity**
- e) does not exist

Limit approaching 4 from the right is infinity.
(at 4, the equation is $4/0$)

If $x = 4.1$, then $\frac{4.1}{(4.1)^2 - 16} = 5.06$


If $x = 4.01$, then $\frac{4.01}{(4.01)^2 - 16} = 50.06$



Thanks for visiting. (Hope it helped!)

If you have questions, suggestions, or requests, let us know.

Cheers,



Also, at Facebook, Google+, Pinterest, TES, and TeachersPayTeachers