

# Introduction to Integrals

Examples and practice questions (with solutions)

Topics include U-Substitution, logarithms, definite and indefinite antiderivatives, trigonometry, partial fractions, tabular integration, and more.

Strategies for finding Antiderivatives

1) "Simplify first"

Example:  $\int \frac{x^2 + 3x + 2}{\sqrt{x}} dx$

Splitting up the trinomial in the numerator creates 3 easier terms to integrate

$$\int \frac{x^2}{\sqrt{x}} + \frac{3x}{\sqrt{x}} + \frac{2}{\sqrt{x}} dx$$

$$\int \frac{x^2}{\sqrt{x}} dx + \int \frac{3x}{\sqrt{x}} dx + \int \frac{2}{\sqrt{x}} dx$$

$$\int x^{3/2} dx + \int 3x^{1/2} dx + \int 2x^{-1/2} dx$$

$$\frac{2}{5} x^{5/2} + 3 \cdot \frac{2}{3} x^{3/2} + 2 \cdot \frac{2}{1} x^{1/2}$$

$$\frac{2}{5} x^{5/2} + 2x^{3/2} + 4x^{1/2} + C$$

Example:  $\int \sin^2(3x) + \cos^2(3x) dx$

trigonometry identity

$$\sin^2 + \cos^2 = 1$$

$$\int 1 dx$$

$$x + C$$

Example:  $\int (x + 7)^2 (x^2 + 3x + 2) dx$

(expand and combine)

$$(x + 7)(x + 7) = x^2 + 14x + 49$$

Then,  $(x^2 + 3x + 2)(x^2 + 14x + 49)$


$$\begin{array}{r} x^4 + 14x^3 + 49x^2 \\ 3x^3 + 42x^2 + 147x \\ 2x^2 + 28x + 98 \\ \hline \end{array}$$

$$x^4 + 17x^3 + 93x^2 + 175x + 98$$

$$\int x^4 + 17x^3 + 93x^2 + 175x + 98 dx$$

$$\frac{x^5}{5} + \frac{17x^4}{4} + \frac{93x^3}{3} + \frac{175x^2}{2} + 98x + C$$

$$\frac{x^5}{5} + \frac{17x^4}{4} + 31x^3 + \frac{175x^2}{2} + 98x + C$$



Strategies for finding Antiderivatives

2) "Derivative beside the function"

Example:  $\int \frac{2x}{\sqrt{3x^2 + 5}} dx$

Note: This technique is similar to *integration by substitution* (or, *U-substitution*)

$$\int 2x \cdot (3x^2 + 5)^{-\frac{1}{2}} dx$$

The 'function' is  $3x^2 + 5...$   
and, its derivative is  $6x$

$$\frac{1}{3} \int 3 \cdot 2x \cdot (3x^2 + 5)^{-\frac{1}{2}} dx$$

Multiply 3 to get  $6x...$   
(and, multiply  $1/3$  to keep the same equation)

$$\frac{1}{3} \int 6x \cdot (3x^2 + 5)^{-\frac{1}{2}} dx$$

*Derivative beside the function then, use power rule...*

'derivative' 'function'

$$\frac{1}{3} \cdot \frac{2}{1} (3x^2 + 5)^{\frac{1}{2}} = \frac{2}{3} \sqrt{3x^2 + 5} + C$$

'function'

('derivative' disappears)

check the answer by taking the derivative!!

$$f(x) = \frac{2}{3} \sqrt{3x^2 + 5} + C$$

$$= \frac{2}{3} (3x^2 + 5)^{\frac{1}{2}} + C$$

use power rule to find derivative

$$f'(x) = \frac{2}{6} (3x^2 + 5)^{-\frac{1}{2}} (6x) + 0$$

('derivative' appears)

$$= \frac{2x}{\sqrt{3x^2 + 5}}$$

Example:  $\int x \cos(3x^2) dx$

There are two terms:  $x$  and  $3x^2$

Since the derivative of  $3x^2$  is  $6x$ ,  
we need a  $6x$  beside it!

$$\frac{1}{6} \int 6x \cos(3x^2) dx$$

$$\frac{1}{6} \sin(3x^2)$$

"derivative beside the function"

When you take the integral of the trig function, the  $6x$  'goes away'

CHECK:  $f(x) = \frac{1}{6} \sin(3x^2)$

$$f'(x) = \frac{1}{6} \cos(3x^2) \cdot (6x)$$

When you take the derivative of  $\cos(3x^2)$ , the  $6x$  'appears'  
(because of the chain rule)

$$f'(x) = x \cos(3x^2) \quad \checkmark$$

Strategies for finding Antiderivatives

2a) "U-Substitution"

Example:  $\int 4(8x + 3)^3 dx$

Expanding the expression  $(8x + 3)^3$  is one approach.  
But,  $(8x + 3)(8x + 3)(8x + 3)$  can get messy...

Using "U-Substitution" is another approach:

identify the 'main function'  
and label u

Let  $u = (8x + 3)$

determine du

then,  $\frac{du}{dx} = 8$        $dx = \frac{du}{8}$

substitute  $\int 4(8x + 3)^3 dx \longrightarrow \int 4(u)^3 \cdot \frac{du}{8} =$

solve  $\frac{4}{8} \int (u)^3 du = \frac{1}{2} \cdot \frac{u^4}{4} + C = \frac{u^4}{8} + C$

substitute  $\frac{(8x + 3)^4}{8} + C.$

(check) derivative of  $\frac{(8x + 3)^4}{8} + C = 4 \frac{(8x + 3)^3}{8} \cdot 8 + 0 = 4(8x + 3)^3$  ✓  
using chain rule

Compare "u-substitution" and "derivative beside function"

Example:  $\int \frac{3x}{\sqrt{x^2 + 8}} dx$

"Derivative beside the function"

$\int 3x(x^2 + 8)^{-\frac{1}{2}} dx$

the 'function' is  $(x^2 + 8)$ , so we need  
2x beside it...

$\frac{3}{2} \int \frac{2}{3} 3x(x^2 + 8)^{-\frac{1}{2}} dx$

$\frac{3}{2} \int \underbrace{2x}_{\text{derivative}} \underbrace{(x^2 + 8)^{-\frac{1}{2}}}_{\text{function}} dx$

now, integrate using power rule...

$\frac{3}{2} \frac{(x^2 + 8)^{\frac{1}{2}}}{\frac{1}{2}} = 3(x^2 + 8)^{\frac{1}{2}}$   
 $= 3\sqrt{x^2 + 8} + C$

"U-Substitution"

$\int 3x(x^2 + 8)^{-\frac{1}{2}} dx$

let  $u = (x^2 + 8)$

$\frac{du}{dx} = 2x \longrightarrow 2x dx = du \longrightarrow x dx = \frac{du}{2}$

$3 \int (x^2 + 8)^{-\frac{1}{2}} x dx$

substitute the terms

$3 \int u^{-\frac{1}{2}} \frac{du}{2} = \frac{3}{2} \int u^{-\frac{1}{2}} du$

$\frac{3}{2} \frac{u^{\frac{1}{2}}}{\frac{1}{2}} + C = 3u^{\frac{1}{2}} + C$

substitute back to x

$3\sqrt{x^2 + 8} + C$

Strategies for finding Antiderivatives

3) Integration by Parts

Review: Product Rule (derivatives)

If  $u = u(x)$  and  $v = v(x)$  and, therefore

$$u' = du = u'(x)dx \quad v' = dv = v'(x)dx$$

Then, derivative of  $u(x) \cdot v(x)$  is  $u'(x)dx \cdot v(x) + v'(x)dx \cdot u(x)$

or, derivative of  $u \cdot v$  is  $u'v + v'u$

This leads to integration by parts:

Since  $u \cdot v = u'v + v'u$

$$\int u \, dv = u \, v - \int v \, du$$

"If you have 2 continuous functions -- where one function's derivative is related to the other function -- try integration by parts."

Since the derivative of  $uv = u'v + v'u$ ,  
then the antiderivative of  $u'v + v'u = uv$

$$\int (u'v + v'u) = uv$$

$$\int u'v + \int v'u = uv$$

so,  $\int v'u = uv - \int u'v$

Example:  $\int 2xe^x \, dx$

Since  $2x$  and  $e^x$  are 'not related' (i.e. neither is the derivative of the other), we'll try integration by parts...

let  $u = 2x$  then,  $u' = 2 \, dx$   
 $v = e^x \, dx$   $v = e^x$

the derivative of  $2x$  is  $2$ ...  
 And  $2$  is related to  $x$ !...

(i.e. the derivative of  $x$  can lead to  $2$ )

$$\int 2xe^x \, dx = 2x \, e^x - \int e^x \, 2 \, dx$$

$$= 2x \, e^x - 2 \, e^x + C$$

$$= 2e^x(x - 1) + C$$

Example:  $\int x \sin x \, dx$

let  $u = x$  then,  $u' = 1 \, dx$   
 $v = \sin x \, dx$   $v = -\cos(x)$

$$\int x \sin x \, dx = x(-\cos(x)) - \int (-\cos(x)) \, dx$$

$$= -x\cos(x) - (-\sin(x)) + C$$

$$= \sin(x) - x\cos(x) + C$$

'derivative of  $x$  is related to  $(x)$ '

Now, suppose we let  $u = \sin x$  then,  $u' = \cos x \, dx$   
 $v = x \, dx$   $v = \frac{x^2}{2}$

$$\int x \sin x \, dx = \sin x \cdot \frac{x^2}{2} - \int \frac{x^2}{2} \cos x \, dx$$

'derivative of  $\sin(x)$  is not related to  $x$ '  
 (the derivative of  $\sin(x)$  cannot lead to  $x$ )

In this case, we would have to use integration by parts again!  
 \*\*\*Note: selecting the correct "u" and "v" is important!

Example:  $\int x^3 e^{-2x} dx$

$u = x^3$        $v' = e^{-2x} dx$   
 $u' = 3x^2 dx$      $v = \frac{-1}{2} e^{-2x}$

Integration by Parts Formula  
 $\int v'u = uv - \int u'v$

$$\begin{aligned} \int x^3 e^{-2x} dx &= x^3 \cdot \frac{-1}{2} e^{-2x} - \int 3x^2 dx \cdot \frac{-1}{2} e^{-2x} \\ &= \frac{-1}{2} \cdot x^3 e^{-2x} - \frac{-1}{2} \int 3x^2 e^{-2x} dx \\ &= \frac{-1}{2} \left[ x^3 e^{-2x} - \int 3x^2 e^{-2x} dx \right] \end{aligned}$$

Now, we'll use integration by parts again...

$u = 3x^2$        $v' = e^{-2x} dx$   
 $u' = 6x dx$      $v = \frac{-1}{2} e^{-2x}$

$$\begin{aligned} &= \frac{-1}{2} \left[ x^3 e^{-2x} - \left( 3x^2 \cdot \frac{-1}{2} e^{-2x} - \int 6x dx \cdot \frac{-1}{2} e^{-2x} \right) \right] \\ &= \frac{-1}{2} \left[ x^3 e^{-2x} - \left( \frac{-1}{2} \cdot 3x^2 e^{-2x} - \frac{-1}{2} \int 6x e^{-2x} dx \right) \right] \\ &= \frac{-1}{2} \left[ x^3 e^{-2x} - \frac{-1}{2} \left( 3x^2 e^{-2x} - \int 6x e^{-2x} dx \right) \right] \end{aligned}$$

Now, we'll use integration by parts again...

$u = 6x$        $v' = e^{-2x} dx$   
 $u' = 6 dx$      $v = \frac{-1}{2} e^{-2x}$


$$\begin{aligned} &= \frac{-1}{2} \left[ x^3 e^{-2x} - \frac{-1}{2} \left( 3x^2 e^{-2x} - \left( 6x \cdot \frac{-1}{2} e^{-2x} - \int 6 dx \cdot \frac{-1}{2} e^{-2x} \right) \right) \right] \\ &= \frac{-1}{2} \left[ x^3 e^{-2x} - \frac{-1}{2} \left( 3x^2 e^{-2x} - \left( \frac{-1}{2} \cdot 6x e^{-2x} - \frac{-1}{2} \int 6 e^{-2x} dx \right) \right) \right] \\ &= \frac{-1}{2} \left[ x^3 e^{-2x} - \frac{-1}{2} \left( 3x^2 e^{-2x} - \frac{-1}{2} \left( 6x e^{-2x} - \int 6 e^{-2x} dx \right) \right) \right] \end{aligned}$$

At last, we can determine the final integral!

$$\begin{aligned} &= \frac{-1}{2} \left[ x^3 e^{-2x} - \frac{-1}{2} \left( 3x^2 e^{-2x} - \frac{-1}{2} \left( 6x e^{-2x} - \left( \frac{-1}{2} \cdot 6 e^{-2x} \right) \right) \right) \right] \\ &= \frac{-1}{2} \left[ x^3 e^{-2x} - \frac{-1}{2} \left( 3x^2 e^{-2x} - \frac{-1}{2} \left( 6x e^{-2x} - \frac{-1}{2} \left( 6 e^{-2x} \right) \right) \right) \right] \\ &= \frac{-1}{2} x^3 e^{-2x} - \frac{1}{4} \cdot 3x^2 e^{-2x} + \frac{-1}{8} \cdot 6x e^{-2x} - \frac{1}{16} \cdot 6 e^{-2x} \end{aligned}$$

Using Tabular Integration:  
 Observe: there is a pattern!

Derivative		Integral
$x^3$	+	$e^{-2x}$
$3x^2$	-	$\frac{-1}{2} e^{-2x}$
$6x$	+	$\frac{1}{4} e^{-2x}$
$6$	-	$\frac{-1}{8} e^{-2x}$
$0$		$\frac{1}{16} e^{-2x}$



4) Utilizing Partial Fractions

Strategies for finding Antiderivatives

Example:  $\int \frac{8x+5}{x^2+3x-10} dx$   
 $(x+5)(x-2)$

$$\frac{A}{(x+5)} + \frac{B}{(x-2)} = \frac{8x+5}{x^2+3x-10}$$

$$\frac{A(x-2)}{(x+5)(x-2)} + \frac{B(x+5)}{(x+5)(x-2)} = \frac{8x+5}{x^2+3x-10}$$

$$A(x-2) + B(x+5) = 8x+5$$

$$Ax - 2A + Bx + 5B = 8x + 5$$

$$(A+B)x - 2A + 5B = 8x + 5$$

Then, we know

$$\begin{array}{r} A + B = 8 \\ -2A + 5B = 5 \end{array}$$

$$\begin{array}{r} 2A + 2B = 16 \\ -2A + 5B = 5 \\ \hline 7B = 21 \\ B = 3 \end{array}$$

then,  $A = 5$

$$\frac{5}{(x+5)} + \frac{3}{(x-2)}$$

$$\int \frac{8x+5}{x^2+3x-10} dx = \int \frac{5}{(x+5)} dx + \int \frac{3}{(x-2)} dx$$

$$5\ln|x+5| + 3\ln|x-2| + C$$

Example:  $\int \frac{-6x^2+3x+5}{x^3-x} dx$   
 $x(x^2-1) = x(x+1)(x-1)$

$$\frac{-6x^2+3x+5}{x^3-x} = \frac{A}{x} + \frac{B}{(x+1)} + \frac{C}{(x-1)}$$

$$= \frac{(x+1)(x-1)A}{x(x+1)(x-1)} + \frac{x(x-1)B}{x(x+1)(x-1)} + \frac{x(x+1)C}{x(x+1)(x-1)}$$

$$-6x^2+3x+5 = (x+1)(x-1)A + x(x-1)B + x(x+1)C$$

$$Ax^2 - 1A + Bx^2 - Bx + Cx^2 + Cx$$

(regroup the terms)

$$(A+B+C)x^2 + (-B+C)x + A(-1)$$

$$-6x^2 + 3x + 5$$

$$\begin{array}{r} A + B + C = -6 \\ -B + C = 3 \\ -A = 5 \end{array}$$

$$\begin{array}{r} A = -5 \\ -5 + B + C = -6 \\ -B + C = 3 \\ \hline 2C = 2 \\ C = 1 \end{array}$$

$$\begin{array}{r} -B + C = 3 \\ -B + 1 = 3 \\ B = -2 \end{array}$$

"Express method"

Let  $x = 0$  (to eliminate B and C)  
 $0 + 0 + 5 = -1A + 0B + 0C$   
 $A = -5$

Let  $x = -1$  (to eliminate A and C)  
 $-6 + (-3) + 5 = 0A + 2B + 0C$   
 $B = -2$

Let  $x = 1$  (to eliminate A and B)  
 $-6 + 3 + 5 = 0A + 0B + 2C$   
 $C = 1$

$$\int \frac{-5}{x} + \frac{-2}{(x+1)} + \frac{1}{(x-1)} dx$$

$$-5\ln|x| - 2\ln|x+1| + \ln|x-1| + C$$

$$\frac{-6x^2+3x+5}{x^3-x} = \frac{-5}{x} + \frac{-2}{(x+1)} + \frac{1}{(x-1)}$$

Example:

$$\int_{13}^{18} x^2 \sqrt{x-9} \, dx$$

Let  $U = x - 9$   $\Rightarrow$  NOTE: the boundaries for x are 13 - 18  
 $x = U + 9$   
 $\frac{dU}{dx} = 1$  so,  $dU = dx$

$$\int_4^9 (U+9)^2 (U)^{\frac{1}{2}} \, dU$$

$$\int_4^9 (U^2 + 18U + 81) (U)^{\frac{1}{2}} \, dU$$

$$\int_4^9 U^{\frac{5}{2}} + 18U^{\frac{3}{2}} + 81U^{\frac{1}{2}} \, dU$$

$$\left. \frac{2}{7} U^{\frac{7}{2}} + \frac{36}{5} U^{\frac{5}{2}} + 54U^{\frac{3}{2}} \right|_4^9$$

Also,

$$\int U^{\frac{5}{2}} + 18U^{\frac{3}{2}} + 81U^{\frac{1}{2}} \, dU$$

$$\frac{2}{7} U^{\frac{7}{2}} + \frac{36}{5} U^{\frac{5}{2}} + 54U^{\frac{3}{2}}$$

put back into terms of x

$$\frac{2}{7} (x-9)^{\frac{7}{2}} + \frac{36}{5} (x-9)^{\frac{5}{2}} + 54(x-9)^{\frac{3}{2}}$$

18  
13

$$\frac{2}{7} (3)^7 + \frac{36}{5} (3)^5 + 54(3)^3 - \left( \frac{2}{7} (2)^7 + \frac{36}{5} (2)^5 + 54(2)^3 \right) = \frac{109672}{35}$$

Example:

$$\int_5^9 t^3 \sqrt{t-4} \, dt$$

Let  $U = t - 4$   
 $t = U + 4$

$$\int_1^5 (U+4)^3 (U)^{\frac{1}{2}} \, dU$$

$\frac{dU}{dt} = 1$   $dU = dt$

$$\int_1^5 (U^3 + 12U^2 + 48U + 64) (U)^{\frac{1}{2}} \, dU$$

$$\int_1^5 U^{\frac{7}{2}} + 12U^{\frac{5}{2}} + 48U^{\frac{3}{2}} + 64(U)^{\frac{1}{2}} \, dU = \left. \frac{2}{9} U^{\frac{9}{2}} + \frac{24}{7} U^{\frac{7}{2}} + \frac{96}{5} U^{\frac{5}{2}} + \frac{128(U)}{3} \right|_1^5 = \frac{79430\sqrt{5}}{63} - \frac{20638}{315}$$

Example:

$$\int x \sqrt{2x+1} \, dx$$

Let  $U = 2x + 1$   $\frac{dU}{dx} = 2$   
 $x = \frac{U-1}{2}$   $dx = \frac{dU}{2}$

$$\int \frac{U-1}{2} \cdot U^{\frac{1}{2}} \cdot \frac{dU}{2}$$

$$\frac{1}{4} \int U^{\frac{3}{2}} + U^{\frac{1}{2}} \, dU = \frac{1}{4} \left( \frac{2U^{\frac{5}{2}}}{5} - \frac{2U^{\frac{3}{2}}}{3} \right) + C \Rightarrow$$

$$(2x+1)^{\frac{3}{2}} \left[ \frac{3x-1}{15} \right] + C$$

$$\frac{1}{10} (2x+1)^{\frac{5}{2}} - \frac{1}{6} (2x+1)^{\frac{3}{2}} + C$$

$$(2x+1)^{\frac{3}{2}} \left[ \frac{1}{10} (2x+1) - \frac{1}{6} \right] + C$$

$$(2x+1)^{\frac{3}{2}} \left[ \frac{3}{30} (2x+1) - \frac{5}{30} \right] + C$$



Example:  $\int y \ln(y) dy$   
 $\int y \cdot \ln(y) dy$

since the derivative of  $\ln y$  is  $1/y$ ,  
 we won't use u-substitution...  
 Instead, we'll try integration by parts...

Integration by Parts

Step 3: Apply the formula

Step 1: Identify "u" and "v"

Since it is easier to take the derivative of  $\ln(y)$ ,

$u = \ln(y)$   
 $v' = y dy$

Step 2: Find other parts

$u' = \frac{1}{y} dy$   
 $v = \int y dy = \frac{y^2}{2}$

$$\int v'u = uv - \int u'v$$

$$\int y dy \cdot \ln(y) = \ln(y) \cdot \frac{y^2}{2} - \int \frac{1}{y} dy \cdot \frac{y^2}{2}$$

$$\ln(y) \cdot \frac{y^2}{2} - \int \frac{y}{2} dy$$

$\ln(y) \cdot \frac{y^2}{2} - \frac{y^2}{4} + C$

Example:  $\int \frac{2x^2 + 7x - 3}{x - 2} dx$   
 $\int 2x + 11 + \frac{19}{x - 2} dx$

Divide using  
 synthetic division  
 Then, integrate...

$$\begin{array}{r|rrr} 2 & 2 & 7 & -3 \\ & & 4 & 22 \\ \hline & 2 & 11 & 19 \end{array}$$

$$2x + 11 + \frac{19}{x - 2}$$

Synthetic Division  
 (or, polynomial division)

$x^2 + 11x + \ln|x - 2| + C$

Example:  $\int (3x + 5)^2 dx$

Evaluate the indefinite integral using a) U-substitution and  
 b) Expanding the term... then, compare your results...

Let  $u = 3x + 5$   
 so,  $\frac{du}{dx} = 3 \quad dx = \frac{1}{3} du$

$\int u^2 \cdot \frac{1}{3} du$   
 $\frac{1}{3} \cdot \frac{u^3}{3} = \frac{1}{9} (3x + 5)^3 + C$

$\frac{1}{9} (27x^3 + 135x^2 + 225x + 125) + C$   
 $3x^3 + 15x^2 + 25x + \frac{125}{9} + C$  ✓

$\int 9x^2 + 30x + 25 dx$   
 $3x^3 + 15x^2 + 25x + C$  ✓

Since C is any constant...  
 C is equivalent to  $\frac{125}{9} + C$

Example:  $\int \frac{10^x}{\ln 10} dx$

(substitute the dx)


Let  $U = 10^x$   
 $\frac{dU}{dx} = (\ln 10)(10^x)$   
 $(\ln 10)(10^x) dx = dU$   
 $dx = \frac{dU}{(\ln 10)(10^x)}$

$\int \frac{10^x}{\ln 10} \frac{dU}{(\ln 10)(10^x)}$   
 $\int \frac{1}{\ln 10} \frac{dU}{(\ln 10)}$

(Remember  $\ln 10$  is a constant/number... So, we're ready to integrate)

$\frac{1}{(\ln 10)^2} \int dU = \frac{1}{(\ln 10)^2} U + C$   

$\frac{1}{(\ln 10)^2} 10^x + C$



Happy Days

*Despite Richie's help, Fonzie dropped out of Calculus.  
(... although he did have some success with velocity and acceleration!)*

Practice ->

Indefinite Integrals Exercise

Evaluate the indefinite integrals. (Check your results by differentiation!)

1)  $\int (x^3 + 4) dx$

2)  $\int \sqrt{x^3} dx$

3)  $\int \frac{1}{6x^2} dx$

4)  $\int \frac{(x+2)^2}{x} dx$

5)  $\int dx$

6)  $\int t^2 (t^3 + 1)^4 dt$

7)  $\int \frac{2x}{\sqrt{x^2 - 3}} dx$

8)  $\int 8(3 + 4x^2)^2 dx$

9)  $\int \cos 2t dt$

For the following, find the specific or general functions:

1)  $f'(x) = 8x^3 + 10x + 5$     $f(1) = 6$

\*\*\*CHALLENGE\*\*\*

2)  $f''(x) = 3 + x^2 + x^5$

Antiderivatives

3)  $f''(x) = 2 - 12x$     $f(0) = 9$     $f(2) = 15$

4)  $f'(x) = \frac{1}{\cos^2 x} + 3^x$

Indefinite Integrals Exercise

Evaluate the indefinite integrals. (Check your results by differentiation!)

SOLUTIONS

$$1) \int (x^3 + 4) dx$$

$$\int x^3 dx + \int 4 dx$$

$$\frac{x^4}{4} + 4x + C$$

the derivative of the solution:

$$\frac{4x^3}{4} + 4 + 0 = x^3 + 4 \checkmark$$

$$4) \int \frac{(x+2)^2}{x} dx$$

(expand numerator)

$$\int \frac{x^2 + 4x + 4}{x} dx$$

$$\int x + 4 + 4\left(\frac{1}{x}\right) dx$$

$$\frac{x^2}{2} + 4x + 4\ln|x| + C$$

$$7) \int \frac{2x}{\sqrt{x^2-3}} dx$$

$$\int 2x(x^2-3)^{-\frac{1}{2}} dx$$

$$u = x^2 - 3$$

$$u' = 2x$$

so, we can use power rule of integration

$$\frac{(x^2-3)^{\frac{1}{2}}}{\frac{1}{2}}$$

$$2\sqrt{x^2-3} + C$$

$$2) \int \sqrt{x^3} dx$$

$$\int (x^3)^{\frac{1}{2}} dx = \int x^{\frac{3}{2}} dx$$

$$\frac{x^{\frac{5}{2}}}{\frac{5}{2}} = \frac{2x^{\frac{5}{2}}}{5} + C$$

(derivative, using power rule)

$$\frac{5}{2} \cdot \frac{2x^{\frac{3}{2}}}{5} + 0 = x^{\frac{3}{2}} \checkmark$$

$$5) \int dx$$

$$\int 1 dx$$

$$x + C$$

$$3) \int \frac{1}{6x^2} dx$$

$$\int \frac{1}{6} x^{-2} dx = \frac{1}{6} \int x^{-2} dx$$

$$= \frac{1}{6} \left( \frac{x^{-1}}{-1} \right) = \frac{-1}{6x} + C$$

derivative of  $-1/6x$  (using quotient rule)

$$\frac{0(6x) - 6(-1)}{(6x)^2} = \frac{6}{36x^2} = \frac{1}{6x^2} \checkmark$$

$$6) \int t^2(t^3+1)^4 dt$$

Let  $u = (t^3+1)$  then, we need  $u' = 3t^2$

$$\frac{1}{3} \int \underset{u'}{3t^2} \underset{u}{(t^3+1)^4} dt = \frac{1}{3} \int \frac{(t^3+1)^5}{5} dt$$

$$= \frac{(t^3+1)^5}{15} + C$$

derivative:

$$\frac{5(t^3+1)^4 \cdot (3t^2+0)}{15} + 0 = (t^3+1)^4 \cdot t^2 \checkmark$$

$$8) \int 8(3+4x^2)^2 dx$$

If  $u = (3+4x^2)$ , then  $u' = 8x$   
since there is no  $x$  term, the power rule of integration cannot be used. instead expand the entire equation.

$$\int 8(9+24x^2+16x^4) dx$$

$$\int 72 + 192x^2 + 128x^4 dx$$

$$72x + 64x^3 + \frac{128x^5}{5} + C$$

$$9) \int \cos 2t dt$$

The derivative of  $\sin t$  is  $\cos t$ , and the derivative of  $\sin 2t$  is  $2\cos 2t$ .

$$\frac{1}{2} \int 2 \cos 2t dt$$

$$\frac{1}{2} \sin 2t + C$$

For the following, find the specific or general functions:

Antiderivatives

$$1) f'(x) = 8x^3 + 10x + 5 \quad f(1) = 6$$

Find the indefinite integral:

$$2x^4 + 5x^2 + 5x + C$$

Then, use the given value to determine C:

$$2(1)^4 + 5(1)^2 + 5(1) + C = 6$$

$$2 + 5 + 5 + C = 6$$

$$C = -6$$

$$f(x) = 2x^4 + 5x^2 + 5x - 6$$

$$3) f''(x) = 2 - 12x \quad f(0) = 9 \quad f(2) = 15$$

Take the antiderivative twice to determine the function:

$$f'(x) = 2x - 6x^2 + C$$

$$f(x) = x^2 - 2x^3 + Cx + D$$

then, use the given values to find the specific function:

$$f(0) = (0)^2 - 2(0)^3 + C(0) + D$$

$$D = 9$$

$$\text{so, } f(x) = x^2 - 2x^3 + Cx + 9$$

$$f(2) = (2)^2 - 2(2)^3 + C(2) + 9$$

$$15 = 4 - 16 + 2C + 9$$

$$C = 9$$

$$f(x) = -2x^3 + x^2 + 9x + 9$$

\*\*\*CHALLENGE\*\*\*

$$2) f''(x) = 3 + x^2 + x^5$$

SOLUTIONS

Find the antiderivative:

$$f'(x) = 3x + \frac{x^3}{3} + \frac{x^6}{6} + C \quad \text{where } C \text{ is a constant}$$

Then, find the antiderivative again:

$$f(x) = \frac{3x^2}{2} + \frac{x^4}{12} + \frac{x^7}{42} + Cx + D$$

where C and D are constants...

$$4) f'(x) = \frac{1}{\cos^2 x} + 3^x$$

to find the antiderivative (integral), separate the terms and rewrite...

$$\int \sec^2 x \, dx + \int 3^x \, dx$$

The derivative of what term equals  $\sec^2 x$  ?

the answer:  $\tan x$

$$\tan x + \int 3^x \, dx$$

The derivative of  $3^x$  is  $3^x \cdot \ln 3$

so, the antiderivative of  $(\ln 3)3^x$  would be  $3^x$


$$\tan x + \frac{1}{\ln 3} \int \ln 3 \cdot 3^x \, dx$$

$$\tan x + \frac{1}{\ln 3} \cdot 3^x = \tan x + \frac{3^x}{\ln 3} + C$$

Thanks for visiting. (Hope it helps!)

If you have questions, suggestions, or feedback, let us know.

Cheers.



Also, at Facebook, Google +, TeacherPayTeachers, and TES, Pinterest

And, Mathplane *Express* for mobile at [mathplane.ORG](http://mathplane.ORG)

---

One more question:

$$\int \sin^5 x \cdot (\cos^{20} x) dx$$

Answer on next page-→

Trigonometry Integral Question  
Steps and Answer

$$\int \sin^5 x \cdot (\cos^{20} x) dx$$

$$\int \sin x \cdot \sin^4 x \cdot (\cos^{20} x) dx$$

$$\int \sin x \cdot (\sin^2 x)^2 \cdot (\cos^{20} x) dx$$

$$\int \sin x \cdot (1 - \cos^2 x)^2 \cdot (\cos^{20} x) dx$$

$$\int \sin x \cdot (1 - 2\cos^2 x + \cos^4 x) \cdot (\cos^{20} x) dx$$

$$\int \sin x \cdot (\cos^{20} x - 2\cos^{22} x + \cos^{24} x) dx$$

$$\int \sin x (\cos^{20} x) - 2\sin x (\cos^{22} x) + \sin x (\cos^{24} x) dx$$

$$\int \sin x (\cos^{20} x) dx + \int -2\sin x (\cos^{22} x) dx + \int \sin x (\cos^{24} x) dx$$

$$- \int -\sin x (\cos^{20} x) dx + \int -2\sin x (\cos^{22} x) dx + - \int -\sin x (\cos^{24} x) dx$$

$$- \left[ \frac{1}{21} \cos^{21} x \right] + \frac{2}{23} \cos^{23} x + - \left[ \frac{1}{25} \cos^{25} x \right]$$

$$\boxed{-\frac{1}{21} \cos^{21} x + \frac{2}{23} \cos^{23} x - \frac{1}{25} \cos^{25} x + C}$$