


# Calculus, Natural Log, and $e$

## Practice Test and Solutions



mathplane.com

*Topics include logarithms, area, tangent lines, implicit differentiation, graphing, inverses, partial fractions, and more.*

Calculus: Logarithms,  $\ln$ , and  $e$

I. Logarithm Review

1) Answer:  $\ln 1 =$   $\ln e =$   $\ln 0 =$

(no calculator)

2)  $\log 4 = .602$   $\log 3 = .477$

Find:  $\log 12 =$   $\log 400 =$   $\log(.75) =$

(no calculator)

3) Solve for x:

A)  $\log_5 x + \log_5 (x-4) = 1$

B)  $3^x = 8$

C)  $2^{6-x} = 4^{2+x}$

Challenge:  $4^x - 2^{x+1} = 1$

II. Calculus:  $e$  and  $\ln$

Calculus: Logarithms,  $\ln$ , and  $e$

1) Find  $\frac{dy}{dx}$

$$y = e^{2x}$$

$$y = -e^{-x}$$

$$y = e^2$$

$$y = \ln(2x + 4)$$

$$y = \ln(3)$$

$$y = \ln(x + 3)^2$$

$$y = \ln((x + 3)^2)$$

$$y = \frac{2}{e^{3x}}$$

2) What is the equation of the line tangent to  $y = e^{2x-3}$  at the point  $(\frac{3}{2}, 1)$  ?

Optional: graph your result

3) What is the equation of the normal to  $y = \ln(x - 2) + 4$  at the point  $(3, 4)$  ?

Optional: graph your result

4)  $\int e^{2x} dx$

$\int \frac{3x}{3x^2 + 2} dx$

$\int \frac{2}{3x + 3} dx$

5) What is the area of the region above the x-axis that is bounded by the y-axis,  $x = 3$ , and  $e^x$  ?

6) What is the area of the region bounded by  $y = \ln(x) + 2$ ,  $y = 2$ , and  $x = 5$  ?  
(Use Calculator)

- 7) Find the equation of the line that is tangent to  $f(x) = 3x^2 - \ln x$  at  $(1, 3)$   
(Optional: Use a graphing calculator to confirm your answer)

8)  $y = \frac{x^2}{2} - \ln x$

What are the extrema?

Points of inflection?

(Optional: Use a graphing calculator  
to check your answers)

III. Inverses and derivatives

1)  $f(x)$  and  $g(x)$  are one to one inverses.

If the slope of  $f(x)$  at  $(3, 8)$  is 2, where is the slope of  $g(x)$  equal to  $1/2$ ?

2)  $f(x) = x^3 - x - 6$

What is  $f^{-1}(0)$ ?

3)  $h(x) = \ln(x) + 4$


a) What is the inverse of  $h(x)$ ?

b)  $h(3) = \ln(3) + 4$  ( $\approx 5.1$ )      What is the slope at  $h(3)$ ?

c) Graph  $h(x)$  and  $h^{-1}(x)$

Sketch the tangent lines at  $(3, 5.1)$  and  $(5.1, 3)$

What are the equations of the tangent lines?



IV. Exponential Functions

Find the first derivatives:


1)  $g(x) = 2^{x+3}$

2)  $f(x) = x^2 e^x$

3)  $g(t) = t^2 2^t$


4) What is the equation of the line tangent to  $y = 2^{-x}$  at  $(0, 1)$ ?

(optional: sketch a graph containing the function and tangent line)



Family  
E-union

"This was taken at my grandpa's 80th birthday party. That's my uncle Ed and aunt Edie in the fourth row..."



"I can see the resemblance!"



LAF #103 9-13-13  
mathplanE.com

SOLUTIONS-→



Calculus: Logarithms, ln, and e

I. Logarithm Review

1) Answer:  $\ln 1 = \log_e 1 = x$        $\ln e = \log_e e = y$        $\ln 0 = \log_e 0 = z$   
 (no calculator)       $e^x = 1$        $e^y = e$        $e^z = 0$   
 $x = 0$        $y = 1$       **No solution**  
 (logarithms cannot be zero or negative)

2)  $\log 4 = .602$        $\log 3 = .477$   
 Find:  $\log 12 = \log(3 \cdot 4)$        $\log 400 = \log(4 \cdot 10)$        $\log(.75) = \log \frac{3}{4}$   
 (no calculator)       $= \log 3 + \log 4$        $= \log 4 + \log 10$        $= \log 3 + \log 4$   
 $= .477 + .602 = 1.079$        $= .602 + 1 = 1.602$        $= .477 - .602 = -.125$

3) Solve for x:

A)  $\log_5 x + \log_5 (x-4) = 1$   
 $\log_5 x(x-4) = 1$   
 $5^1 = x(x-4)$   
 $5 = x^2 - 4x$   
 $x^2 - 4x - 5 = 0$   
 $(x-5)(x+1) = 0$   
 $x = 5, -1$   $\log(-1)$  does not exist

B)  $3^x = 8$   
 $\log 3^x = \log 8$   
 $x \log 3 = \log 8$   
 $x = \frac{\log 8}{\log 3}$   
 $= \frac{.903}{.477} = 1.89$   
 quick check:  
 $3^{1.89} = 8$  ✓

C)  $2^{6-x} = 4^{2+x}$   
 $2^{6-x} = (2^2)^{2+x}$  quick check:  
 $2^{6-x} = 2^{4+2x}$   $2^{5.33} = 4^{2.67}$   
 $6-x = 4+2x$   $40.3 = 40.3$  ✓  
 $x = \frac{2}{3}$

Challenge:  $4^x - 2^{x+1} = 1$

$4^x - 2^{x+1} - 3 = 0$   
 $(2^2)^x - (2^x)(2^1) - 3 = 0$   
 $(2^x)^2 - (2^x)(2^1) - 3 = 0$

therefore,  
 $2^x = -1$  and  $3$   
 $-1$  is extraneous!  
 $2^x = 3$

approximately 1.585

Let  $y = 2^x$   
 $y^2 - 2y - 3 = 0$   
 $(y-3)(y+1) = 0$   
 $y = -1, 3$

$2^x = 3$	$2^x = -1$
$X \log 2 = \log 3$	$x \log 2 = \log(-1)$
$X = \log 3 / \log 2$	$x = \log(-1) / \log(2)$
$X = 1.5849625$	$x$ does not exist

Check:  
 $4^{1.585} - 2^{2.585} =$   
 $9 - 6 = 3$  ✓

II. Calculus:  $e$  and  $\ln$

- 1) Find  $\frac{dy}{dx}$  for  $y = e^u$  "derivative of exponent times itself"  
 for  $y = \ln(u)$  "derivative over itself"

$y = e^{2x}$

$2e^{2x}$

$y = -e^{-x}$

$(-1)(-e^{-x})$

$e^{-x}$  or  $\frac{1}{e^x}$

$y = e^2$

0

$e^2$  is a constant

$y = \ln(2x + 4)$

$\frac{2}{2x + 4}$  or  $\frac{1}{x + 2}$

$y = \ln(3)$

0

$\ln(3)$  is a constant

$y = \ln(x + 3)^2$

$2(\ln(x + 3))^1 \cdot \frac{1}{(x + 3)}$

$\frac{2\ln(x + 3)}{(x + 3)}$

$y = \ln((x + 3)^2)$

$\frac{2(x + 3)}{(x + 3)^2}$

$\frac{2}{x + 3}$

$y = \frac{2}{e^{3x}}$

$2 \cdot e^{-3x}$

$-6e^{-3x}$  or  $-\frac{6}{e^{3x}}$

- 2) What is the equation of the line tangent to  $y = e^{2x-3}$  at the point  $(\frac{3}{2}, 1)$  ?

Optional: graph your result

To find the equation of a line, we need a point and the slope.

Point:  $(\frac{3}{2}, 1)$


Slope: rate of change at  $x = \frac{3}{2}$

$y' = 2 \cdot e^{2x-3}$

at  $(\frac{3}{2}, 1)$

$y' = 2 \cdot e^0 = 2$

tangent line:  $y - 1 = 2(x - \frac{3}{2})$



- 3) What is the equation of the normal to  $y = \ln(x - 2) + 4$  at the point  $(3, 4)$  ?

Optional: graph your result

The normal is perpendicular to the tangent line

Point:  $(3, 4)$


Find  $\frac{dy}{dx}$  to get instantaneous rate of change (i.e. slope)

$\frac{dy}{dx} = \frac{1}{x - 2} + 0$

slope at  $(3, 4)$  is 1 (tangent slope)

therefore, opposite reciprocal is -1 (normal slope)

normal line:  $y - 4 = -1(x - 3)$



$$4) \int e^{2x} dx$$

$$\frac{1}{2} \int 2e^{2x} dx$$

$$\frac{1}{2} e^{2x} + C$$

$$\int \frac{3x}{3x^2+2} dx$$

$$\frac{1}{2} \int \frac{2 \cdot 3x}{3x^2+2} dx$$

$$\frac{1}{2} \ln(3x^2+2) + C$$

$$\int \frac{2}{3x+3} dx$$

$$\int \frac{2}{3(x+1)} dx$$

$$\frac{2}{3} \int \frac{1}{(x+1)} dx = \frac{2}{3} \ln(x+1) + C$$

or  $\ln(x+1)^{\frac{2}{3}} + C$

5) What is the area of the region above the x-axis that is bounded by the y-axis, x = 3, and  $e^x$ ?

A quick sketch will show the enclosed region (and its boundaries)


The endpoints of the integral will be  $x = 0$  and  $x = 3$

and, the upper boundary will be  $y = e^x$

and the lower boundary will be  $y = 0$

$$\int_0^3 e^x dx = e^x \Big|_0^3 = e^3 - e^0 \approx 19.09$$

$$\int_0^3 e^x dx - \int_0^3 0 dx$$



6) What is the area of the region bounded by  $y = \ln(x) + 2$ ,  $y = 2$ , and  $x = 5$ ?

(Use Calculator)

First, draw a sketch

note: a region with boundaries  $y = \ln(x)$ , the x-axis, and  $x = 5$  would have the same area. (a downward shift of 2)


Second, establish boundaries of integral (i.e. interval of the *integrand*)

The right boundary is  $x = 5$

The left boundary is  $x = 1$

because

$$\begin{aligned} y &= \ln(x) + 2 \\ y &= 2 \end{aligned} \quad \begin{aligned} &\text{find intersection:} \\ &\text{(set equations equal)} \\ 2 &= \ln(x) + 2 \\ 0 &= \ln(x) \\ x &= 1 \end{aligned}$$



Then, evaluate the definite integral

$$\int_1^5 \ln(x) + 2 dx - \int_1^5 2 dx = \int_1^5 \ln(x) dx = x \ln(x) - x \Big|_1^5 = 5 \ln(5) - 5 - (0 - 1) \approx 4.047$$

upper boundary                      lower boundary

7) Find the equation of the line that is tangent to  $f(x) = 3x^2 - \ln x$  at  $(1, 3)$

(Optional: Use a graphing calculator to confirm your answer)

Point:  $(1, 3)$


Slope: find instantaneous rate of change (derivative)

$$f'(x) = 6x - \frac{1}{x}$$

then, slope at  $(1, 3)$  is  $f'(1) = 5$

Equation of tangent line:

$$y - 3 = 5(x - 1) \quad \text{or} \quad y = 5x - 2$$




8)  $y = \frac{x^2}{2} - \ln x$

What are the extrema?

Points of inflection?

(Optional: Use a graphing calculator to check your answers)



To find extrema (max. or min), set first derivative equal to zero.

$$y' = x - \frac{1}{x} \quad x - \frac{1}{x} = 0 \quad \text{multiply both sides by } x$$

$$x^2 - 1 = 0 \quad \text{factor}$$

$$(x + 1)(x - 1) = 0$$

$x = -1$  and  $1$       since  $\ln(-1)$  does not exist,  $-1$  is extraneous!

at  $x = 0$ , derivative is  $< 0$  (decreasing)

at  $x = 2$ , derivative is  $> 0$  (increasing)

$x = 1$  is a minimum

$$y'' = 1 + \frac{1}{x^2} \quad 1 + \frac{1}{x^2} = 0$$

$$\frac{1}{x^2} = -1$$

No solution, so there is no point of inflection!

### III. Inverses and derivatives


- 1)  $f(x)$  and  $g(x)$  are one to one inverses.

If the slope of  $f(x)$  at  $(3, 8)$  is 2, where is the slope of  $g(x)$  equal to  $1/2$ ?

Since  $f(x)$  and  $g(x)$  are inverses, they reflect over  $y = x$ .

the rate of change (slopes) of reflection points will be reciprocals...

the reflection point of  $(3, 8)$  is  $(8, 3)$



- 2)  $f(x) = x^3 - x - 6$

What is  $f^{-1}'(0)$ ?

We need to find two parts: 1) the value of  $f^{-1}(0)$  If  $x^3 - x - 6 = 0$   
 $x = 2$

$$f^{-1}'(a) = \frac{1}{f'(f^{-1}(a))}$$

2)  $f'(x) \quad f' = 3x^2 - 1$

$$\frac{1}{3(2)^2 - 1} = \frac{1}{11}$$

- 3)  $h(x) = \ln(x) + 4$

- a) What is the inverse of  $h(x)$ ?

$y = \ln(x) + 4$  "switch x and y"

$x = \ln(y) + 4$

$\ln(y) = x - 4$  "solve for y"

$\log_e(y) = (x - 4)$

$y = e^{x-4}$

$$h^{-1}(x) = e^{x-4}$$

- b)  $h(3) = \ln(3) + 4 \quad (\approx 5.1)$  What is the slope at  $h(3)$ ?

$h'(x) = \frac{1}{x} + 0 \quad h'(3) = \frac{1}{3}$

$h^{-1}(5.1) \approx 3$  What is the slope at  $h^{-1}(5.1)$ ?

$h^{-1}'(x) = e^{x-4} \quad h^{-1}'(5.1) = e^{5.1-4} = e^{1.1} \approx 3$


- c) Graph  $h(x)$  and  $h^{-1}(x)$

Sketch the tangent lines at  $(3, 5.1)$  and  $(5.1, 3)$

What are the equations of the tangent lines?

$y - 5.1 = \frac{1}{3}(x - 3)$  and  $y - 3 = 3(x - 5.1)$

$$y = \frac{1}{3}x + 4.1 \quad y = 3x - 12.3$$



#### IV. Exponential Functions

Find the first derivatives:

1)  $g(x) = 2^{x+3}$

using logarithmic differentiation:

$$y = 2^{x+3}$$

$$\ln y = \ln(2^{x+3})$$

$$\ln y = (x+3)\ln 2$$

$$\frac{1}{y} \cdot y' = (1+0)\ln 2 + 0(x+3)$$

$$y' = y \ln 2$$

$$y' = 2^{x+3} \cdot \ln 2$$

using the definition/formula:

$$u = x + 3$$

$$\frac{du}{dx} = 1 \quad \frac{d}{dx}(2^{x+3}) = (1)(2^{x+3}) \ln 2$$

$$a = 2$$

$$\frac{d}{dx}(a^u) = \frac{du}{dx}(a^u) \ln a$$

4) What is the equation of the line tangent to  $y = 2^{-x}$  at  $(0, 1)$ ?

(optional: sketch a graph containing the function and tangent line)

To find the equation of a line, we need the

point:  $(0, 1)$

slope:  $y' = (-1)(2^{-x})(\ln 2)$

at  $x = 0$ , the slope is  $-(\ln 2) \approx -.693$

$$\text{tangent line: } y = -.693x + 1$$

#### SOLUTIONS

3)  $g(t) = t^2 2^t$

2)  $f(x) = x^2 e^x$

product rule:


$$f'(x) = 2x(e^x) + e^x(x^2)$$


$$u' v + v' u$$

$$= x e^x (x + 2)$$

$$g'(t) = 2t \cdot 2^t + 2^t (\ln 2) \cdot t^2$$

$$= 2^t \left( (\ln 2)t^2 + 2t \right)$$





Implicit Differentiation and Logarithm extras---→

Implicit differentiation and natural log

Example: Find  $\frac{dy}{dx}$  :  $x^2 + 3\ln y + y^2 = 10$

$$2x + 3 \cdot \frac{1}{y} \frac{dy}{dx} + 2y \frac{dy}{dx} = 0$$

$$\frac{3}{y} \frac{dy}{dx} + 2y \frac{dy}{dx} = -2x$$

$$\frac{dy}{dx} \left( \frac{3}{y} + 2y \right) = -2x$$

$$\frac{dy}{dx} = \frac{-2x}{\frac{3 + 2y^2}{y}} = \frac{-2xy}{3 + 2y^2}$$

Example: Find the equation of the line tangent to  $x + y - 1 = \ln(x^2 + y^2)$   
at (1, 0)

To find equation of a line, we need slope and a point.

Point: (1, 0)

Slope: Take the derivative and evaluate the point of tangency

Implicit differentiation:  $1 + (1) \frac{dy}{dx} - 0 = \frac{2x + (2y) \frac{dy}{dx}}{(x^2 + y^2)}$  cross-multiply

$$(x^2 + y^2) + (x^2 + y^2) \frac{dy}{dx} = 2x + (2y) \frac{dy}{dx}$$
 collect dy/dx to one side

$$(x^2 + y^2) - 2x = (2y) \frac{dy}{dx} - (x^2 + y^2) \frac{dy}{dx}$$
 factor out the dy/dx

$$\frac{(x^2 + y^2) - 2x}{(2y) - (x^2 + y^2)} = \frac{dy}{dx}$$


To find slope at point of tangency, substitute (1, 0) for (x, y)

$$\frac{(1+0) - 2}{0 - (1+0)} = 1$$

Equation of the tangent line:  $y - 0 = 1(x - 1)$

or

$$y = x - 1$$





Derivatives of logarithms (other than e)

Example:  $f(x) = 5^x$  find  $f'(x)$

$$\frac{d}{dx} (a^x) = (a^x) \ln a$$

(the base  $a$  is a constant)

using the definition:

$$5^x \cdot \ln 5$$

using logarithmic differentiation:

$$y = 5^x$$

log of both sides

$$\ln y = \ln 5^x$$

logarithm power rule

$$\ln y = x \cdot \ln 5$$

$$\frac{1}{y} \cdot y' = (1)\ln 5 + 0(x) \quad \text{derivative}$$

$$y' = y \ln 5$$

substitution

$$y' = 5^x \cdot \ln 5$$

Example:  $y = 3^{x^2}$  find  $\frac{dy}{dx}$

$$\frac{d}{dx} (a^u) = \frac{du}{dx} (a^u) \ln a$$

(the base  $a$  is a constant  
and  $u$  is a function)

using the definition:

$$u = x^2$$

$$a = 3$$

$$\frac{du}{dx} = 2x$$

$$\frac{dy}{dx} 3^{x^2} = (2x) \cdot 3^{x^2} \cdot \ln 3$$

using logarithmic differentiation:

$$y = 3^{x^2}$$

$$\ln y = \ln 3^{x^2}$$

$$\ln y = x^2 \ln 3$$

$$\ln y = 1.098x^2$$

$$\frac{1}{y} \cdot y' = 2.196x$$

$$\frac{1}{y} \cdot y' = 2x(\ln 3)$$

$$y' = 2.196xy$$

$$y' = 2x(\ln 3)y$$

$$y' = 2.196x (3^{x^2})$$

$$y' = 2x \cdot \ln 3 \cdot 3^{x^2}$$

Example: What is the derivative of  $e^x$  ?

$$y = e^x$$

$$y' = e^x \cdot \ln(e)$$

$$y' = e^x \cdot 1 = e^x$$

## Comparing Logarithmic Differentiation

*Example:*  $y = (2x + 3)^2 (x^2 + 1)$

$$\ln y = \ln \left[ (2x + 3)^2 (x^2 + 1) \right]$$

$$\ln y = \ln (2x + 3)^2 + \ln (x^2 + 1)$$

$$\ln y = 2\ln(2x + 3) + \ln(x^2 + 1)$$

$$\frac{1}{y} \frac{dy}{dx} = 2 \frac{2}{(2x + 3)} + \frac{2x}{(x^2 + 1)}$$

$$\frac{1}{y} \frac{dy}{dx} = \frac{4}{(2x + 3)} + \frac{2x}{(x^2 + 1)}$$

$$\frac{dy}{dx} = \left[ \frac{4}{(2x + 3)} + \frac{2x}{(x^2 + 1)} \right] y$$


$$\frac{dy}{dx} = \left[ \frac{4}{(2x + 3)} + \frac{2x}{(x^2 + 1)} \right] \left[ (2x + 3)^2 (x^2 + 1) \right]$$

$$\begin{matrix} f & g \\ (2x + 3)^2 & (x^2 + 1) \end{matrix}$$

$$2(2x + 3)^1 \cdot 2 \cdot (x^2 + 1) + 2x \cdot (2x + 3)^2$$

$$\begin{matrix} f' & g & g' & f \end{matrix}$$

$$4(2x + 3)(x^2 + 1) + 2x(2x + 3)^2$$



The left uses logarithms and implicit differentiation...

The right uses power rule, product rule, and chain rule...

*Example:*

$$y = \sqrt[4]{\frac{(x - 2)^3 (x^2 + 1)}{(2x + 5)^3}}$$

$$\ln y = \ln \left[ \frac{(x - 2)^3 (x^2 + 1)}{(2x + 5)^3} \right]^{\frac{1}{4}}$$

$$\ln y = \frac{1}{4} \ln \left[ \frac{(x - 2)^3 (x^2 + 1)}{(2x + 5)^3} \right]$$

$$\ln y = \frac{1}{4} \left[ \ln (x - 2)^3 + \ln (x^2 + 1) + \ln (2x + 5)^3 \right]$$

$$\ln y = \frac{1}{4} \left[ 3 \ln (x - 2) + \ln (x^2 + 1) + 3 \ln (2x + 5) \right]$$

$$\frac{1}{y} \frac{dy}{dx} = \frac{1}{4} \left[ \frac{3}{(x - 2)} + \frac{2x}{(x^2 + 1)} + \frac{3 \cdot 2}{(2x + 5)} \right]$$

$$\frac{dy}{dx} = \frac{1}{4} \left[ \frac{3}{(x - 2)} + \frac{2x}{(x^2 + 1)} + \frac{3 \cdot 2}{(2x + 5)} \right] y$$

$$\frac{dy}{dx} = \frac{1}{4} \left[ \frac{3}{(x - 2)} + \frac{2x}{(x^2 + 1)} + \frac{3 \cdot 2}{(2x + 5)} \right] \left[ \frac{(x - 2)^3 (x^2 + 1)}{(2x + 5)^3} \right]^{\frac{1}{4}}$$

Calculus: Logarithm Extras

$$\frac{d}{dx} (a^x) = (a^x) \ln a$$

(the base  $a$  is a constant)

Derivative of an exponential function (other than  $e^x$ )

*Example:* Find the equation of the line tangent to  $y = 4^x$  at  $(1, 4)$

To determine the equation of a line, we need a point and the slope.


Point:  $(1, 4)$

Slope: Find the derivative

$$y' = 4^x \cdot \ln(4)$$

and, at  $x = 1$  the slope is  $4^{(1)} \cdot \ln(4) \approx 5.545$

$$y - 4 = 5.545(x - 1)$$



Logarithmic Differentiation: Using logarithms and implicit differentiation to find a derivative.

*Example:*  $y = x^{\sin x}$  Find  $\frac{dy}{dx}$

Since there is a variable in the term AND the exponent, it cannot be directly differentiated.

But, we can take the natural log of both sides...

$$\ln(y) = \ln(x^{\sin x})$$

Logarithm properties: power rule  $\ln(y) = \sin x \cdot \ln(x)$

Implicit differentiation  $\frac{1}{y} \frac{dy}{dx} = \cos x \cdot \ln(x) + \sin x \cdot \frac{1}{x}$

multiply both sides by  $y$   
(to isolate the  $dy/dx$ )  $\frac{dy}{dx} = y \left( \cos x \ln(x) + \frac{\sin x}{x} \right)$

substitute the  $y$  with the original terms

$$\frac{dy}{dx} = x^{\sin x} \left( \cos x \ln(x) + \frac{\sin x}{x} \right)$$

Example:  $\int \frac{8x+5}{x^2+3x-10} dx$

Using partial fractions to find the integral

$$\frac{A}{(x+5)} + \frac{B}{(x-2)} = \frac{8x+5}{x^2+3x-10}$$

$$\frac{A(x-2)}{(x+5)(x-2)} + \frac{B(x+5)}{(x+5)(x-2)} = \frac{8x+5}{x^2+3x-10}$$

$$A(x-2) + B(x+5) = 8x+5$$

$$Ax - 2A + Bx + 5B = 8x + 5$$

$$(A+B)(x) - 2A + 5B = 8x + 5$$

Then, we know

$$\begin{aligned} A + B &= 8 \\ -2A + 5B &= 5 \end{aligned}$$

$$\begin{aligned} 2A + 2B &= 16 \\ -2A + 5B &= 5 \\ \hline 7B &= 21 \\ B &= 3 \end{aligned}$$

then,  $A = 5$

$$\int \frac{8x+5}{x^2+3x-10} dx = \int \frac{5}{(x+5)} dx + \int \frac{3}{(x-2)} dx$$

$$\frac{5}{(x+5)} + \frac{3}{(x-2)}$$

$$5\ln|x+5| + 3\ln|x-2| + C$$

Example:  $\int \frac{-6x^2+3x+5}{x^3-x} dx$

$$x(x^2-1) = x(x+1)(x-1)$$

$$\frac{-6x^2+3x+5}{x^3-x} = \frac{A}{x} + \frac{B}{(x+1)} + \frac{C}{(x-1)}$$

$$= \frac{(x+1)(x-1)A}{x(x+1)(x-1)} + \frac{x(x-1)B}{x(x+1)(x-1)} + \frac{x(x+1)C}{x(x+1)(x-1)}$$

$$-6x^2+3x+5 = (x+1)(x-1)A + x(x-1)B + x(x+1)C$$

$$Ax^2 - 1A + Bx^2 - Bx + Cx^2 + Cx$$

(regroup the terms)

$$(A+B+C)x^2 + (-B+C)x + A(-1)$$

$$-6x^2 + 3x + 5$$

$$\begin{aligned} A + B + C &= -6 \\ -B + C &= 3 \\ -A &= 5 \end{aligned}$$

$$\begin{aligned} A &= -5 & -5 + B + C &= -6 & -B + C &= 3 \\ & & -B + C &= 3 & & \\ & & \hline & & 2C &= 2 & & \\ & & & & & & B &= -2 \\ & & & & & & C &= 1 \end{aligned}$$

"Express method"

Let  $x = 0$  (to eliminate B and C)

$$0 + 0 + 5 = -1A + 0B + 0C$$

$$A = -5$$

Let  $x = -1$  (to eliminate A and C)

$$-6 + (-3) + 5 = 0A + 2B + 0C$$

$$B = -2$$

Let  $x = 1$  (to eliminate A and B)

$$-6 + 3 + 5 = 0A + 0B + 2C$$

$$C = 1$$

$$\int \frac{-5}{x} + \frac{-2}{(x+1)} + \frac{1}{(x-1)} dx$$


$$-5\ln|x| + -2\ln|x+1| + \ln|x-1| + C$$

$$\frac{-6x^2+3x+5}{x^3-x} = \frac{-5}{x} + \frac{-2}{(x+1)} + \frac{1}{(x-1)}$$

Thanks for visiting. (Hope it helped!)

If you have questions, suggestions, or requests, let us know.

Enjoy



Also, at Facebook, Google+, and TeachersPayTeachers, TES, and Pinterest