


Logarithm and Exponents 2:

Solving equations



(Answer in the back)

Topics include change of base, inverses, inequalities, factoring, intercepts, graphing, and more

Example: $\log_4(x-1) = -1 + \log_4(x)$

$$\log_4(x-1) - \log_4(x) = -1 \quad (\text{Logarithm Quotient Rule})$$

$$\log_4\left(\frac{x-1}{x}\right) = -1 \quad (\text{Change to Exponential Form})$$

$$4^{-1} = \frac{x-1}{x}$$

$$\frac{1}{4} = \frac{x-1}{x} \quad (\text{Cross Multiply})$$

$$4x - 4 = x$$

$$x = \frac{4}{3}$$

Example: $\log_{\sqrt{a}}(5) = \log_a x$ find x:

$$\frac{\log 5}{\log(\sqrt{a})} = \frac{\log x}{\log a}$$

OR

$$\frac{\log 5}{\frac{1}{2} \log a} = \frac{\log x}{\log a}$$

$$\log 5 = \frac{1}{2} \log x$$

$$\log 5 = \log \sqrt{x}$$

$$x = 25$$

$$\log 5 \cdot \log a = \log x \cdot \log \sqrt{a}$$

$$\frac{\log 5}{\log x} = \frac{\log \sqrt{a}}{\log a}$$

$$\log_x 5 = \log_a \sqrt{a}$$

$$\log_x 5 = \frac{1}{2}$$

$$x = 25$$

Example: $\log(\sqrt[3]{x}) = \sqrt{\log(x)}$

$$\log x^{\frac{1}{3}} = \sqrt{\log(x)} \quad (\text{Logarithm Power Rule})$$

$$\frac{1}{3} \log(x) = \sqrt{\log(x)} \quad (\text{Square both sides})$$

$$\frac{1}{3} \log(x) \cdot \frac{1}{3} \log(x) = \log(x)$$

$$\frac{1}{9} (\log(x))^2 - \log(x) = 0$$

$$\log(x) \cdot \left(\frac{1}{9} \log(x) - 1\right) = 0$$

$$\log(x) = 0 \quad \frac{1}{9} \log(x) - 1 = 0$$

$$x = 10^0 \quad \frac{1}{9} \log(x) = 1$$

$$x = 1$$

$$\log(x) = 9$$

$$x = 10^9$$

Example: $e^{3x} = \left(\frac{7}{e}\right)^{x+1}$

$$e^{3x} = (7 \cdot e^{-1})^{x+1}$$

$$e^{3x} = 7^{(x+1)} \cdot e^{-(x+1)}$$

$$\frac{e^{3x}}{e^{-(x+1)}} = 7^{(x+1)}$$

$$e^{4x+1} = 7^{(x+1)}$$

$$\ln e^{4x+1} = \ln 7^{(x+1)}$$

$$(4x+1)\ln e = (x+1)\ln 7$$

$$4x+1 = 1.946x + 1.946$$

$$2.054x = .946$$

$$x = .461 \quad \text{approximately}$$

Check:

$$e^{(3 \cdot .461)} = \left(\frac{7}{e}\right)^{(.461+1)}$$

$$e^{1.383} = \left(\frac{7}{e}\right)^{1.461}$$

$$3.987 = 3.983 \quad (\text{approx.}) \checkmark$$

Example: $\log_7(x+5) = \log_7(x-1) - \log_7(x+1)$

$$\log_7(x+5) = \log_7\left(\frac{x-1}{x+1}\right) \quad (\text{Logarithm Quotient Rule})$$

$$\frac{(x+5)}{1} = \left(\frac{x-1}{x+1}\right) \quad (\text{Drop the logarithms})$$

$$(x+5)(x+1) = x-1$$

$$x^2 + 5x + 6 = 0$$

$$(x+2)(x+3) = 0 \quad x = -2 \text{ or } -3$$

However, logarithms cannot be negative... Therefore there is

NO SOLUTION!

Example: $\log(4x) - \log(24 + \sqrt{x}) = 2$

$$\log_{10} \frac{4x}{(24 + \sqrt{x})} = 2$$

$$\frac{4x}{(24 + \sqrt{x})} = 100$$

$$4x = 2400 + 100\sqrt{x}$$

$$4x - 100\sqrt{x} - 2400 = 0$$

$$x - 25\sqrt{x} - 600 = 0$$

(let $A = \sqrt{x}$)

$$\text{Check: } \log(4(1600)) - \log(24 + \sqrt{1600}) = 2$$

$$\log(6400) - \log(64) = 2 \quad \checkmark$$

$$\log(4(225)) - \log(24 + \sqrt{225}) = 2$$

$$\log(900) - \log(39) = 2 \quad \times$$

$$A^2 - 25A - 600 = 0$$

$$(A-40)(A+15) = 0$$

$$A = 40 \text{ or } -15$$

$$\text{therefore, } \sqrt{x} = 40 \text{ or } -15$$

$$x = 1600 \text{ or } 225$$

Exponents and Logarithms

Example: Find $3^x = 21$

Method 1: Convert to logarithmic form...

$$3^x = 21$$

$$\log_3 21 = x \quad \text{Then, input into a calculator...} \quad x = 2.771244$$

Method 2: Use the common log (base 10)

$$3^x = 21$$

$$\log(3^x) = \log(21)$$

$$x \log(3) = \log(21)$$

"raise" both sides to the common log

logarithm power rule...

$$3^x = 21$$

$$\log_3 21 = x$$

"Change of Base Formula"

$$x = \frac{\log 21}{\log 3} = \frac{1.3222}{.4771}$$


Method 3: Graphing each side

The intersection of

$$y = 3^x$$

and

$$y = 21 \text{ is the solution...}$$



Method 4: Guess and check

$$3^x = 21 \quad \text{If } x = 2, \text{ then } 3^2 = 9 \quad \text{Greater...}$$

$$\text{If } x = 3, \text{ then } 3^3 = 27 \quad \text{Less...}$$

$$\text{If } x = 2.8, \text{ then } 3^{2.8} = 21.67 \quad \text{Less...}$$

$$\text{If } x = 2.7, \text{ then } 3^{2.7} = 19.4 \quad \text{Greater...}$$

$$\text{If } x = 2.75, \text{ then } 3^{2.75} = 20.52 \quad \text{Greater...}$$

We've determined the answer is between 2.75 and 2.8

$$\text{If } x = 2.77, \text{ then } 3^{2.77} = 20.97$$

Example: $36 = 10 \left(1 + \frac{.08}{4}\right)^{4x}$

NOTE: this is a model of a compounding interest function!
 "how long will it take 10 to grow to 36 if compounded at 8% quarterly?"

Rule of 72: $72/8 = 9$
 it will take approx. 9 years to double...
 10... 20 (9 years)... 40 (18 years)
 so, the answer should be a bit under 18 years!

Let's see....

$$\log 3.6 = (4x)\log(1.02)$$

$$.5563025 = (4x)(.00860017)$$

$$x = 16.17 \text{ (approximately)}$$

Example: Find the inverse of $g(x) = 2^{(x-4)} + 6$

Using \log_2 $y = 2^{(x-4)} + 6$ change $g(x)$ to y
 $x = 2^{(y-4)} + 6$ switch x and y
 $x - 6 = 2^{(y-4)}$ solve for y

$$\log_2(x - 6) = \log_2(2^{(y-4)})$$

$$\log_2(x - 6) = (y - 4)\log_2 2$$

$$\log_2(x - 6) = (y - 4)(1)$$

$$\log_2(x - 6) + 4 = y$$

$$g^{-1}(x) = \log_2(x - 6) + 4$$

Using \log $y = 2^{(x-4)} + 6$ change $g(x)$ to y
 (\log_{10}) $x = 2^{(y-4)} + 6$ switch x and y
 $x - 6 = 2^{(y-4)}$ solve for y

$$\log(x - 6) = \log 2^{(y-4)}$$

$$\log(x - 6) = (y - 4)\log 2$$

$$\frac{\log(x - 6)}{\log 2} = (y - 4)$$

$$\log_2(x - 6) = (y - 4)$$

Example: Solve algebraically... Then, support your answer graphically.

$$\log_3 x + 7 = 4 - \log_5 x$$

$$\log_3 x + \log_5 x = 4 - 7$$

$$\frac{\log x}{\log 3} + \frac{\log x}{\log 5} = -3$$

$$.477 \log x + .699 \log x = -3$$

$$2.10 \log x = -3$$

$$\log x = -.85$$

$$x = .141$$

x	y
(.141,	5.22)

To solve on TI-Nspire CX CAS

"solve($\log_3 x + \log_5 x + 3 = 0$, x)"

"enter"



solve graphically on calculator

graph $\log_3 x + 7$: $\frac{\log x}{\log 3} + 7$

then,

graph $4 - \log_5 x$: $4 - \frac{\log x}{\log 5}$

The intersection is the solution!



I. Logarithm rules and properties

Simplify 1) $\ln e^3 + (\ln e)^2 - \ln(4e^2) =$

2) $2\log_4 8 + (\log_3 162 - \log_3 2) =$

Solve for x 3) $\log(x + 3) = \log x + \log 3$

4) $6 + \log(x^2 - 80) = 6$

5) $2 \log_2 x + \log_2 \left(\frac{1}{x-1} \right) = 5$

6) $\log_2(x + 7) - \log_2(x - 7) = 3$

7) $3\log_2 x = -\log_2 27$

8) $\log_3(-81) = x$

II. Exponentials and Bases

Solve for x :

1) $8^{5x} = 16^{3x+4}$

2) $4^{3-x} \cdot \left(\frac{1}{8}\right)^{2x+5} = 16^{x+3}$

3) $2^{x+1} = 3^{x-1}$

4) $2^{x+3} = 3^{2x-1}$

5) $4^{3x+1} = 5^{x-2}$

6) $2^{2/\log_5 x} = \frac{1}{16}$

Solve for x and y :

7) $4^{x+y} = 64$

$2^{2x-y} = 128$

8) $5^{2x+y} = 21$

$7^{4x-y} = 25$

III. Using Change of Base

Simplify:

1) $\log_{10} 11 \cdot \log_{11} 12 \cdot \log_{12} 13 \cdot \dots \cdot \log_{999} 1000 =$

2) $\frac{\log_{25}(3)}{\log_5(81)}$

Solve for x:

3) $\log_4 x + \log_{16} x = 1$

4) $3^{x-9} = \frac{\log_5 8}{\log_5 2}$

Find y:

5) $(\log_3 x)(\log_x 4x)(\log_{4x} y) = \log_x x^2$

6) $\log_9 \left(\frac{1}{27} \right) = \frac{y}{2}$

IV. Factoring exponentials

Solve for x :

1) $2^{2x} - 2^x - 6 = 0$

2) $3^{2x+1} - 7 \cdot 3^x + 2 = 0$

3) $4^x - 2^{x+1} = 3$

4) $e^x - 6e^{-x} = 1$

5) $(\log_3 x)^2 - \log_3(x^2) = 3$

V. Exponential and Logarithm inequalities

1) $\ln(x+2)^2 > 3$

2) $6^{n-1} < 11^n$

3) $\ln(x^2) \geq \ln(x+2)$

4) $2\ln 3 - \ln(x+3) > \ln 6$

5) When is $\log_2(x-2) > \log_4(x)$?

VI. Miscellaneous Questions

- 1) What are the intercepts? (x-intercept and y-intercept)

$$y = \log_3 (x + 9) - 3$$

- 2) The vertical asymptote is at $x = 2$
containing point $(18, -5)$

What is the function in the log form

$$f(x) = \log_4 (x + A) + B ?$$

- 3) $\log_{10} 2 = .30$ What is $\log_3 4$?
 $\log_{10} 3 = .48$ (no calculator)

- 4) Rewrite using base 5:

a) $y = 2(25)^{0.4x}$

b) $y = (4)^{-0.2x}$

5) Find the inverses:

$$f(x) = 4e^{(x+2)} + 16$$

$$h(x) = 3 - \log(2 + x)$$

6) Word Problems

A) You deposit \$10,000 into an investment account that earns 7% interest. How many years will it take to increase to \$30,000?

a) Use the "rule of 72" to get an estimate...

b) Use logarithms to get an actual value....

B) A six year old savings account has \$21,000... It has been compounding interest continuously at 4%.

What was the original savings deposit?

C) If 300 mg of a sample decays to 200 mg in 48 hours, find the half-life of the sample...


***VII. Challenge Questions

1) $3^x \cdot \frac{-4}{3^{x+1}} = 8$

2) $\log_5(x+3) = \log_5(x-1) + \log_3 9 + 6^{\log_6 2}$

3) $2\log_4(x) = \log_4(11x+4) - .5\log_4 9$

4) Graph $\log_3(9x)$ (hint: 9x is "9 times x")



5) $x + 7x^{(2/3)} + 10x^{(1/3)} = 0$



SOLUTIONS-→

I. Logarithm rules and properties

SOLUTIONS

Simplify 1) $\ln e^3 + (\ln e)^2 - \ln(4e^2) =$
 $3\ln e + (1)^2 - (\ln 4 + \ln e^2)$
 $4 - \ln 4 - 2\ln e$
 $2 - \ln 4$

2) $2\log_4 8 + (\log_3 162 - \log_3 2) =$
 $\log_4 8^2 + (\log_3 \frac{162}{2})$
 $3 + 4 = 7$

Solve for x 3) $\log(x + 3) = \log x + \log 3$
 $\log(x + 3) = \log(x \cdot 3)$
 $x + 3 = 3x$
 $3 = 2x$
 $x = 3/2$

4) $6 + \log(x^2 - 80) = 6$
 $\log(x^2 - 80) = 0$
 $10^0 = x^2 - 80$
 $x^2 - 81 = 0$
 $x = 9 \text{ and } -9$

5) $2 \log_2 x + \log_2 \left(\frac{1}{x-1} \right) = 5$

6) $\log_2(x + 7) - \log_2(x - 7) = 3$

logarithm power rule $\log_2 x^2 + \log_2 \left(\frac{1}{x-1} \right) = 5$

$\log_2 \frac{(x+7)}{(x-7)} = 3$

logarithm product rule $\log_2 \left(\frac{x^2}{x-1} \right) = 5$

$2^3 = \frac{(x+7)}{(x-7)}$

change to exponential form $\frac{x^2}{x-1} = 32$

$8x - 56 = x + 7$

cross multiply $x^2 = 32(x - 1)$

$7x = 63$

quadratic formula $x^2 - 32x + 32 = 0$

$x = 9$

$x = 1.033 \text{ or } 30.967$

7) $3\log_2 x = -\log_2 27$

8) $\log_3(-81) = x$

$\log_2 x^3 = \log_2 27^{-1}$

no solution!

$x^3 = \frac{1}{27}$

$3^x \text{ cannot equal } -81$

$x = \frac{1}{3}$

II. Exponentials and Bases

SOLUTIONS

Logarithm 2 Practice Test

Solve for x:

$$1) \quad 8^{5x} = 16^{3x+4}$$

$$(2^3)^{5x} = (2^4)^{3x+4}$$

$$2^{15x} = 2^{12x+16}$$

$$15x = 12x + 16$$

$$3x = 16$$

$$x = 16/3$$

$$4) \quad 2^{x+3} = 3^{2x-1}$$

take the log of both sides:

$$\log 2^{x+3} = \log 3^{2x-1}$$

$$(x+3)\log 2 = (2x-1)\log 3$$

$$.301x + .903 = .653x - .477$$

$$1.380 = .653x$$

$$x = 2.11$$

Solve for x and y:

$$7) \quad 4^{x+y} = 64 \quad 4^{x+y} = 4^3$$

$$2^{2x-y} = 128 \quad 2^{2x-y} = 2^7$$

$$x + y = 3$$

$$2x - y = 7$$

$$x = 10/3$$

$$y = -1/3$$

$$2) \quad 4^{3-x} \cdot \left(\frac{1}{8}\right)^{2x+5} = 16^{x+3}$$

$$\left(2^2\right)^{3-x} \cdot \left(2^{-3}\right)^{2x+5} = \left(2^4\right)^{x+3}$$

$$2^{6-2x} \cdot 2^{-6x-15} = 2^{4x+12}$$

$$2^{-8x-9} = 2^{4x+12}$$

$$-8x-9 = 4x+12$$

$$-21 = 12x$$

$$x = -21/12 = -7/4$$

$$5) \quad 4^{3x+1} = 5^{x-2}$$

one method:
take log (base 4) of both sides...

$$\log_4 4^{3x+1} = \log_4 5^{x-2}$$

$$3x+1 = (x-2)(\log_4 5)$$

$$3x+1 = (x-2)(1.16)$$

$$1.84x = -3.32$$

$$x = -1.80 \text{ approximately}$$

check: $4^{3(-1.80)+1} = 5^{-1.80-2}$
 $.00224 \approx .00221$ ✓

$$3) \quad 2^{x+1} = 3^{x-1}$$

$$\log 2^{x+1} = \log 3^{x-1}$$

$$(x+1)\log 2 = (x-1)\log 3$$

$$(x+1)(.301) = (x-1)(.477)$$

$$.301x + .301 = .477x - .477$$

$$.778 = .176x$$

$$x = 4.42 \text{ (approx.)}$$

Check: $2^{4.42+1} = 3^{4.42-1}$
 $2^{5.42} = 3^{3.42} \text{ (approximately)}$
 $42.81 = 42.82$ ✓

$$6) \quad 2^{2/\log_5 x} = \frac{1}{16}$$

$$2^{2/\log_5 x} = 2^{-4}$$

$$\frac{2}{\log_5 x} = -4$$

$$2 = (-4) \log_5 x$$

$$\frac{-1}{2} = \log_5 x$$

$$x = 5^{-1/2} \text{ or } \frac{1}{\sqrt{5}}$$

$$8) \quad 5^{2x+y} = 21$$

$$7^{4x-y} = 25$$

$$\log_5 (21) = 2x+y$$

$$\log_7 (25) = 4x-y$$

$$1.8917 = 2x+y$$

$$+ 1.6542 = 4x-y$$

$$3.5459 = 6x$$

solve system:
combination/
elimination method

$$x = .591 \quad y = .710$$

check: $5^{2(.591)+(.710)} = 21.01$
 $7^{4(.591)-(.710)} = 24.99$

III. Using Change of Base

SOLUTIONS

Logarithm 2 Practice Test

Simplify:

1) $\log_{10} 11 \cdot \log_{11} 12 \cdot \log_{12} 13 \cdot \dots \cdot \log_{999} 1000 =$

Using change of base formula:

$$\frac{\log 11}{\log 10} \cdot \frac{\log 12}{\log 11} \cdot \frac{\log 13}{\log 12} \dots \frac{\log 999}{\log 998} \cdot \frac{\log 1000}{\log 999}$$

$$\frac{\log 1000}{\log 10} = \frac{3}{1} = 3$$

Solve for x:

3) $\log_4 x + \log_{16} x = 1$

$\log_4 x + \frac{\log_4 x}{\log_4 16} = 1$ use change of base (to base 4)

$\log_4 x + \frac{\log_4 x}{2} = 1$

$2\log_4 x + \log_4 x = 2$ log power rule

$\log_4 x^2 + \log_4 x = 2$ log product rule

$\log_4 x^3 = 2$ convert to exponential form

$x^3 = 16$

$x = 2\sqrt[3]{2}$

Find y:

5) $(\log_3 x)(\log_x 4x)(\log_{4x} y) = \log_x x^2$

Using change of base formula: $\frac{\log x}{\log 3} \cdot \frac{\log 4x}{\log x} \cdot \frac{\log y}{\log 4x} = \log_x x^2$

Simplify: $\frac{\log y}{\log 3} = \log_x x^2$

$\frac{\log y}{\log 3} = 2$

$\log_3 y = 2$

$y = 3^2 = 9$

2) $\frac{\log_{25}(3)}{\log_5(81)}$

change of base $\frac{\frac{\log 3}{\log 25}}{\frac{\log 81}{\log 5}}$

$\frac{\log 3}{\log 25} \cdot \frac{\log 5}{\log 81}$

$\frac{\log 3}{\log 81} \cdot \frac{\log 5}{\log 25}$

$\log_{81}(3) \cdot \log_{25}(5)$

$\frac{1}{4} \cdot \frac{1}{2} = \frac{1}{8}$

4) $3^{x-9} = \frac{\log_5 8}{\log_5 2}$

$\frac{\frac{\log 8}{\log 5}}{\frac{\log 2}{\log 5}} = \frac{\log 8}{\log 2}$

$3^{x-9} = \frac{\log_2 8}{\log_2 2}$

Instead of using base 10, let's use base 2...

$3^{x-9} = \frac{3}{1}$

$3^{x-9} = 3^1$

$x = 10$

6) $\log_9 \left(\frac{1}{27} \right) = \frac{y}{2}$

change of base (to base 3)

$\frac{\log_3 \left(\frac{1}{27} \right)}{\log_3 9} = \frac{y}{2}$

$\frac{-3}{2} = \frac{y}{2}$

$y = -3$

OR change to exponential form

$\frac{y}{9} = \frac{1}{27}$

$(3^2)^{\frac{y}{9}} = \frac{1}{27}$

$3^{\frac{2y}{9}} = 3^{-3}$

$\frac{2y}{9} = -3$

$y = -3$

IV. Factoring exponentials

SOLUTIONS

Solve for x:

1) $2^{2x} - 2^x - 6 = 0$

Hint: $2^{2x} = (2^x)^2$

$$(2^x)^2 - 2^x - 6 = 0 \quad A^2 - A - 6 = 0$$

$$(2^x - 3)(2^x + 2) = 0 \quad (A - 3)(A + 2) = 0$$

$$A = 3, -2$$

$$2^x = 3$$

$$x = \frac{\log 3}{\log 2} \quad \text{approx. 1.585}$$

$$2^x = -2 \quad \text{No solution}$$

3) $4^x - 2^{x+1} = 3$

$$4^x - 2^{x+1} - 3 = 0$$

$$(2^2)^x - (2^x)(2^1) - 3 = 0$$

$$(2^x)^2 - (2^x)(2^1) - 3 = 0$$

Let $y = 2^x$

$$y^2 - 2y - 3 = 0$$

Check:

$$(y - 3)(y + 1) = 0$$

$$y = -1, 3$$

therefore, $2^x = -1$ and 3

approx. 1.585

-1 is extraneous!

$$2^x = 3$$

5) $(\log_3 x)^2 - \log_3 (x^2) = 3$

$$(\log_3 x)^2 - 2(\log_3 x) = 3$$

$$(\log_3 x)^2 - 2(\log_3 x) - 3 = 0$$

$$A^2 - 2A - 3 = 0$$

$$(\log_3 x - 3)(\log_3 x + 1) = 0$$

$$(A - 3)(A + 1) = 0$$

$$(\log_3 x - 3) = 0 \quad \log_3 x = 3 \quad x = 27$$

$$(\log_3 x + 1) = 0 \quad \log_3 x = -1 \quad x = 1/3$$

2) $3^{2x+1} - 7 \cdot 3^x + 2 = 0$

Hint: recognize 3^x as a term and use exponent rules

$$3^{2x+1} = 3^{2x} \cdot 3^1$$

$$3A^2 - 7A + 2 = 0$$

$$\text{Let } A = 3^x$$

$$(3A - 1)(A - 2) = 0$$

$$\text{then, } 3^{2x} = A^2$$

$$A = 1/3 \text{ or } 2$$

$$3^x = 1/3 \quad x = -1$$

$$3^x = 2 \quad x = \frac{\log 2}{\log 3} \quad x = .63 \text{ (approx.)}$$

(substitute into original equation to check!)

4) $e^x - 6e^{-x} = 1$

$$e^x \cdot (e^x - 6e^{-x} - 1) = 0 \cdot e^x$$

$$e^{2x} - 6e^0 - e^x = 0$$

$$e^{2x} - e^x - 6 = 0$$

let $A = e^x$

$$A^2 - A - 6 = 0$$

$$(A - 3)(A + 2) = 0$$

$$A = -2, 3$$

$$e^x = -2 \text{ or } 3$$

-2 is extraneous, because e^x will never be negative.

$$e^x = 3$$

take natural log of each side

$$\ln e^x = \ln 3$$

$$x \ln e = 1.0986 \text{ (approximately)}$$

$$x = 1.0986 \text{ (approximately)}$$

V. Exponential and Logarithm inequalities

SOLUTIONS

1) $\ln(x+2)^2 > 3$

$$\log_e (x+2)^2 = 3$$

$$e^3 = (x+2)^2$$

$$\pm\sqrt{20.08} = x+2$$

$$x = -2 \pm\sqrt{20.08}$$



2) $6^{n-1} < 11^n$

$$(n-1)\log 6 = n\log 11$$

$$n\log 6 - \log 6 = n\log 11$$

$$n\log 6 - n\log 11 = \log 6$$

$$n(\log 6 - \log 11) = \log 6$$

$$n\log(6/11) = \log 6$$

$$n = \log 6 / \log(6/11)$$

$$n > -2.9560$$

If $n = 0$,

then $6^{0-1} < 11^0$

$$\frac{1}{6} < 1 \quad \checkmark$$

3) $\ln(x^2) \geq \ln(x+2)$


$x^2 > (x+2)$ assume terms are equal to determine the 'critical values'

$$x^2 - x - 2 = 0$$

$$(x-2)(x+1) = 0 \quad x = -1, 2$$

$$(-2, -1] \cup [2, \infty)$$

Then, test values in each region...



\ln cannot be 0 or be negative...

4) $2\ln 3 - \ln(x+3) > \ln 6$

$$\ln 9 - \ln(x+3) = \ln 6$$

$$\ln \frac{9}{x+3} = \ln 6$$

$$\frac{9}{x+3} = 6$$

$$6x + 18 = 9$$

$$x = -3/2$$

Test $x = -2$

$$2\ln 3 - \ln(-2+3) > \ln 6 \quad ?$$

$$\ln 9 - \ln(1) > \ln 6$$

$$\ln 9 > \ln 6 \quad \text{YES}$$

Test $x = 0$

$$2\ln 3 + \ln(0+3) > \ln 6 \quad ?$$

$$\ln 9 + \ln 3 > \ln 6$$

$$\ln \frac{9}{3} > \ln 6 \quad \text{NO}$$



So, $x < -3/2$... BUT, it must be greater than -3 (otherwise, $\ln(x+3)$ is undefined)

$$-3 < x < -3/2$$

5) When is $\log_2(x-2) > \log_4(x)$?

(first, find where sides are equal...)

$$\frac{\log_2(x-2)}{\log_2 2} = \frac{\log_2 x}{\log_2 4} \quad \text{use change of base formula}$$

$$\frac{\log_2(x-2)}{1} = \frac{\log_2 x}{2}$$

$$2\log_2(x-2) = \log_2 x$$

$$\log_2(x-2)^2 = \log_2 x$$

$$(x-2)^2 = x$$

$$x^2 - 5x + 4 = 0$$

$$(x-4)(x-1) = 0$$

$$x = 1, 4$$

But, we eliminate 1, because $\log_2(x-2)$ does not exist when $x = 1$

test $x = 3$...

and, the inequality does not work..



$$x \geq 4$$

VI. Miscellaneous Questions

SOLUTIONS

- 1) What are the intercepts? (x-intercept and y-intercept)

$$y = \log_3(x + 9) - 3$$

y-intercept occurs when $x = 0$ (0, ?)

$$(0, -1)$$

$$y = -1$$

x-intercept occurs when $y = 0$ (?, 0)

$$3 = \log_3(x + 9)$$

$$(18, 0)$$

$$x + 9 = 27 \quad x = 18$$

- 2) The vertical asymptote is at $x = 2$
containing point (18, -5)

What is the function in the log form

$$f(x) = \log_4(x + A) + B ?$$

since asymptote is $x = 2$,

$$f(x) = \log_4(x - 2) + B$$

then, to find B, substitute the point (18, -5)

$$-5 = \log_4(18 - 2) + B$$

$$-5 - B = \log_4(16)$$

$$f(x) = \log_4(x - 2) - 7$$

$$B = -7$$

3) $\log_{10} 2 = .30$

$\log_{10} 3 = .48$

What is $\log_3 4$?
(no calculator)

$$\log_3(2)^2$$

$$2 \cdot \log_3(2)$$

$$2 \cdot \frac{\log 2}{\log 3} = 2 \cdot \frac{.30}{.48} = \frac{.60}{.48} = 1.25$$

- 4) Rewrite using base 5:

a) $y = 2(25)^{0.4x}$

$$y = 2(5^2)^{0.4x}$$

Find $5^x = 2$

$$\log 5^x = \log 2$$

$$x = \frac{\log 2}{\log 5} = .43$$

$$y = 5^{(0.8x + .43)} \quad (\text{approx})$$

$$y = 2(5)^{0.8x}$$

$$5^{.43} = 2$$

b) $y = (4)^{-0.2x}$

Find $5^x = 4$

$$\log 5^x = \log 4$$

$$x = \frac{\log 4}{\log 5} = .86$$

$$y = (5)^{-.17x} \quad (\text{approx})$$

$$y = (5^{.86})^{-0.2x}$$

5) Find the inverses:

SOLUTIONS

Logarithm 2 Practice Test

$$f(x) = 4e^{(x+2)} + 16$$

$$y = 4e^{(x+2)} + 16 \quad \text{switch x and y}$$

$$x = 4e^{(y+2)} + 16$$

$$x - 16 = 4e^{(y+2)}$$

$$\frac{x-16}{4} = e^{(y+2)}$$

$$\ln \frac{x-16}{4} = y+2$$

$$f^{-1}(x) = \ln \frac{x-16}{4} - 2$$

$$h(x) = 3 - \log(2+x)$$

$$y = 3 - \log(2+x) \quad \text{switch x and y}$$

$$x = 3 - \log(2+y) \quad \text{solve for y}$$

$$3 - x = \log(2+y)$$

$$10^{3-x} = 2+y$$

$$h^{-1}(x) = 10^{3-x} - 2$$

6) Word Problems

A) You deposit \$10,000 into an investment account that earns 7% interest. How many years will it take to increase to \$30,000?

a) Use the "rule of 72" to get an estimate...

"rule of 72" estimates it'll take 72/7, or approx. 10 years to double..
\$10,000 to \$20,000 will take 10 years...
\$20,000 to \$40,000 will take another 10 years...

b) Use logarithms to get an actual value....

Since we are looking for an estimate for \$30,000, half-way, it takes approx 15 years...

$$A = Pe^{rt} \quad 30,000 = 10,000e^{.07t}$$

$$3 = e^{.07t}$$

$$\ln 3 = \ln e^{.07t}$$

$$t = \frac{\ln 3}{.07} = 15.69 \text{ years (approx)}$$

B) A six year old savings account has \$21,000... It has been compounding interest continuously at 4%.

$$A = Pe^{rt}$$

$$21,000 = Pe^{(.04)(6)}$$

$$21,000 = P(1.27)$$

$$P = 16,519$$

What was the original savings deposit?

C) If 300 mg of a sample decays to 200 mg in 48 hours, find the half-life of the sample...

Step 1: Find the rate r

$$A = Pe^{rt}$$

$$200\text{mg} = 300\text{mg}(e)^{r(48)}$$

$$\frac{2}{3} = e^{48r}$$

$$\ln\left(\frac{2}{3}\right) = 48r(\ln e)$$

$$r = -.008447$$

Step 2: Find the half-life (t)

$$A = Pe^{rt}$$

$$150\text{mg} = 300\text{mg}(e)^{-.008447t}$$

$$\ln\left(\frac{1}{2}\right) = -.008447t$$

$$82 \text{ hours}$$

***VII. Challenge Questions

SOLUTIONS

1) $3^x \cdot \frac{-4}{3^{x+1}} = 8$

$$-4 \cdot \frac{3^x}{3^{x+1}} = 8$$

$$\frac{3^x}{3^{x+1}} = -2$$

NO SOLUTION

$$3^{-1} = -2$$

3) $2\log_4(x) = \log_4(11x + 4) - .5\log_4 9$

$$\log_4(x)^2 = \log_4(11x + 4) - \log_4 9^{.5}$$

$$\log_4(x^2) = \log_4 \frac{(11x + 4)}{3}$$

$$3x^2 = 11x + 4$$

$$3x^2 - 11x - 4 = 0$$

$$(3x + 1)(x - 4) = 0$$

$$x = 4 \text{ or } -1/3$$

ONLY $x = 4$

2) $\log_5(x + 3) = \log_5(x - 1) + \log_3 9 + 6^{\log_6 2}$

$$\log_5(x + 3) - \log_5(x - 1) = 2 + 2$$

$$\log_5 \frac{(x + 3)}{(x - 1)} = 4$$

$$\frac{(x + 3)}{(x - 1)} = 625$$


$$x = \frac{157}{156}$$

$$625x - 625 = x + 3$$

$$624x = 628$$

4) Graph $\log_3(9x)$ (hint: $9x$ is "9 times x ")

$$\log_3(9) + \log_3(x) = 2 + \log_3(x)$$



Points include: (1, 2) (9, 4) and (1/9, 0)

5) $x + 7x^{(2/3)} + 10x^{(1/3)} = 0$

Use substitution

(choose the "smallest variable exponent")

$$\text{Let } U = x^{(1/3)}$$

$$U^3 + 7U^2 + 10U = 0$$

$$U(U + 2)(U + 5) = 0$$

$$U = -2, -5, 0$$

$$U = -2: -2 = x^{(1/3)}$$

$$U = 0: 0 = x^{(1/3)}$$

$$U = -5: -5 = x^{(1/3)}$$

$$x = -8$$

$$x = 0$$

$$x = -125$$

(plug in solutions to original equation to check)

$$-8 + 7(4) + 10(-2) = 0$$


$$0 + 7(0) + 10(0) = 0$$

$$-125 + 7(25) + 10(-5) = 0$$

Thanks for visiting. (Hope it helped!)


If you have questions, suggestions, or requests, let us know.

Cheers



Also, at Teacherspayteachers, Facebook, Google+, TES, and Pinterest

"Use the change of base formula!"



$$\log_2 3 \cdot \log_3 4 \cdot \log_4 5 \cdot \log_5 6 \cdot \log_6 7 \cdot \log_7 8$$

Using Change of Base Formula:

$$\frac{\log 3}{\log 2} \cdot \frac{\log 4}{\log 3} \cdot \frac{\log 5}{\log 4} \cdot \frac{\log 6}{\log 5} \cdot \frac{\log 7}{\log 6} \cdot \frac{\log 8}{\log 7}$$

$$\frac{\log 8}{\log 2} = \log_2 8 = \boxed{3}$$