

Automated Western Blot Processor



How BlotCycler Mini works:

BlotCycler Mini is a compact bench-top device that automates processing of 1 to 8 western blots simultaneously. Blots are placed in specialized trays - choose from delta, mini, midi or mega sizes. Add primary and secondary antibodies, and washing buffers and you're ready to go!

Protocol steps are programmed using the touch screen; set the blocking time, washing steps, addition of antibodies, and incubation time. Each step is performed precisely, from the timing to the management of solutions and

BlotCycler Mini specifications:

Input voltage:	110 – 120V or 220 – 240V
Input voltage:	4A max
Polyurethane trays	5
Delta	4 cm x 10 cm
Mini	10 cm x 8 cm
Midi	15 cm x 10 cm
Mega	30 cm x 10 cm
Dimensions:	44 cm W x 30 cm H x 26 cm D
Weight:	15 lbs.

buffers. The membranes are agitated efficiently and consistently, buffers are thoroughly drained, and antibody is automatically added.

Primary antibodies may be collected and saved for reuse. Once blots are removed, a cleaning cycle prepares the system for the next run. *BlotCycler Mini* works with established protocols, uses standard reagents, requires only about 16" x 10" of bench space and works with standard electrical power. It's built in the USA using patented technology.

- Eliminate variations that occur when protocols are performed manually
- Fits into existing workflow using standard protocols and reagents
- Program up to 100 protocol steps, and save up to 40 protocols in the internal memory
- Performs washing steps automatically after each test
- Delivers worry-free performance using patented fluid control handling

1-440-482-5005

E: order@onsbio.com







Automated Western Blot Processor


Traditional western blot analysis is a time-consuming process. *BlotCycler Mini* changes everything by automating the fluid changes and timing associated with blocking, washing and antibody incubation.


There's no more need to watch over the western blot procedure. With **BlotCycler Mini**, simply set up your protocol, and walk away!

Blot-to-blot consistency. Time after time.

The first lane in each blot shown at left contains a HeLa whole cell lysate, the second lane contains a recombinant human IL-2, and the third lane a mixture of HeLa whole cell lysate and hIL2









RT

The NEW Western Blot Workflow:

- 1. Sample preparation
- 2. Gel electrophoresis
- 3. Electrotransfer (blotting)
- **Blocking and** antibody incubation is performed automatically by BlotCycler Mini
- 5. Quantification

About Precision Biosystems

BlotCycler and **BlotCycler Mini** were developed by Precision Biosystems, a Massachusetts-based company that provides solutions to help researchers automate time-consuming tasks in their laboratories. 082019

Authorized distributor www.onsbio.com

1-440-482-5005

E: order@onsbio.com



www.blotcycler.com

