Jennifer Hoch, DVM Diplomate ACVS



CHECKLIST FOR SURGERY DAY - Orthopedics

- <u>MVSS</u> Surgical Consent Form completed, found at MVSS.info
- $\circ~$ Record vitals on CONSENT form then,
- $\circ \ \ Email \ \ consent \ to \ MVSS for pets @gmail.com$
- Confirm pet is fasted
- Surgical leg painted, shaved or marked with owner
- Surgical leg confirmed, circle: RIGHT or LEFT
- Check skin for pyoderma
- IV catheter placed
- IV fluids. 5-10 ml/kg/hour <u>during surgery</u> if no underlying disease.
- o Anesthesia recommendations
 - o Premedication-
 - Cerenia 1mg/kg IV slow or SQ *Highly recommend for Brachycephalic patient.
 - Choose 1 analgesic
 - Hydromorphone 0.1-<u>0.2</u> mg/kg
 - Morphine 0.25-1.0 mg/kg IM or SQ (canine only)
 - Buprenex 0.02 mg/kg (higher dose for advanced sx)
 - Midazolam or Diazepam 0.1-0.2 mg/kg
 - Other Sedatives IF needed and young/healthy
 - Acepromazine 0.02 mg/kg IV
 - Dexdomitor ¹/₄ IV dose
 - Induction agent: Choose most comfortable protocol
 - Propofol 4-6 mg/kg to effect only
 - Alfaxalone 2mg/kg dog, 5 mg/kg cat to effect only
 - Ketamine/Midazolam or Diazepam
 - \circ Gas anesthesia
 - If possible to bring machine into X-ray room (or have syringe of induction agent available for x-rays)
- Antibiotics
 - Cefazolin 22 mg/kg IV at INDUCTION, then again at 90 min Concentrated 100mg/ml for IV dose (9.6ml Sterile Water)

Jennifer Hoch, DVM Diplomate ACVS



Please have these items out and readily available before we arrive, this helps us to be most efficient and extend our care to more patients per day

- Surgical prep supplies ready at prep area
 - Items to keep patient warm: Socks for feet, towels, hot water bottles. *Most heat is lost during prep time*
 - Designated surgery clippers or Clean/new blade
 - Exam gloves, 1-inch tape roll (sticky tape preferred)
 - Chlorhex scrub and alcohol gauze (non-sterile premade is fine)
 - IV pole to hang the leg, (with nothing hanging or attached)
- **Operating room** supplies ready in OR

*Please be sure to remove clutter and unneeded equipment from the OR

- $\circ~$ Extra Towels (not sterile) 3-4 for floor and positioning
- Surgical gloves, size 6.5 & 7; 2 pair each
- Bottle of 70% Isopropyl Alcohol
- Chlorhexidine scrub <u>**4%**</u>, we will supply if you do not have
- $\circ~$ Saline bottle preferred (or bag of IV fluids) NEW/UNopened
- Surgical Pack (general or spay is fine)
- Sterile bowl w/gauze (can be clean dog bowl sterilized)
- Suture- (generic okay)
 - PDS on taper needle, sizes vary 0, 2-0, or 3-0
 - Monocryl on reverse cutting needle, 3-0 or 4-0
- o Sterile Syringe 12cc; Needles- 21 gauge, 22-20 gauge 1.5in
- \circ Telfa non-adherent pad
- **E-collar** (hard and plastic) to go home
 - $\circ~$ Must be at least 1" past nose when cone pushed down neck
- Medications to go home
 - Antibiotics Cephalexin 30 mg/kg BID for 2 weeks (or Cefpodoxime)
 - $\circ~$ NSAIDs for 2 weeks (if appropriate)
 - Gabapentin 10 mg/kg every 8-12 hours for 7 days
 - Sedative if pet is very excitable or anxious
 - Acepromazine or Trazadone (do not use w/Tramadol)
- Discharge papers and ICEPACK, +/- LICK SLEVE for client
 - MVSS to provide via email in addition to YOUR office discharge