

MOVIPREP

DAY BEFORE COLONOSCOPY OR COLON SURGERY

1. ONE WEEK BEFORE THE EXAMINATION:

- A. STOP all blood thinners (i.e. Coumadin), aspirin and aspirin containing products.
- B. ONE WEEK PRIOR TO EXAM: do not eat products containing Clean or Olestra for example Fat-Free Pringle chips or WOW chips.

2. TWO DAYS BEFORE THE EXAMINATION:

- A. STOP any iron pills and/or fiber supplements such as Citrucel, Metamucil, perdiem or Fibercon.

3. ONE DAY BEFORE THE EXAMINATION:

- A. You must remain on a clear liquid diet the entire day before the examination. You must not eat any solids.

A "Clear Liquid Diet" consists of the following:

- Soft drinks such as ginger ale, cola, sprite, 7-up, strained fruit juices without pulp such as apple juice, white grape juice, lemonade, water, tea or black coffee, NO MILK OR MILK PRODUCTS OR NON-DAIRY CREAMERS, fat free low sodium chicken or beef bouillon/broth.
- Hard candies
- Jell-O (lemon or lime) no fruit or toppings
- Popsicles, Italian ice – no sherbets or fruit bars
- **DO NOT EAT ANYTHING COLORED RED, PURPLE OR ORANGE**

4. ON THE DAY OF EXAMINATION:

- A. Nothing *After Midnight*
- B. Bring someone to drive you home. The procedure will not be done if you do not have a driver

APPOINTMENT INFORMATION

NAME: _____ DATE: _____

LOCATION: _____ REPORT OT ADMITTING @: _____

(If you are using Public Transportation (i.e. KATZ, SOONERRIDE, TAXI SERVICE ETC..) you **must** have a family member or friend with you at your procedure or your examination will be rescheduled **NO EXCEPTIONS**


Patient Instructions

The MoviPrep carton contains 4 pouches and a disposable container for mixing. You must complete the entire prep to ensure the most effective cleansing.

Beginning at: 5 PM


STEP 1 MIX FIRST DOSE

- Empty 1 Pouch A and 1 Pouch B into the disposable container
 - Add lukewarm drinking water to the top line of the container. Mix to dissolve
- If preferred, mix solution ahead of time and refrigerate prior to drinking. The reconstituted solution should be used within 24 hours.*



STEP 2 DRINK FIRST DOSE

- The MoviPrep container is divided by 4 marks. Every 15 minutes, drink the solution down to the next mark (approximately 8 oz), until the full liter is consumed
- Drink 16 oz of the clear liquid of your choice. This is a necessary step to ensure adequate hydration and an effective prep




Clear liquids include water, ginger ale, apple juice, Gatorade®, lemonade, and broth. No red or purple liquids.

Ask your doctor if you have any questions about whether a particular drink is acceptable.

Beginning at: 8 PM


STEP 3 MIX SECOND DOSE

- Empty 1 Pouch A and 1 Pouch B into the disposable container
 - Add lukewarm drinking water to the top line of the container. Mix to dissolve
- If preferred, mix solution ahead of time and refrigerate prior to drinking. The reconstituted solution should be used within 24 hours.*



STEP 4 DRINK SECOND DOSE

- The MoviPrep container is divided by 4 marks. Every 15 minutes, drink the solution down to the next mark (approximately 8 oz), until the full liter is consumed
- Drink 16 oz of the clear liquid of your choice. This is a necessary step to ensure adequate hydration and an effective prep



A colonoscopy prep causes the body to lose a significant amount of fluid and can result in sickness due to dehydration. It's important that you prepare your body by drinking extra clear liquids before the prep. Stay hydrated by drinking all required clear liquids during the prep. Replenish your system by drinking clear liquids after returning home from your colonoscopy.

If you have any questions, please call our office at **918.682.0700**.

*Gatorade is a registered trademark of Stokely-Van Camp, Inc, Chicago, IL.



Web site: www.salix.com 8510 Colonnade Center Drive, Raleigh, NC 27615 Tel • 866.669.SLXP (7597)
©2012 Salix Pharmaceuticals, Inc. All rights reserved. Printed in USA. MOV 11/43-1

MoviPrep®
(PEG-3350, Sodium Sulfate, Sodium Chloride, Potassium Chloride,
Sodium Ascorbate and Ascorbic Acid for Oral Solution)

www.MoviPrep.com

OFFICE INFORMATION

Procedure date:

Arrival time:

Location:

Bowel Preparation for Colonoscopy With MoviPrep

Colonoscopy is a routine, generally safe procedure used to examine the colon. There are steps that you, the patient, must take before the procedure to ensure the highest level of safety and effectiveness.

- ① Make arrangements to have someone drive you home after the procedure
 - Colonoscopy involves sedation, and you will not be allowed to leave unaccompanied
- ② Follow your physician's instructions regarding medicines to avoid and diet to follow before your procedure
- ③ Finish the entire bowel prep regimen as described on the following page, unless otherwise directed by a physician



MoviPrep is available at most retail pharmacy outlets. If your pharmacy does not have MoviPrep in stock, it can be ordered and delivered within 24 hours.

Information for your pharmacist: NDC 65649-201-75

MoviPrep
(PEG-3350, Sodium Sulfate, Sodium Chloride, Potassium Chloride,
Sodium Ascorbate and Ascorbic Acid for Oral Solution)

www.MoviPrep.com

Important Safety Information about MOVIPREP

MOVIPREP® (PEG-3350, sodium sulfate, sodium chloride, potassium chloride, sodium ascorbate and ascorbic acid for oral solution) is an osmotic laxative indicated for cleansing of the colon as a preparation for colonoscopy in adults 18 years of age or older. MOVIPREP is contraindicated in patients with gastrointestinal (GI) obstruction, bowel perforation, gastric retention, ileus, toxic colitis or toxic megacolon, and patients who have had a severe hypersensitivity reaction to any of its components. MOVIPREP should be used with caution in patients at risk of or with fluid and electrolyte abnormalities, hyponatremia, arrhythmias, seizures, in patients with impaired renal function or patients taking concomitant medications that affect renal function, patients with known or suspected inflammatory bowel disease, patients with suspected GI obstruction or perforation, patients at risk for aspiration, and patients with glucose-6-phosphate dehydrogenase deficiency. Most common adverse reactions for split dosing (incidence $\geq 5\%$) are malaise, nausea, abdominal pain, vomiting, and upper abdominal pain. The most common adverse reactions for evening only dosing (incidence $\geq 5\%$) are abdominal distension, anal discomfort, thirst, nausea, abdominal pain, sleep disorder, rigors, hunger, malaise, vomiting, and dizziness. MOVIPREP contains 233 mg of phenylalanine per treatment. Advise patients to hydrate adequately before, during, and after the use of MOVIPREP. Please see accompanying full Prescribing Information for MoviPrep.