

Patient Informed Consent for Botox Cosmetic® Treatment

Patient Label

Dr. Matthew Ward, MD, CCFP and Dr. Galina Turpin, MD, CCFP

Botox® Cosmetic® (Allergan®) is generally a safe, temporary and effective treatment for various cosmetic indications of the upper face. Results take up to 14 days and last 3+ months. Injection intervals should be no more frequent than 3 months with a maximum cumulative dose of 360 units during this time (6U/kg).

As the patient, I have read the following and agree/understand each point (please initial each box):

- I am over the age of 18.
- I have not had any Botox® treatment(s) within the past 3 months.
- I am not pregnant and there is no chance that I could be pregnant at this time. I am not breast-feeding.
- I am not aware of any personal medical history of hypersensitivity or allergic reactions to Botox®.
- I do not have any active skin infections and generally feel well. I am not taking antibiotics.
- I do not take blood thinners (e.g., warfarin, Apixaban/Eliquis, Rivaroxaban/Xarelto, Pradaxa/Dabigatran) or antiplatelet drugs (e.g., Plavix/Clopidogrel, Brilinta/Ticagrelor). Please notify the physician if you take regular “baby aspirin” (ASA 81mg) or regular anti-inflammatory medications for pain (e.g., Advil, Aspirin, Celebrex, Motrin, Aleve, etc...).
- I do not have a known bleeding disorder.
- I do not have a neuromuscular condition (e.g., Myasthenia Gravis, Cerebral Palsy, ALS, previous Polio infection, Lambert-Eaton syndrome, etc.).
- I am aware this product contains human serum albumin which is a derivative of human blood.
- I am aware of the more common, temporary, side effects of Botox® injection(s), including:
 - Inflammation and/or bruising and/or localized pain around the injection site(s)
 - Swelling/edema, bleeding, tenderness, paresthesias (“pins and needles”)
 - Cosmetic asymmetry, not fully-improved “wrinkles,” need for additional treatment(s)
- I am aware of rare, but potentially serious side effects of Botox® treatment, not limited to:
 - Infection
 - Ptosis (drooping eyelid) and/or diplopia (double vision)
 - Extremely rare: distant spread of botulinum toxin resulting in secondary problems such as swallowing or speech disorders, generalized weakness, respiratory failure or complications, cardiovascular events, immunogenic reactions (e.g., anaphylaxis), deleterious neurological effects, and other skin reaction(s).

I am aware that the practice of medicine and surgery is not an exact science and I acknowledge that no guarantees have been given to me concerning the results of any Botox® procedure. Alternative treatment options are also available by a different physician, including: no treatment, filler injections, chemical peels, laser resurfacing, etc. I also give my consent for photographs to be taken as needed before and after the procedure for medical documentation.

Patient Signature

Date